# $PETCT\_NEMA\_IQ.MlAdult\_PET6$

27. februar 2013

# Indhold

1	Fopogram           1.1 Routine	1 1 1
2	Copogram           2.1 Routine            2.2 Scan	1 1 2
3	CT Body 3.1 Routine	2 3 3 4
4	Pause	4
<ul><li>5</li><li>6</li></ul>	PET Control 5.1 Routine	4 5 5 5
7		6 6 6 7
8	Pause	8
9	PET LM2 14  9.1 Routine	8 8 8 8
10	Pause	9

11 PET LM3 15	9
11.1 Routine	. 9
11.2 Scan	. 10
11.3 Recons	. 10
11.3.1 Recon 1	. 10

## 1 Topogram

### 1.1 Routine

• mA: 35

• kV: 120

• Topogram length: 512 mm

• Tube position: Top

### 1.2 Scan

• mA: 35

• kV: 120

• Delay: 4s

• Topogram length: 512 mm

• Direction: Craniocaudal

• Tube position: Top

• API: None

• Kernel: 80s shar

• Window: Topogram Body

## 2 Topogram

### 2.1 Routine

• mA: 35

• kV: 120

• Topogram length: 512 mm

• Tube position: Lateral

• mA: 35

• kV: 120

• Delay: 4s

• Topogram length: 512 mm

• Direction: Caudocranial

• Tube position: Lateral

• API: None

• Kernel: 80s shar

• Window: Topogram Body

# 3 CT Body

### 3.1 Routine

• Eff. mAs: 170

• kV: 120

• CARE Dose4D: On

• CareDoseType: CareDoseAEC

• CTDlvol: 11.4779mGy

 $\bullet$  Scan time: 5.630 s

• Delay: 4.000 s

• Slice: 2.00 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

• Quality ref. mAs: 170

• Eff. mAs: 170

• kV: 120

 $\bullet$  Scan time: 5.630 s

• Rotation time: 0.500 s

• Delay: 4.000 s

• Slice: 2.00 mm

• Pitch: 0.8000000

• Direction: Craniocaudal

### 3.3 Recons

#### 3.3.1 Recon 1

• Series description: AC CT

• Slice: 2.00

• Kernel: B19f PET very smooth

• Window: Abdomen

• Extended FoV: On

• FoV: 700

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 2.000

• No. of images: 111

#### 3.3.2 Recon 2

• Series description: CT Body 2.0 B30f

• Slice: 2.00

• Kernel: B30f medium smooth

• Window: Abdomen

• Extended FoV: Off

• FoV: 500

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 2.000

• No. of images: 111

### 4 Pause

### 5 PET Control

#### 5.1 Routine

• Isotope: F-18

• Pharm.: FDG

- Inj. Dose: 10 Millicuries
- Scan mode: Sinogram
- Scan range: Match CT FOV
- No. of beds: 1
- Scan duration/bed: 60 Seconds

- Autoload: On
- Rebinner LUT: Off
- Scan output: Sinogram
- Sinogram mode: Trues
- Input trigger signal: None
- LLD (keV): 435
- ULD (keV): 650

#### 5.3 Recons

### 5.3.1 Recon 1

- Series description: PET WB
- Recon range (bed): -1 to -1
- Output image type: Corrected
- Recon method: Iterative
- Iterations: 4
- Subsets: 8
- Image size: 168
- Zoom: 1
- Filter: Gaussian

- FWHM (mm): 5
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: On (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

### 6 Pause

### 7 PET LM1 13

### 7.1 Routine

- Isotope: F-18
- Pharm.: FDG
- Inj. Dose: 10 Millicuries
- Scan mode: List mode
- Scan range: Match CT FOV
- No. of beds: 1
- Scan duration/bed: 13 Minutes

#### 7.2 Scan

- Autoload: On
- Rebinner LUT: Off
- Scan output: List mode
- Sinogram mode: Trues
- Input trigger signal: None

- LLD (keV): 435
- ULD (keV): 650

### 7.3 Recons

#### 7.3.1 Recon 1

- Series description: PET WB
- Recon range (bed): -1 to -1
- Output image type: NoRecon
- Recon method: Iterative
- Iterations: 4
- Subsets: 8
- Image size: 168
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 5
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: On (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

### 8 Pause

### 9 PET LM2 14

### 9.1 Routine

• Isotope: F-18

• Pharm.: FDG

• Inj. Dose: 10 Millicuries

• Scan mode: List mode

• Scan range: Match CT FOV

• No. of beds: 1

• Scan duration/bed: 14 Minutes

### 9.2 Scan

• Autoload: On

• Rebinner LUT: Off

• Scan output: List mode

• Sinogram mode: Trues

• Input trigger signal: None

• LLD (keV): 435

• ULD (keV): 650

#### 9.3 Recons

#### 9.3.1 Recon 1

• Series description: PET WB

• Recon range (bed): -1 to -1

• Output image type: NoRecon

- Recon method: Iterative
- Iterations: 4
- Subsets: 8
- Image size: 168
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 5
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: On (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

### 10 Pause

### 11 PET LM3 15

### 11.1 Routine

- Isotope: F-18
- Pharm.: FDG
- Inj. Dose: 10 Millicuries
- Scan mode: List mode
- Scan range: Match CT FOV
- No. of beds: 1
- Scan duration/bed: 15 Minutes

• Autoload: On

• Rebinner LUT: Off

• Scan output: List mode

• Sinogram mode: Trues

• Input trigger signal: None

• LLD (keV): 435

• ULD (keV): 650

#### 11.3 Recons

#### 11.3.1 Recon 1

• Series description: PET WB

• Recon range (bed): -1 to -1

• Output image type: NoRecon

• Recon method: Iterative

• Iterations: 4

• Subsets: 8

• Image size: 168

• Zoom: 1

• Filter: Gaussian

• FWHM (mm): 5

• Offset X: 0 mm

• Offset Y: 0 mm

• Attenuation correction: On (1)

• Scatter correction: On

• Match CT slice location: On

• Save intermediate data: Off