# $4\_PET\_CT\_FDG\_Brain.Adult\_PET4$

23. januar 2013

# Indhold

1	Top	ogram	1		
	1.1	Routine	1		
	1.2	Scan	1		
<b>2</b>	CT Brain				
	2.1	Routine	1		
	2.2	Scan	2		
	2.3	Recons	2		
		2.3.1 Recon 1	2		
		2.3.2 Recon 2	3		
		2.3.3 Recon 3	4		
3	Pau	se ·	4		
4	PE	Brain	4		
	4.1	Routine	4		
	4.2	Scan	5		
	4.3		5		
			5		
			6		

## 1 Topogram

## 1.1 Routine

• mA: 35

• kV: 120

• Topogram length: 256 mm

• Tube position: Lateral

#### 1.2 Scan

• mA: 35

• kV: 120

• Delay: 4s

• Topogram length: 256 mm

• Direction: Craniocaudal

• Tube position: Lateral

• API: API

• Kernel: 20

• Window: Topogram

## 2 CT Brain

## 2.1 Routine

• Eff. mAs: 380

• kV: 120

• CARE Dose4D: On

• CareDoseType: CareDoseAEC

 $\bullet$  CTDlvol: 64.904mGy

• Scan time: 69.330 s

• Delay: 2.000 s

• Slice: 2 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

## 2.2 Scan

• Quality ref. mAs: 380

• Eff. mAs: 380

• kV: 120

 $\bullet$  Scan time: 69.330 s

• Rotation time: 1.000 s

• Delay: 2.000 s

• Slice: 2 mm

• Pitch: 0.55

• Direction: Craniocaudal

### 2.3 Recons

#### 2.3.1 Recon 1

• Series description: tmp CT Brain RTD

 $\bullet$  Slice: 5

• Kernel: H19s

• Window: Cerebrum

• Extended FoV: Off

• FoV: 300

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 5.000

• No. of images: 45

#### 2.3.2 Recon 2

• Series description: AC CT

• Slice: 2

• Kernel: H19s

• Window: Cerebrum

• Extended FoV: Off

• FoV: 300

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 2.000

• No. of images: 112

#### 2.3.3 Recon 3

• Series description: Cranie VIR 2.0 H30S

• Slice: 2

• Kernel: H30s

• Window: Cerebrum

• Extended FoV: Off

• FoV: 300

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 2.000

• No. of images: 112

## 3 Pause

## 4 PET Brain

#### 4.1 Routine

• Isotope: F-18

• Pharm.: Fluorodeoxyglucose

 $\bullet\,$  Inj. Dose: 200000000 Bequerels

• Scan mode: Sinogram

• Scan range: Do not match CT FOV

- No. of beds: Not given. Check recon range
- Scan duration/bed: 10.000000 Minutes

#### 4.2 Scan

- Autoload: On
- Rebinner LUT: Off
- Scan output: Sinogram
- Sinogram mode: false
- Input trigger signal: None
- LLD (keV): n/a
- ULD (keV): n/a

### 4.3 Recons

#### 4.3.1 Recon 1

- Series description: PET Brain Corrected 3 mm zoom 3
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: Iterative3D
- Iterations: 4
- Subsets: 24
- Image size: 400
- Zoom: 3.000000
- Filter: Gaussian
- FWHM (mm): 3.000000
- Offset X: 0 mm
- Offset Y: 0 mm

- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

#### 4.3.2 Recon 2

- Series description: Neurostat
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: Iterative3D
- Iterations: 4
- Subsets: 8
- Image size: 128
- Zoom: 3.000000
- Filter: Gaussian
- FWHM (mm): 8.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off