WB_FDG_PET_barn_4.Child_PET4

30. maj 2013

Indhold

1	Top	Topogram															1						
	1.1	Routir	ne																				1
	1.2	Scan																					1
2	LavDosis CT																						
	2.1	Routin	ne																				1
	2.2	Scan																					1
	2.3	Recon	s																				2
		2.3.1	Recon 1																				2
		2.3.2	Recon 2																				2
		2.3.3	Recon 3																				2
3	Pau	se																					3
4	PET WB															3							
	4.1	Routir	ne																				3
	4.2	Scan																					3
	4.3		S																				3
		4.3.1	Recon 1																				3
		4.3.2	Recon 2																				4

1 Topogram

1.1 Routine

mA: 20kV: 80

Topogram length: 1536 mmTube position: Bottom

1.2 Scan

mA: 20kV: 80Delay: 4s

Topogram length: 1536 mmDirection: CraniocaudalTube position: Bottom

API: APIKernel: 80

• Window: Topogram

2 LavDosis CT

2.1 Routine

• Eff. mAs: 50

• kV: 80

• CARE Dose4D: On

• CareDoseType: CareDoseAEC

CTDlvol: 1.02557mGyScan time: 11.011 s

Delay: 6.000 sSlice: 2 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

2.2 Scan

• Quality ref. mAs: 50

• Eff. mAs: 50

• kV: 80

Scan time: 11.011 sRotation time: 0.500 s

Delay: 6.000 sSlice: 2 mmPitch: 1.20

• Direction: Craniocaudal

2.3 Recons

2.3.1 Recon 1

- Series description: LavDosis CT RTD
- Slice: 4
- Kernel: B19f
- Window: AbdomenExtended FoV: Off
- FoV: 500
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 4.000
- No. of images: 119

2.3.2 Recon 2

- Series description: AC CT
- Slice: 2
- Kernel: B19f
- Window: Abdomen
- Extended FoV: Off
- FoV: 500
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 2.000
- No. of images: 237

2.3.3 Recon 3

- Series description: LavDosis CT 3.0 B30f
- Slice: 3
- Kernel: B30f
- Window: Abdomen
- Extended FoV: Off

- FoV: 500
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 3.000
- No. of images: 158

3 Pause

4 PET WB

4.1 Routine

- Isotope: F-18
- Pharm.: Fluorodeoxyglucose
- Inj. Dose: 0 Bequerels
- Scan mode: Sinogram
- Scan range: Do not match CT FOV
- No. of beds: Not given. Check recon range
- Scan duration/bed: 2.000000 Minutes

4.2 Scan

- Autoload: On
- Rebinner LUT: Off
- Scan output: Sinogram
- Sinogram mode: false
- Input trigger signal: None
- LLD (keV): n/a
- ULD (keV): n/a

4.3 Recons

4.3.1 Recon 1

- Series description: PET WB image check
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: TrueX
- Iterations: 2
- Subsets: 12
- Image size: 200

- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 3.000000
- Offset X: 0 mmOffset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

4.3.2 Recon 2

- Series description: PET WB Corr-TrueX
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: TrueXTOF
- Iterations: 2
- Subsets: 21
- Image size: 400
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 2.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off