
Table of Contents

Example 1 - Load the consolidated sleep diary data (load)	1
Example 2 - Calculate the number of students in section 2 (==)	1
Exercise 1	1
Example 3 - Find the average number of minutes students in section 2 took to fall asleep (indexing)	2
Exercise 2	2
Example 4 - Calculate the number of men in the cohort (strcmp)	3
Exercise 3	3
Example 5 - Calculate the % of men in the cohort	3
Exercise 4	3
Example 6 - Calculate the number of men in section 2 using the And operator (&)	3
Example 7 - Calculate total students in sections 2 or 3 (- OR)	4

Example 1 - Load the consolidated sleep diary data (load)

```
load diaries.mat; % Load the sleep diaries
```

Example 2 - Calculate the number of students in section 2 (==)

```
sect2 = (section == 2); % sect2 has 1's corresponding to section 2 students
totalSect2 = sum(sect2); % Add up the true's (1's) to find number of students
fprintf('%g students in section 2\n', totalSect2);
```

```
46 students in section 2
```

Exercise 1

```
sect1 = (section == 1);
totalSect1 = sum(sect1);
fprintf('%g students in section 1\n', totalSect1);
```

```
45 students in section 1
```

Example 3 - Find the average number of minutes students in section 2 took to fall asleep (indexing)

```
minutesSect2 = toSleepMinutes(:, sect2); % Pick columns of section 2
averMinutes2 = mean(minutesSect2(:)); % Find overall average of section 2
fprintf('Average minutes to sleep for section 2 students = %g \n', ...
averMinutes2);
```

Average minutes to sleep for section 2 students = 16.9482

Exercise 2

```
alarmSect2 = useAlarm(:,sect2);
averAlarmuse2 = mean(alarmSect2(:));
fprintf('Average alarm use suring studying for section 2 students = %g \n',
averAlarmuse2)
```

Average alarm use suring studying for section 2 students = 0.637681

Example 4 - Calculate the number of men in the cohort (strcmp)

```
men = strcmp(gender, 'male'); % 1's in positions corresponding to males
totalMen = sum(men); % Add to find number of men
fprintf('%g men in the cohort\n', totalMen);
```

70 men in the cohort

Exercise 3

```
women = strcmp(gender, 'female');
totalWomen = sum(women);
fprintf('%g women in the cohort\n', totalWomen);
```

74 women in the cohort

Example 5 - Calculate the % of men in the cohort

```
totalStudents = length(gender); % gender has an entry for each student
percentMen = 100.*totalMen./totalStudents;
fprintf('%g%% of the students in the cohort are men\n', percentMen);
```

48.6111% of the students in the cohort are men

Exercise 4

```
percentWomen = 100.*totalWomen./totalStudents;
fprintf('%g%% of the students in the cohort are women\n', percentWomen);
```

51.3889% of the students in the cohort are women

Example 6 - Calculate the number of men in section 2 using the And operator (&)

```
menSect2 = men & sect2; % 1's in positions of men in section 2
totalMen2 = sum(menSect2); % Add up the trues (1's)
fprintf('%g men in section 2\n', totalMen2);
```

25 men in section 2

Example 7 - Calculate total students in sections 2 or 3 (| - OR)

```
sect2or3 = (section == 2) | (section == 3); % 1's for section 2 or 3  
fprintf('%g students in sections 2 or 3\n', sum(sect2or3));
```

```
98 students in sections 2 or 3
```

Published with MATLAB® R2023b