

CPSC 354 Report

Drew Floyd

Contents

1	The MU-Puzzle	2
2	Rewriting Assignment	3

1 The MU-Puzzle

MI \rightarrow MU

Rule 1: If you possess a string whose last letter is I, add U.

Rule 2: Suppose you have Mx, you may add Mxx.

Rule 3: If III occurs in one of the strings, you may make a new string with U in place of III.

Rule 4: If UU, you can drop it.

MI
MII *Mxx*
MIII *Mxx*
MIIIIIIII *Mxx*
MUIIU *MIU*
 \emptyset

MI \rightarrow use *Mxx* rule ∞ times
MIIII...

No matter what Rule you use you will never be able to get 0 Mod3, because I will always be 1 mod 3 or 2 mod 3

MUUU
MIII

Rule 1 does not affect # of I's.

Rule 2 does not give 0 mod 3.

Rule 3 does not solve the problem as removing 3 I's does not change the output of mod3.

Rule 4 does not change the # of I's.

We can never get rid of all of the I's, 0 mod 3 is not possible. Thus you cannot get MU from MI.

2 Rewriting Assignment

1. $A = \{\}$
 $R = \{\}$




This diagram is terminating because there are no infinite loops, confluent because all paths lead to the same result, and has a unique normal form as there is only one final state.

2. $A = \{a\}$
 $R = \{\}$




This diagram is terminating because there are no infinite loops, confluent because all paths lead to the same result, and has a unique normal form as there is only one final state.

3. $A = \{a\}$
 $R = \{(a, a)\}$



This diagram is not terminating due to the presence of infinite loops, confluent because all paths merge, but does not have a unique normal form as multiple results are possible.

4. $A = \{a, b, c\}$
 $R = \{(a, b), (a, b)\}$




This diagram is terminating as there are no infinite loops, not confluent because paths diverge, and does not have a unique normal form due to multiple end states.

5. $A = \{a, b\}$
 $R = \{(a, a), (a, b)\}$




This diagram is not terminating due to the presence of infinite loops, confluent because all paths lead to b, it has a unique normal form as the single end state is b.

6. $A = \{a, b, c\}$
 $R = \{(a, b), (b, b), (a, c)\}$



This diagram is not terminating due to the presence of infinite loops, not confluent because paths diverge, and has a unique normal form due to having a single end state on c.

7. $A = \{a, b, c\}$
 $R = \{(a, b), (b, b), (a, c), (c, c)\}$



This diagram is not terminating due to the presence of infinite loops, not confluent because paths diverge, and does not have a unique normal form due to no end states.

Properties

T	C	U	Example(s)	Explanation
True	True	True	1, 2	These examples terminate, are confluent, and have a unique normal form because all paths lead to a single final state without divergence or loops.
True	True	False	None	This state is impossible because confluence implies a unique normal form when termination is true.
True	False	True	None	This state is impossible because non-confluence means paths diverge, which contradicts having a unique normal form.
True	False	False	4	This example terminates but is not confluent due to diverging paths and does not have a unique normal form as multiple end states exist.
False	True	True	5	This example does not terminate because a is a loop, but since it loops on itself it can still be confluent and point to b to result in a unique normal form.
False	True	False	3	This example does not terminate but is confluent because all paths merge, though it does not have a unique normal form due to multiple results.
False	False	True	None	This state is impossible because non-termination and non-confluence contradict having a unique normal form.
False	False	False	6, 7	These examples do not terminate, are not confluent due to diverging paths, and do not have a unique normal form as no single end state exists.

T: Terminating, C: Confluent, U: Unique Normal Form