

Image Quilting

OVERVIEW

Image Quilting for Texture Synthesis and Transfer

- Our project is based on the paper here (title above):
<http://graphics.cs.cmu.edu/people/efros/research/quilting/quilting.pdf>
- Image Quilting Algorithm*:
 - Go through the image to be synthesized in raster scan order in steps of one block (minus the overlap).
 - For every location, search the input texture for a set of blocks that satisfy the overlap constraints (above and left) within some error tolerance. Randomly pick one such block.
 - Compute the error surface between the newly chosen block and the old blocks at the overlap region. Find the minimum cost path along this surface and make that the boundary of the new block. Paste the block onto the texture. Repeat

** Cited from the above paper*

Minimum Error Boundary Cut

$$E_{i,j} = e_{i,j} + \min(E_{i-1,j-1}, E_{i-1,j}, E_{i-1,j+1})$$

- After DP above, in the end, the minimum value of the last row in E will indicate the end of the minimal vertical path through the surface and we can trace back and find the path of the best cut through the overlapped region.

Texture Transfer

- If we modify the synthesis algorithm by requiring that each patch satisfy a desired correspondence map, C , as well as satisfy the texture synthesis requirements, we can use it for texture transfer.
- The correspondence map is a spatial map of some corresponding quantity over both the texture source image and a controlling target image.
- For texture transfer, image being synthesized must respect two independent constraints:
 - (a) the output are legitimate, synthesized examples of the source texture
 - (b) that the correspondence image mapping is respected.
- Hence, we modify the error term by the use of an 'alpha' parameter.

OUR RESULTS


Texture Synthesis




Texture Synthesis



Texture Synthesis



Texture Synthesis



Texture Synthesis



Texture Synthesis



Texture Synthesis


... of a visual cortical neuron—their
... describing the response of that neuro:
... ht as a function of position—is perhap
... functional description of that neuron.
... seek a single conceptual and mathem:
... scribe the wealth of simple-cell recep
... and neurophysiologically¹⁻³ and inferred
... especially if such a framework has the
... it helps us to understand the functio
... leeper way. Whereas no generic mo
... ussians (DOG), difference of offset C
... rivative of a Gaussian, higher derivati
... function, and so on—can be expect
... simple-cell receptive field, we noneth

... of a visual cortical neuron—function of position—is perhaps (DOG), difference
... describing the response of that neuronal description of that neuron of a Gaussian, high
... ht as a function of position—is single conceptual and mathematical, and so on—can
... functional description of that neuronal wealth of simple-cell receptive function of posit
... seek a single conceptual and neurophysiologically¹⁻³ and inferential description c
... scribe the wealth of simple-cells if such a framework has the a single conceptual
... and neurophysiologically¹⁻³ and in us to understand the function—ie wealth of si
... especially if such a framework way. Whereas no generic model of that physiologically¹
... it helps us to understand the s (DOG), difference of offset and y if such a fram
... leeper way. Whereas no generic of a Gaussian, higher derivat of position—is esta
... ussians (DOG), difference of ion, and so on—can be expectation of that reas
... rivative of a Gaussian, higher derivat of position, higher derivat of position, higher derivat
... ally if such a framework description of that neuronal wealth of simple-cell receptive field, we noneth
... ps us to understand single conceptual and single concept has been expected¹⁻³ and in on—
... way. Whereas no generic model and the function framework has the neuronal concept
... ns (DOG), difference of offset and no generic model and the function and the wealth of
... ve of a Gaussian, higher derivat of offset of offset and no generic model and the wealth of
... h of simple-cell neuronal description of simple-cell receptive field, we noneth
... gically¹⁻³ and infer a single conceptually if we expect to expect, higher derivat of
... a framework describe the wealth ps us to noneth noneth—can be expectation of other
... understand the function neurophysiological way. Whereas no generic Gaussian, higher function
... ereas no generic especially if such ns (DOG), difference of offset on—can be natural de
... difference of offset helps us to understand a framework, higher derivat of function.ingle
... ssian, higher deeper way. Whereas understand the function describe the wealth ns (DOG)
... o on—can be ssians (DOG), difference ereas no generic neurophysiological of physi
... single conceptual and mathematical of that neuronal wealth of offset especially if such a fram
... ne wealth of simple-cell position—is perhaps, higher derivat helps us to understand us to
... physiologically¹⁻³ and in on of that neuron—can be expected way. Whereas noay.
... ly if such a framework derivative of a Gaussian, higher ssians (DOG), difference of
... us to understand the function, and so on—can be derivative of a Gaussian, higher



Texture Synthesis



Texture Synthesis



Texture Synthesis



Texture Synthesis




Texture Synthesis



Texture Transfer




Texture Transfer



Texture Transfer



Texture Transf



THANK YOU!