



Antibiotic resistance in *Pseudomonas aeruginosa* – Mechanisms, epidemiology and evolution

João Botelho^{b,c,*}, Filipa Grosso^a, Luísa Peixe^{a,**}

^a UCIBIO/REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal

^b Max Planck Institute for Evolutionary Biology, 24306 Plön, Germany

^c Evolutionary Ecology and Genetics, CAU Kiel, 24118 Kiel, Germany



ARTICLE INFO

Keywords:

Pseudomonas aeruginosa

Antibiotic resistance

Integrative and conjugative elements

Plasmids

Fitness cost

ABSTRACT

Antibiotics are powerful drugs used in the treatment of bacterial infections. The inappropriate use of these medicines has driven the dissemination of antibiotic resistance (AR) in most bacteria. *Pseudomonas aeruginosa* is an opportunistic pathogen commonly involved in environmental- and difficult-to-treat hospital-acquired infections. This species is frequently resistant to several antibiotics, being in the “critical” category of the WHO's priority pathogens list for research and development of new antibiotics. In addition to a remarkable intrinsic resistance to several antibiotics, *P. aeruginosa* can acquire resistance through chromosomal mutations and acquisition of AR genes. *P. aeruginosa* has one of the largest bacterial genomes and possesses a significant assortment of genes acquired by horizontal gene transfer (HGT), which are frequently localized within integrons and mobile genetic elements (MGEs), such as transposons, insertion sequences, genomic islands, phages, plasmids and integrative and conjugative elements (ICEs). This genomic diversity results in a non-clonal population structure, punctuated by specific clones that are associated with significant morbidity and mortality worldwide, the so-called high-risk clones. Acquisition of MGEs produces a fitness cost in the host, that can be eased over time by compensatory mutations during MGE-host coevolution. Even though plasmids and ICEs are important drivers of AR, the underlying evolutionary traits that promote this dissemination are poorly understood. In this review, we provide a comprehensive description of the main strategies involved in AR in *P. aeruginosa* and the leading drivers of HGT in this species. The most recently developed genomic tools that allowed a better understanding of the features contributing for the success of *P. aeruginosa* are discussed.

1. *Pseudomonas aeruginosa*: the making of a superbug

Pseudomonas aeruginosa is a Gram-negative rod-shaped gamma-proteobacterium. It is a ubiquitous microorganism present in multiple ecological niches, thriving in aquatic and soil habitats due to its metabolic versatility. *P. aeruginosa* can colonize several living sources, such as plants, animals and humans (Gellatly and Hancock, 2013; Moradali et al., 2017; Silby et al., 2011). It is also an opportunistic human pathogen associated with an ever-widening array of life-threatening acute and chronic infections, including cystic fibrosis (CF), ventilator-associated pneumonia, urinary tract infections, otitis externa, burn and wound injuries, bone and joint infections, bacteremia and systemic infections (Shortridge et al., 2019; Bassetti et al., 2018; Burrows, 2018; Nguyen et al., 2018; Parkins et al., 2018). *P. aeruginosa*

is one of the “ESKAPE” pathogens – *Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *P. aeruginosa* and *Enterobacter* species –, emphasizing its impact on hospital infections and the ability of this microorganism to “escape” the activity of antibacterial drugs (Fig. 1) (Boucher et al., 2009). This species is also in the “critical” category of the World Health Organization (WHO)'s priority list of bacterial pathogens for which research and development of new antibiotics is urgently required (Fig. 1) (Tacconelli et al., 2018). Moreover, according to the International Nosocomial Infection Control Consortium report, comprising a surveillance study from January 2010 to December 2015 in 703 intensive care units (ICU) worldwide, nosocomial infections caused by *P. aeruginosa* have become a healthcare concern, mainly due to the high level of resistance to several antibiotics (Rosenthal et al., 2016). The European Centre for Disease Prevention

* Corresponding author at: Department of Evolutionary Ecology and Genetics, Zoological Institute, Christian-Albrechts-Universität zu Kiel, 24118 Kiel, Germany. Antibiotic Resistance Evolution Group, Max-Planck-Institute for Evolutionary Biology, 24306 Plön, Germany.

** Corresponding author at: Laboratório de Microbiologia, Faculdade de Farmácia da Universidade do Porto, Rua Jorge Viterbo Ferreira n° 228, 4050-313 Porto, Portugal.

E-mail addresses: botelho@evolbio.mpg.de (J. Botelho), lpeixe@ff.up.pt (L. Peixe).

<https://doi.org/10.1016/j.drug.2019.07.002>

Received 23 April 2019; Received in revised form 11 July 2019; Accepted 12 July 2019

1368-7646/ © 2019 Elsevier Ltd. All rights reserved.




Fig. 1. Mnemonics assigned to *P. aeruginosa*. WHO classifies *P. aeruginosa* as a “critical” pathogen for which research and development of new antibiotics is urgently required, whereas the US Center for Disease Control flagged multi-drug-resistant *P. aeruginosa* as a serious threat and as one of the ESKAPE pathogens causing nosocomial infections worldwide.

and Control (ECDC) Epidemiological Report 2016 on healthcare-associated infections (HAI) acquired in ICU, comprising data submitted by 15 European Union countries (plus Iceland, Liechtenstein and Norway) in 2014, revealed that *P. aeruginosa* was the most frequently isolated microorganism in ICU-acquired pneumonia episodes and one of the most prevalent in ICU-acquired urinary tract infections and ICU-acquired bloodstream infections (ECDC, 2018a). Point prevalence surveys of HAI and antimicrobial use in European long-term care facilities and acute care hospitals showed that *P. aeruginosa* was one of the microorganisms most frequently isolated from HAI (ECDC, 2013, 2014).

Metabolic versatility, the production of a myriad of virulence factors, the formation of biofilms and antibiotic resistance (AR) comprise the critical traits contributing to the pathogenicity of *P. aeruginosa* (Balasubramanian et al., 2013; Granato et al., 2018; Juan et al., 2017a). The pathogenic potential of this species is governed by the transcriptional, post-transcriptional and post-translational regulation of several systems. An intricate network of transcription factors, quorum-sensing (QS), two-component regulatory systems, non-coding RNAs and unidentified environmental signals controls the expression of these traits (Andersen et al., 2018; Balasubramanian et al., 2014; Chambonnier et al., 2016; Chen et al., 2019; Darch et al., 2018; Ferreira et al., 2019; Koeppen et al., 2016; Kollaran et al., 2019; Lee and Zhang, 2015; Mikkelsen et al., 2011; Nadal Jimenez et al., 2012; Talà et al., 2019). QS is the central regulatory mechanism, controlling biofilm formation, secretion systems, efflux pumps, motility and AR in a population density-dependent manner. The production of several virulence factors, such as alginate, toxins, proteases and siderophores is also regulated by QS. QS controls the expression of multiple virulence factors in a population-dependent manner and is activated by several regulators such as LasR, RhlR, AmpR, MvfR and VqsR (Balasubramanian et al., 2013). Additionally, *P. aeruginosa* is intrinsically resistant to a wide variety of antibiotics (Aleksun and Levy, 2007; Blair et al., 2015; Breidenstein et al., 2011; Burrows, 2018; Chevalier et al., 2017; Du et al., 2018; Hancock, 1998; Juan et al., 2017b; Li et al., 2015; Livermore, 2001; Moradali et al., 2017; Nguyen et al., 2018), and can easily switch from a planktonic to a sessile lifestyle in response to environmental changes, leading to the formation of biofilms, the hallmark of *P. aeruginosa* disease progression and chronic infections (Juan et al., 2017a; Lee and Zhang, 2015; Mikkelsen et al., 2011; Moradali et al., 2017; Schick and Kassen, 2018; Secor et al., 2018).

In this review, we explore the main strategies involved in AR in *P. aeruginosa* and the most important drivers of horizontal gene transfer (HGT) in this species, with emphasis on the barely exploited field analyzing the role of integrative and conjugative elements in the dissemination of carbapenemase-encoding genes. We also sought to provide a brief explanation on the most recently developed genomic tools that allowed a better understanding of the features contributing to the success of *P. aeruginosa*. While this review focuses on *P. aeruginosa*, the concepts discussed herein are also important to other bacteria. Novel therapeutic strategies have been extensively reviewed elsewhere (S.J. Baker et al., 2018; Bassetti et al., 2018; Burrows, 2018; Lynch et al., 2017; Nguyen et al., 2018; Pang et al., 2019).

2. Omics approaches to explore *P. aeruginosa* lifestyle

The genome of *P. aeruginosa* PAO1 reference strain, isolated more than 50 years ago from a patient wound in Melbourne, Australia, was the first to be sequenced, following a community-aided approach that was initiated on 1977 (Stover et al., 2000). The 6.3 mega base pair (Mb) genome comprising 5570 predicted open reading frames was the largest bacterial genome to be sequenced back then. Interestingly, no evidence of recent gene duplication was found, leading to the conclusion that the large genome resulted from genetic and functional diversity. Additionally, a high percentage of predicted regulatory genes (9.4%) was identified in *P. aeruginosa* PAO1 strain and directly linked to the genome size, a trend frequently encountered in bacterial species thriving to survive in several niches (Mathee et al., 2008; Stover et al., 2000). Moreover, about 150 genes identified on *P. aeruginosa* PAO1 strain are predicted to encode for outer membrane proteins involved in adhesion, motility, antibiotic and virulence factors export, an outsized number compared with other genomes (Chevalier et al., 2017). A disproportionately large number of genes (nearly 300) encoding for transport systems and enzymes involved in the uptake of nutrients was also reported (Chevalier et al., 2017; Stover et al., 2000). Genome size, predicted regulatory genes and complexity are consistent with the remarkable versatility frequently described in this species and reflects evolutionary adaptation to multiple distinct niches (Mathee et al., 2008; Stover et al., 2000). Indeed, the expansion on nutritionally restricted environments through acquisition of new metabolic functions most likely comprises a key player shaping the *P. aeruginosa* genome (Mathee et al., 2008; Schick and Kassen, 2018), an evolution pattern contrasting to the observed genome reduction upon adaptation of pathogens to a parasitic existence (Mathee et al., 2008; Moran, 2002; Moran et al., 2009; Weinert and Welch, 2017). A recent genomic and phenotypic analysis of ten different laboratory *P. aeruginosa* PAO1 isolates showed variable production of molecular products, suggesting alterations in transcriptional and translational regulation and the ongoing microevolution during culture and storage that may affect reproducibility when using the reference strain in different laboratories (Chandler et al., 2018).

Comparative genomic analysis performed by Mathee et al. and later revisited, revealed that *P. aeruginosa* genome presents a picture of a mosaic genome, composed of a large number of core genes interspersed by strain-specific blocks of genes (Fischer et al., 2016; Freschi et al., 2018b; Hilker et al., 2015; Klockgether et al., 2011; Kung et al., 2010; Mathee et al., 2008; Valot et al., 2015). The latter segments were defined as regions of genomic plasticity (Mathee et al., 2008). The sum of these unique regions comprise the accessory genome. As such, the *P. aeruginosa* pangenome is currently envisioned as the result of the sum of the (1) core genome, comprising genes that are identified in almost all *P. aeruginosa* strains and the (2) accessory genome, encompassing strain-specific genes usually inserted *en bloc* and clustered in certain loci that may encode niche adaptation properties (Freschi et al., 2018a, 2018b; Kung et al., 2010; Mathee et al., 2008; McInerney et al., 2017;

Valot et al., 2015). Genes are classified as unique in comparative genomics when identified in a single genome. The pangenome of a given species depends on the number of sequenced genomes, and for species with an extended and diverse accessory genome such as *P. aeruginosa*, each new sequenced genome will enlarge this pool of genes and will naturally decrease those of the core genome (Freschi et al., 2018b; Klockgether et al., 2011). Klockgether et al. found that the core genome is largely collinear among *P. aeruginosa* strains and that the accessory genome is the main contributor for genome diversity (Klockgether et al., 2011). Intra-clonal genome diversity of *P. aeruginosa* major clones showed that the core genome was highly conserved, while the accessory genome was equally variable at the intra- and inter-clonal level, suggesting a stronger purifying selection in the core genome than in the accessory genome (Fischer et al., 2016). Comparing with the genome of two major pathogens *Escherichia coli* and *S. aureus*, *P. aeruginosa* displays a larger average size (Boissy et al., 2011; Touchon et al., 2009). Curiously, genome dynamics among these pathogens is remarkably different, reflecting different host-bacteria interactions. Indeed, *E. coli* displays a very large pangenome (17838 genes) and a small core genome (1976 genes), encompassing a variety of commensal and highly pathogenic strains (Touchon et al., 2009). On the other hand, the pangenome of *S. aureus* (3221 genes) is quite close to the average and core genomes size (3118 and 2245 genes, respectively), probably reflecting an optimized capacity to maintain core phenotypes and to survive the human host's conditions (Boissy et al., 2011).

According to the EzBioCloud website (<http://www.ezbiocloud.net/>) (Yoon et al., 2017), the mean values for size, guanine-cytosine (GC) content and number of coding sequences of the 3684 quality controlled *P. aeruginosa* genomes (of which only 165 represent complete genomes) are around 6.6 Mb, 66.2% and 6140, respectively (accessed July 30, 2019). The Pathosystems Resource Integration Center (PATRIC, <https://www.patricbrc.org/>) database harbors a wider collection of *P. aeruginosa* genomes, comprising 4974 whole-genome sequencing (WGS) projects and 422 complete genomes (accessed July 30, 2019) (Wattam et al., 2017). The *Pseudomonas* Genome Database (<http://Pseudomonas.com/>), besides providing significant updates to the *P. aeruginosa* PAO1 genome annotations, also allows large scale comparative curated genome analysis and visualization (Winsor et al., 2016). Another interesting repository of *P. aeruginosa* genomes, the International *Pseudomonas* Consortium Database (<https://ipcd.ibis.ulaval.ca/>), currently contains genomic and phenotypic data regarding more than 1700 isolates from soil, water, plants, animals and human infections (accessed July 30, 2019). The main purpose was to perform a metadata analysis linking bacterial genotype, phenotype and clinical data, as well as to build a user-friendly pipeline for genome evolution, AR and virulence (Freschi et al., 2015). BACTOME (<https://bactome.helmholtz-hzi.de/cgi-bin/h-intro.cgi>) is a recently developed database for the identification of genotype–phenotype correlations of *P. aeruginosa* clinical isolates, to explore sequence variation and the transcriptional landscape of this species (Hornischer et al., 2018).

Apart from genomics, other omics yielded new insights into the organization of the *P. aeruginosa* genome as well as its biology, diversity and virulence (Dötsch et al., 2015; Dunphy et al., 2019; Gao et al., 2018; Gonzalez et al., 2018; Han et al., 2019; Jensen et al., 2017; Rossi et al., 2018; Skurnik et al., 2013; Tognon et al., 2019; Wurtzel et al., 2012). Koeppen et al. used RNA-seq to characterize a novel mechanism whereby *P. aeruginosa* reduces the host immune response through packaged intracellular small RNAs in outer membrane vesicles (Koeppen et al., 2016). A recent transcriptomics study revealed that several AR genes had substantially higher expression in human infections than under laboratory conditions, most likely explaining why AR assays often underestimate resistance in patients (Cornforth et al., 2018). Turner et al. conducted RNA-seq and genome-wide insertion mutant fitness profiling (Tn-seq) analyses to investigate the genetic requirements for acute and chronic pathogenesis in *P. aeruginosa* infections, and demonstrated that the combination of these techniques is

quite helpful for the study of physiology, metabolism and virulence (Turner et al., 2014). In addition, Limoli et al. highlighted the important role of alginate overproduction by mucoid strains of *P. aeruginosa* in reducing the production of anti-*S. aureus* effectors such as siderophores and rhamnolipids, promoting their co-existence in a Model of CF Respiratory Infection (Limoli et al., 2017). Lassek et al. provided a quantitative proteomics analysis from *P. aeruginosa* clinical isolates derived from several infection sites, and hypothesized that common and highly expressed proteins might be considered as promising targets for antimicrobial therapy (Lassek et al., 2016).

3. Antibiotic resistance in *P. aeruginosa*

Antibiotics are powerful medicines used in the treatment of bacterial infections. The inappropriate use of these drugs has promoted the spread of AR in most bacteria (WHO, 2015; Cassini et al., 2018; CDC, 2017; Chatterjee et al., 2018; Davies and Davies, 2010; de Kraker et al., 2016; Núñez-Núñez et al., 2018; Political Declaration, 2016; Temkin et al., 2018; Versporten et al., 2018; Waglechner and Wright, 2017; Tacconelli et al., 2018; O'Neill, 2016; Collignon et al., 2018; European Centre for Disease Prevention and Control, 2018b; ECDC, 2018b; Gillings et al., 2017; Holmes et al., 2016; Klein et al., 2018; Livermore, 2018). The WHO's report on AR provided the first ever global picture and national surveillance capacity with data provided by 114 countries, exposing high levels of AR in all regions of the world and significant gaps in tracking AR (WHO, 2015). The report also lists infection control actions to prevent a post-antibiotic era, in which common infections and minor injuries can once again kill. Maintaining the trend that has been registered throughout the years, the 12th edition of the World Economic Forum's Global Risks Report included rapid and massive spread of infectious diseases, as a result of AR, among the top societal risks confronting the world ("The Global Risks Report, 2017). According to recent estimates, nearly 700,000 people die every year as a consequence of AR infections worldwide (O'Neill, 2016). Additionally, according to the latest ECDC AR surveillance report, the European population-weighted mean for *P. aeruginosa* isolates with combined resistance (resistance to three or more antibiotic groups, including carbapenems) was 12.9%, a steady-state value at least since 2012 (ECDC, 2017). Recently, several approaches have been explored to constrain AR evolution (Durão et al., 2018), such as cellular hysteresis and antibiotic combination efficacy networks (Brochado et al., 2018; Barbosa et al., 2018, 2017; Imamovic et al., 2018; Roemhild et al., 2018), 'evolution-proof' antibiotics (Bell and MacLean, 2018; Ling et al., 2015), inactivation of evolvability factors (Ragheb et al., 2018), preventing the evolution of antibiotic tolerance (Brauner et al., 2016; Levin-Reisman et al., 2017; Meredith et al., 2018) and functional metagenomics combined with next-generation approaches (Crofts et al., 2017).

Eight categories of antimicrobial agents are frequently used to treat *P. aeruginosa* infections: aminoglycosides (gentamicin, amikacin, netilmicin, tobramycin), carbapenems (imipenem, meropenem, doripenem), cephalosporins (ceftazidime, cefepime), fluoroquinolones (ciprofloxacin, levofloxacin), penicillins with β -lactamase inhibitors (ticarcillin-clavulanic acid, piperacillin-tazobactam), monobactams (aztreonam), phosphonic acids (fosfomycin) and polymyxins (colistin and polymyxin B). Ceftazidime-avibactam and ceftolozane-tazobactam were already approved by the FDA and are available for use, while cefiderocol and imipenem-cilastatin/relebactam are currently in development (Barnes et al., 2018; Bassetti et al., 2018; Giani et al., 2018; Nguyen et al., 2018). According to Magiorakos et al., *P. aeruginosa* strains are defined as multidrug-resistant (MDR) if non-susceptible to ≥ 1 agent in ≥ 3 antimicrobial categories, extensively drug-resistant (XDR) if non-susceptible to ≥ 1 agent in all but ≤ 2 categories and pandrug-resistant (PDR) if non-susceptible to all antimicrobial agents listed (Magiorakos et al., 2012).

As stated in the aforementioned reports, MDR bacteria are an increasing healthcare concern worldwide, and many of the involved mechanisms have been unveiled over the last decades. One of the main

drivers for the emergence of MDR in the majority of Gram-negative and Gram-positive bacteria is horizontal gene transfer (HGT). An intricate network of several factors is responsible for the evolution, transmission (vertical and/or horizontal) and maintenance of AR in bacterial populations, such as i) mutation frequency, ii) the resistance level conferred by a given mechanism, iii) fitness cost associated with the burden of AR, iv) epistatic interactions, v) the development of co-, cross- or pleiotropic resistance and vi) the nature of the selective pressure (Baltrus, 2013; Bell and MacLean, 2018; Cantón and Ruiz-Garbajosa, 2011; Estrela and Brown, 2018; Frost et al., 2018; Good et al., 2017; Knopp and Andersson, 2018; Knöppel et al., 2017; Lehtinen et al., 2017; Sánchez-Diener et al., 2017; Schenk et al., 2013; Sommer et al., 2017; Yokoyama et al., 2018). Besides sub-inhibitory concentrations of antibiotics, non-antibiotic pharmaceuticals may also promote AR through HGT (Jutkina et al., 2018; Maier et al., 2018; Wang et al., 2018). Curiously, intrinsic resistance and naturally occurring tools were present in bacteria from all environments (including those without human influence) long before the appearance of the first antibiotics (Box 1) (Clemente et al., 2015; D'Costa et al., 2011; Dantas et al., 2008; Karkman et al., 2018). D'Costa et al. performed metagenomics of ancient environmental DNA from 30,000-year-old Beringian permafrost sediments and identified several genes encoding resistance to antibiotics, providing the first evidence that the presence of genes encoding for AR is indeed an ancient phenomenon that predates our use of antibiotics (D'Costa et al., 2011). These enzymes displayed 53 and 84% identity with known β -lactamases, and are frequently expressed at low levels.

Even though HGT plays a decisive role in the emergence of AR among several species, spontaneous mutation is the main driver in *P. aeruginosa* resistance evolution (Lister et al., 2009). *P. aeruginosa* can develop resistance to a wide range of antibiotics, mainly due to a combination of intrinsic, acquired and/or adaptive resistance (Bassetti et al., 2018; Breidenstein et al., 2011; Chevalier et al., 2017; Juan et al., 2017b; Moradali et al., 2017; Nguyen et al., 2018; Oliver et al., 2015). The main intrinsic resistance feature in *P. aeruginosa* is its low outer membrane permeability, which is ~12–100-fold lower than that reported for *E. coli* (Hancock, 1998). Together with the production of AmpC and MexAB-OprM, these strategies largely contribute to the low basal susceptibility of *P. aeruginosa* to antibiotics. Alongside the genes encoding for AmpC, GyrA/GyrB and ParC/ParE, those encoding for the four clinically relevant AR efflux pumps (MexAB-OprM, MexCD-OprJ, MexXY and MexEF-OprN) were all identified in the core genome (Valot

et al., 2015). Efflux pumps are in fact ancient elements present in bacteria long before the antibiotic era, being responsible for the extrusion of several toxic compounds (Du et al., 2018; Martinez et al., 2009). The constitutive expression of MexAB-OprM is responsible for conferring a basal low susceptibility to nearly all β -lactams (with the exception of imipenem) and fluoroquinolones (Li et al., 2015). Over-expression and/or modification of this pump, alongside deletions of specific chromosomal regions and mutations leading to β -lactam target modification may promote AmpC-independent β -lactam resistance (Cabot et al., 2018). Besides AmpC, *P. aeruginosa* presents other two chromosome-encoded β -lactamase that influence the basal level of susceptibility to β -lactams: the OXA-50 class D β -lactamase and the metallo- β -lactamase PIB-1 (Fajardo et al., 2014; Girlich et al., 2004; Juan et al., 2017b). FosA homologues are present in the majority of genomes in some species, including *P. aeruginosa*, and contribute to intrinsic fosfomycin resistance (Ito et al., 2017b, 2017a).

Adaptive resistance is induced by the presence of antibiotics or other environmental factors, as biocides, pH, anaerobiosis, as well as social bacterial activities for instance swarming motility and biofilm formation (Breidenstein et al., 2011; Moradali et al., 2017; Spalding et al., 2018). These triggers may affect the basal expression level of AmpC and efflux pumps. The inducible expression of AmpC is involved in the intrinsic resistance of this species to aminopenicillins and cephalosporins and in the basal resistance level to imipenem, since these molecules prompt the expression of this β -lactamase (Livermore, 1992). The inducible expression of MexXY confers decreased susceptibility to aminoglycosides (Li et al., 2015). Given the large pool of regulatory genes identified on *P. aeruginosa* genome (Stover et al., 2000), the adaptive phenomenon is expected to play a major role in the outcome of resistance.

Mutational inactivation or loss of outer membrane protein-encoding *oprD* gene plays a key role in imipenem resistance and also reduces susceptibility to meropenem, an antibiotic of the same class (cross-resistance) (Cantón and Ruiz-Garbajosa, 2011; Chevalier et al., 2017; Skurnik et al., 2013). *P. aeruginosa* frequently develops resistance to penicillins, cephalosporins and monobactams due to AmpC over-expression by means of specific mutations in peptidoglycan-recycling genes, such as *dacB*, *ampR*, *ampD* and its homologues *ampDh2* and *ampDh3* (Cabot et al., 2011; Juan et al., 2006; Moya et al., 2009). Furthermore, target mutations in *ampC* were shown to increase resistance to cephalosporins (Berrazeg et al., 2015; Rodríguez-Martínez et al., 2009). Regulation of AmpC (frequently designated as PDC, for

Box 1

Evolution of antimicrobial resistance (AR).

A recent study conducted on the host-associated microbiome of Amerindians with no known exposure to antibiotics found multiple AR genes, revealing that these functional genes may be a natural feature of the human microbiome (Clemente et al., 2015). Additionally, environmental metagenomics studies have identified several AR genes in different niches, suggesting that besides the availability of functional AR genes, the successful transmission of these genes into the clinical setting is also related to their association with mobile genetic elements (MGE), the fitness cost of the gene and the interactions between the donor and recipient niches (D'Costa et al., 2011; Forsberg et al., 2014; Munck et al., 2015; Sommer et al., 2009). An interesting tactic to circumvent this problem was recently proposed by Sommer et al.: functional metagenomics of the human microbiome with which a specific pathogen is most likely to interact should be inspected prior to the development of a new antimicrobial drug (Sommer et al., 2017). With this approach, it would be possible to screen the pool of AR genes (resistome, as coined by D'Costa et al. (D'Costa et al., 2006)) of a given environment and to evaluate the risk assessment of a new antibiotic. Besides intrinsic and acquired resistance genes, the resistome encompasses proto-resistance genes, with the potential to evolve to a resistance function, and cryptic resistance genes, which are not obviously associated with resistance due to low or lack of expression (Crofts et al., 2017).

Back in 1973, the theory that environmental antibiotic-producing bacteria were the evolutionary origin of clinically relevant AR genes was starting to gain attention (Benveniste and Davies, 1973). A recent study by Forsberg et al. developed a functional metagenomics pipeline that unveiled the first evidence of shared multi-drug resistant genes between non-pathogenic soil bacteria and human pathogens (Forsberg et al., 2012). However, resistome composition is correlated with the phylogenetic and taxonomic structure of a microbial community, and the synteny of AR genes with MGE appears to be less frequent in environmental metagenomes than in the genome of human pathogens, suggesting that these genes may not be transferred between environmental bacteria as willingly as between clinically relevant pathogens (Forsberg et al., 2014). Due to the ancient origins of both naturally occurring antibiotics and AR genes, it is not surprising that clinical and agricultural use of antibiotics has potentiated the dissemination of resistance (Knapp et al., 2010). Future studies should prioritize increased surveillance on hotspots for HGT of AR genes, such as the agricultural settings, hospitals, wastewater and sewage settings, through functional metagenomics, next-generation sequencing and computational analyses (Pärnänen et al., 2019; Crofts et al., 2017; Munck et al., 2015).

Pseudomonas-derived cephalosporinase) comprises one of the most intricate repression-derepression systems, since the transcription of *bla_{PD}* is controlled by the aforementioned peptidoglycan-recycling genes (Juan et al., 2006; Rodríguez-Martínez et al., 2009). AmpR is a major transcriptional regulator that also plays a role in switching from the acute to the chronic infection traits, by controlling the expression of virulence factors through QS, biofilm formation and the production of alginate (Balasubramanian et al., 2014; Gifford et al., 2018).

Specific mutations in DNA gyrase and topoisomerase IV-encoding genes *gyrA*, *gyrB*, *parC* and *parE* are responsible for fluoroquinolone resistance in *P. aeruginosa* (Bruchmann et al., 2013). Mutations on efflux pump regulatory genes, such as *nalC* and *nfxB*, may affect antibiotics of the same class or from different classes (pleiotropic resistance), including some β -lactams, fluoroquinolones and aminoglycosides (Cantón and Ruiz-Garbajosa, 2011; Fernández and Hancock, 2012; Livermore, 2001). Evolved colistin resistance in *P. aeruginosa* is a complex and multistep process, that likely emerges through mutation in at least five independent loci with epistatic interactions (Döbelmann et al., 2017; Jochumsen et al., 2016; Lee et al., 2016). Recently, the transferable colistin resistance gene *mcr* was described for the first time in *P. aeruginosa* within a Tn3-like transposon (Snesrud et al., 2018).

4. Carbapenemases identified on *P. aeruginosa*

Due to their high importance for human medicine, carbapenems are considered by the WHO as critically-important antimicrobials that should be reserved for the treatment of infections caused by multidrug resistant Gram-negative bacteria in humans (EFSA, 2013). Recent data from ECDC, presenting the latest information on European antimicrobial consumption, revealed that the population-weighted mean

consumption of carbapenems did not show any significant change during 2011–2015 (ECDC, 2018c). According to the latest ECDC AR surveillance report, the European population-weighted mean for *P. aeruginosa* isolates with resistance to carbapenems was 17.4% (ECDC, 2018c). The latest ECDC epidemiological report on HAI acquired in ICU showed that the percentage of carbapenem-resistant isolates in *P. aeruginosa* was 27.7%, which represents the treatment challenges for ICU patients (European Centre for Disease Prevention and Control, 2018a).

Even though chromosomal mutations (such as those leading to the loss or inactivation of OprD porin and/or to the overexpression of efflux pumps) are still the most important promoters of carbapenem resistance in *P. aeruginosa*, the acquisition of carbapenemase-encoding genes (CEGs) is extremely relevant (Botelho et al., 2015, 2017b, 2018a, 2018b, 2018c; Breidenstein et al., 2011; Diene and Rolain, 2014; Moradali et al., 2017; Oliver et al., 2015; Partridge et al., 2018b; Queenan and Bush, 2007). These β -lactamases are able to hydrolyze carbapenems and confer resistance to virtually all β -lactam antibiotics. A recent survey conducted on 14 European countries revealed that 20% of carbapenem-resistant *P. aeruginosa* isolates were carbapenemase producers (Castanheira et al., 2014). Carbapenemases are not intrinsically expressed by *P. aeruginosa*, but rather encoded by heterologous genes acquired by HGT. Only two resident carbapenemases were identified in *Pseudomonas*: POM-1 from *Pseudomonas otitidis* and PAM-1 from *Pseudomonas alcaligenes* (Suzuki et al., 2014; Thaller et al., 2011). Similar to the majority of the carbapenemases reported among this genus, these resident carbapenemases are metallo- β -lactamases (MBL). These Ambler class B β -lactamases present a pan- β -lactam resistance phenotype with the exception of aztreonam, high carbapenemase activity and resistance to β -lactamase inhibitors (Berglund et al., 2017; Cornaglia et al., 2011; McGeary et al., 2017). Amid this class, several families of enzymes presenting at least 30% of amino acid similarity




Fig. 2. Carbapenemases identified in all *P. aeruginosa* genomes available on NCBI (accessed on the 11/4/2019). 4532 genomes were downloaded and blasted against an in-house database of 688 carbapenemases (adapted from (Botelho et al., 2018c)). 397 carbapenemases were identified and country information was retrieved from the genbank files. When more than one variant was identified (such as in GES, IMP and KPC), the family name was included on the figure legend. VIM-2 was separated from the remaining VIM variants, to highlight its prevalence. The size of the circles reflects the total number of sequenced genomes with the given carbapenemase.

among them were described in *P. aeruginosa*: VIM, IMP, NDM, SPM, DIM, GIM and FIM (Fig. 2) (Berglund et al., 2017; Cornaglia et al., 2011; Nascimento et al., 2016; Oliver et al., 2015; Poirel et al., 2010; Queenan and Bush, 2007; Turton et al., 2015). The *bla*_{VIM}, *bla*_{IMP} and *bla*_{NDM-1} are widely disseminated, opposed to *bla*_{SPM-1}, *bla*_{GIM-1}, *bla*_{FIM-1} and *bla*_{AIM-1}, which appear to be restricted to Brazil, Germany, Italy and Australia, respectively (Cornaglia et al., 2011; Oliver et al., 2015). Even though *bla*_{SPM-1} has been identified in Switzerland and in the United Kingdom (Hopkins et al., 2016; Salabi et al., 2010), both patients had been previously hospitalized in Brazil. The *bla*_{VIM-2} is by far the most frequently reported CEG in *P. aeruginosa* (Botelho et al., 2018c; Cornaglia et al., 2011; Oliver et al., 2015). A few strains were reported to carry more than one carbapenemase (Botelho et al., 2018c). Despite the fact that the origin of these acquired MBL remains unknown, environmental bacteria sharing the same ecologic niche as clinically important pathogens are the most likely source (Cornaglia et al., 2011). Recent approaches to discover MBL inhibitors have been developed (McGeary et al., 2017).

In addition to MBL, serine carbapenemases belonging to Ambler class A β -lactamases, such as KPC-2 and several GES variants, have also been reported in *P. aeruginosa* (Fig. 2) (Botelho et al., 2015, 2018a, 2018c; Dai et al., 2016; Naas et al., 2013; Oliver et al., 2015; Queenan and Bush, 2007). Besides the intrinsic oxacillinase OXA-50, Ambler class D β -lactamases are rarely identified in this species. To date, only three class D carbapenemases have been reported in *P. aeruginosa*: OXA-40, OXA-181 and OXA-198 (El Garch et al., 2011; Evans and Amyes, 2014; Meunier et al., 2016; Sevillano et al., 2009).

5. The role of the *P. aeruginosa* accessory genome in the dissemination of resistance

As aforementioned, the accessory genome is composed of strain-specific genes that may encode niche adaptation properties. In *P. aeruginosa*, the accessory genome comprises integrons, transposons, insertion sequences, genomic islands (GIs), prophages and phage-like elements, plasmids and integrative and conjugative elements (ICEs) (Kung et al., 2010). Larger genomes are more prone to be the recipient of HGT events from divergent organisms, a correlation that might be related to the decrease effect of codon bias with total genomic size (Baltrus, 2013; diCenzo and Finan, 2017; Guglielmini et al., 2011). Since *P. aeruginosa* comprises one of the largest bacterial genomes, it is not surprising to observe the large pool of genes acquired by HGT. Most of these elements are able to mobilize, and frequently possess a mosaic structure comprising modules from different mobile genetic elements (MGEs). Since this species exhibits a high GC content, foreign DNA from other species will be frequently associated with a lower GC content. Other features such as codon and tetranucleotide usage should be analyzed in order to correctly address this distinction (Langille et al., 2010). However, once integrated into the chromosome, genes and elements acquired by HGT will experience the same selective pressure as the core genome and may lose those sequence compositional differences (Kung et al., 2010). For instance, Valot et al. found that recently acquired prophages displayed a lower GC content when compared to long-standing ones, which experienced a genetic drift and reached a similar content as that of the core genome (Valot et al., 2015).

The MGEs frequently integrate into the chromosome by targeting specific regions of genomic plasticity within the core genome. These regions act as hotspots for the insertion of MGEs (Botelho et al., 2018c; Kung et al., 2010; Mathee et al., 2008; Oliveira et al., 2017). These elements may harbor genes i) that improve bacterial persistence within different ecological niches, ii) involved in AR, iii) that encode for virulence determinants, and so their acquisition by HGT may confer advantageous traits and must be seen as a major contributor to shape genome evolution (Kung et al., 2010). However, MGE are essentially self-interested entities, promoting their self-propagation and may cause

a burden to the host (Ghaly and Gillings, 2018; Rankin et al., 2011). Taking into account the significant value that mutational events already represent for AR in *P. aeruginosa*, the spread of MGE harboring AR genes illustrates the dire public health consequences of such events.

6. Integrons and “jumping” genes

Most CEGs identified in *P. aeruginosa* are associated with class I integrons (Oliver et al., 2015; Partridge et al., 2018b). Integrons are genetic elements composed of an integrase-encoding *intI* gene, an *attI* recombination site and a Pc promoter (Domingues et al., 2012; Partridge et al., 2018b). These elements present a variable region flanked by two conserved regions (5'CS and 3'CS). The integrase mediates site-specific recombination responsible for the acquisition or excision of gene cassettes harboring AR genes. Several gene cassettes may be captured by the same integron into the variable region and their expression will be further regulated by the Pc promoter (Domingues et al., 2012; Partridge, 2011; Partridge et al., 2018b). Integrons *per se* are not mobilizable, but these elements may be inserted on MGEs such as transposons, plasmids and/or ICEs (Kung et al., 2010; Partridge et al., 2018b). Also, these elements may be mobilized when a complete transposition module is provided in *trans* (Domingues et al., 2012). Related integrons may be identified in different contexts, which also suggest an important contribution of homologous recombination in promoting the evolution of integrons (Domingues et al., 2012; Partridge et al., 2018b).

The most clinically relevant class I integrons identified are ‘sul1-type’ integrons (Domingues et al., 2012; Partridge, 2011; Partridge et al., 2018b). These elements present a *qac* Δ *1**sul1* gene fusion, exposing a successful link between an antiseptic and a sulphonamide resistance gene and its further association with several AR genes, creating a so-called winner effect (Baquero, 2004; Domingues et al., 2012; Partridge, 2011). Class I integrons are derived from different lineages, comprising variable ISs. Among these, *Int4*-like integrons are particularly relevant in *P. aeruginosa*. These elements present a partially duplicated IS6100 insertion sequence downstream the 3'CS region. Variants lacking part or all of the 3'CS region, almost certainly resulting from IS6100-mediated deletions into the conserved region, were also identified (Partridge, 2011; Partridge et al., 2001). Surprisingly, most of these integrons are associated with a Tn402-like transposon which lost part of its transposition module (Domingues et al., 2012; Partridge, 2011; Partridge et al., 2018b). These defective transposons may however be mobilized in *trans* and are capable of targeting the *res* site of Tn3-like transposons and plasmids, establishing a Matryoshka-like arrangement that may contribute for their mobilization (Nicolas et al., 2015; Partridge, 2011; Partridge et al., 2018b; Petrovski et al., 2011). The Tn3 family of transposons are flanked by 38-bp inverted repeats and comprise genes encoding for a transposase, resolvase and a resolution site. Tn3-type transposons are frequently associated with mercury resistance genes. Its dissemination throughout different bacterial species provided the bedrock for the emergence of class I integrons harboring AR genes (Nicolas et al., 2015; Partridge, 2011; Partridge et al., 2018b; Petrovski et al., 2011).

In *P. aeruginosa*, most acquired carbapenemases are present on class I integrons within Tn402-like transposons of chromosomal location, with the important exceptions of *bla*_{SPM-1}, *bla*_{NDM-1} and *bla*_{KPC-2} (Botelho et al., 2018c; Dai et al., 2016; Ding et al., 2018; Naas et al., 2013; Nascimento et al., 2016; Oliver et al., 2015; Queenan and Bush, 2007). Importantly, most integrons bearing CEGs co-harbor gene cassettes mediating resistance to other antibiotic classes (Botelho et al., 2018c; Cornaglia et al., 2011; Oliver et al., 2015; Partridge et al., 2018b). Dissemination of these integrons may then contribute to the occurrence of MDR phenotypes. Besides CEGs, integrons often harbor genes encoding extended-spectrum β -lactamases (ESBLs) mediating resistance to penicillins, third-generation cephalosporins and aztreonam (Bouheraoua et al., 2018; Garza-Ramos et al., 2015; Laudy

et al., 2017; Picao et al., 2009). Recently, aminoglycoside-modifying enzymes (AMEs, conferring selective resistance) and 16S rRNA methyltransferases (*rmt*, mediating class-wide resistance) were found to be prevalent among this species and other clinically relevant Gram-negative pathogens (Costello et al., 2018; Jackson et al., 2013). *rmt* genes were often located within class I integrons (Silveira et al., 2016; Tada et al., 2017). A quinolone resistance determinant *qnrVC1* was identified within a class I integron co-harboring *bla_{VIM-2}* and genes conferring resistance to other antibiotics (Belotti et al., 2015). This was the first description of a *qnr* determinant in a *P. aeruginosa* strain.

Several carbapenemase-harboring transposons have been identified in *P. aeruginosa*, such as KPC-2-encoding Tn4401, a Tn3-like transposon assuring high-frequency transposition; *bla_{VIM}*-harboring Tn6001, Tn6162, Tn6249, Tn6352 and Tn6356 transposons; a Tn5051-type transposon housing a *bla_{IMP-13}* (Botelho et al., 2017a, 2017b; Chowdhury et al., 2017; Cuzon et al., 2011; Di Pilato et al., 2015; Naas et al., 2013; Papagiannitsis et al., 2017; Pfennigwerth et al., 2017; Roy Chowdhury et al., 2009; Samuelsen et al., 2009; Shimizu et al., 2015; Toleman et al., 2003; Tseng et al., 2009; van der Zee et al., 2018; Zhan et al., 2018). Besides CEGs, transposons may carry genes conferring resistance to other antibiotics (Dubois et al., 2002; Feng et al., 2016; Subedi et al., 2018; Uemura et al., 2010). A transferable colistin resistance gene (*mcr-5*, and reclassified as *mcr-5.1* according to the nomenclature proposed by Partridge et al. (Partridge et al., 2018a)) was recently reported within a Tn3-like transposon in a *P. aeruginosa* clinical isolate (Snesrud et al., 2018). This was the first report of *mcr* in colistin-susceptible *P. aeruginosa*.

Recently, a transposable element typing pipeline named TETyper (<https://github.com/aesheppard/TETyper>) was developed (Sheppard et al., 2018). This tool classifies variation and genetic contexts of transposable elements from short-read WGS data. Accurate annotation of integrons and transposable elements may be achieved by Galileo™ AMR (<https://galileoamr.arcbio.com/mara/>) (Arc Bio, Cambridge, MA) (Partridge and Tsafnat, 2018). The INTEGRALL database (<http://integrall.bio.ua.pt/>) provides an easy access to the integron DNA sequences and genetic arrangements (Moura et al., 2009). IntegronFinder (https://github.com/gem-pasteur/Integron_Finder) is a bioinformatics tool to find integrons in bacterial genomes (Cury et al., 2016).

7. Plasmids

Transposons or “jumping genes” are MGEs which integrate into and excise from the chromosome but are not capable of being horizontally transferred (Roberts et al., 2008; Roberts and Mullany, 2011). To move between cells, transposons are frequently encountered within plasmids. These autonomous self-replicating elements may be transferred from cell to cell by conjugation (Garcillán-Barcia et al., 2011; Rozwandowicz et al., 2018; San Millan, 2018; San Millan and MacLean, 2017; Smillie

et al., 2010). Halary et al. reconstructed a network of MGEs and concluded that conjugation is more frequent than transduction, a bacteriophage-mediated DNA transfer mode, emphasizing the dominant role of plasmids in HGT (Halary et al., 2010; Lopatkin et al., 2017; Ramsay and Firth, 2017). The plasmid backbone is mainly taken up by genes encoding for its own propagation and stability (Hülter et al., 2017). The replication module defines the copy number of plasmid cells and its own survival in several hosts (Ilhan et al., 2018). Low copy-number plasmids are subjected to a higher frequency of plasmid loss, due to random assortment at cell division (Garcillán-Barcia et al., 2011; Smillie et al., 2010). As so, extra stability modules such as toxin-antitoxin and partition systems may be required (Baxter and Funnell, 2014; Díaz-Orejas et al., 2017).

Mobility of plasmids will depend on the set of backbone genes, and so these extrachromosomal elements may be conjugative, mobilizable or non-transmissible. Conjugative plasmids carry all the machinery necessary for self-propagation: (i) a relaxase, a key protein in conjugation; (ii) an origin of transfer site, a small DNA sequence that is recognized by the relaxase and that is required in *cis* for a plasmid to be properly transferred; (iii) a set of genes encoding for the type-IV secretion system (T4SS), which provides a membrane-spanning secretion channel connecting the donor to the recipient strain; and (iv) a type-IV coupling protein-encoding gene, associated with the link between the relaxosome and the mating channel (Garcillán-Barcia et al., 2011; Smillie et al., 2010). Mobilizable plasmids can spare the complete set of genes encoding for the T4SS, and may use those of a helper plasmid present in the cell to be successfully transferred. Conjugative plasmids tend to be low copy number and large-sized, while mobilizable plasmids are frequently high copy number and low-sized (< 30 kb) (Garcillán-Barcia et al., 2011; Smillie et al., 2010). Interestingly, Smillie et al. found that more than half of the plasmids are non-transmissible (Smillie et al., 2010), and these elements may in fact use diverse strategies to facilitate their mobilization (Peña-Miller et al., 2015; Ramsay and Firth, 2017; Wein et al., 2019).

Plasmids may harbor an accessory module (comprising for example virulence-encoding factors and/or AR genes) that confers an adaptive advantage upon their host (Loftie-Eaton et al., 2017; Rozwandowicz et al., 2018; San Millan, 2018; Svara and Rankin, 2011). Table 1 summarizes the main characteristics of the CEG-bearing plasmids that have been reported in *P. aeruginosa*. Besides CEGs, plasmids may carry genes conferring resistance to other antibiotics (Chávez-Jacobo et al., 2018; Jeannot et al., 2018).

To date, fourteen incompatibility groups (IncP-1 to IncP-14) were characterized amongst *Pseudomonas* plasmids (Partridge et al., 2018b; Thomas and Haines, 2004). Narrow host range plasmids comprise IncP types 2, 5, 7, 10, 12 and 13 and cannot be transferred onto *E. coli*. On the other hand, other groups seem to display a broad host range, being also included in the plasmid type scheme of *Enterobacteriaceae*: IncP-1

Table 1

List of relevant CEG-harboring plasmids in *P. aeruginosa* deposited on the National Center for Biotechnology Information (NCBI).

Plasmid	Carbapenemase	Inc type(s)	Size(s) (kb)	GC content (%)	Country	Origin	Year	References
pAMBL1 and pAMBL2	VIM-1	–	26 and 24	63.5 and 60.4	Spain	Clinical	2006–07	(San Millan et al., 2015a)
pNOR2000	VIM-2	–	22	62.8	France	Clinical	1996	(Bonnin et al., 2013)
pDCPR1	VIM-2	Unknown	18	58.4	Argentina	Clinical	2005	(Vilacoba et al., 2014)
pJB12	VIM-2	Unknown	30	62.6	Portugal	Clinical	2006	(Botelho et al., 2017a)
pJB37	VIM-2	IncP-2	465	57.2	Portugal	Clinical	2008	(Botelho et al., 2017b)
pS04 90	VIM-2	Unknown	159	57.7	Netherlands	Clinical	–	(van der Zee et al., 2018)
p07-406	VIM-7	–	24	63.8	USA	Clinical	2001	(Li et al., 2008)
pP378-IMP	IMP-4	IncN	51	50.5	China	Clinical	2009–2013	(Feng et al., 2016)
pOZ176	IMP-9	IncP-2	501	57.0	China	Clinical	2000	(Xiong et al., 2013)
pBM413	IMP-45	Unknown	423	56.4	China	Clinical	2012	(J. Liu et al., 2018)
pHN39-SIM	SIM-2	Unknown	282	56.9	China	Clinical	2012	(Sun et al., 2016)
pCOL-1 and pPA-2	KPC-2	IncP-6 and IncU	32 and 8	60.0 and 56.0	Colombia	Clinical	–	(Naas et al., 2013)
p10265-KPC	KPC-2	IncP-6	39	58.2	China	Clinical	2010	(Dai et al., 2016)
pOXA-198	OXA-198	IncP-11	49	60.5	Belgium	Clinical	2010 and 2013	(Bonnin et al., 2018)

(IncP), IncP-3 (IncA/C), IncP-4 (IncQ), IncP-6 (IncG) (Partridge et al., 2018b; Thomas and Haines, 2004). Contrarily to what occurred with *Enterobacteriaceae*, a replicon-based PCR for *P. aeruginosa* plasmids has not been created yet. Moreover, plasmid typing among *P. aeruginosa* is particularly challenging due to insertion, deletion, co-integration and exchange events that will determine low phylogenetic concordance between plasmid core genes (Orlek et al., 2017; Tazzyman and Bonhoeffer, 2014). The advent of WGS enabled the *in silico* analyses of a wide array of plasmids, most of them provided by assembly of short-read sequencing data (see References on Table 1). However, plasmid reconstruction can be quite puzzling (Arredondo-Alonso et al., 2017). Performing plasmid isolation before sequencing may optimize this task, but is laborious. The most appropriate solution is to undergo long-read sequencing (such as Pacific Biosciences or Oxford Nanopore MinION), which allows a more accurate reconstruction. Several *in silico* platforms are available to analyze plasmids: PLACNET (<https://sourceforge.net/projects/placnet/>), a reference-dependent graph-based tool and its recently developed user-friendly web-tool (<https://castillo.dicom.unican.es/upload/>) (Lanza et al., 2014; Vielva et al., 2017); the automated and reference-independent tools plasmidSPAdes (<http://cab.spbu.ru/software/spades/>) and Recycler (<https://github.com/Shamir-Lab/Recycler>) (Antipov et al., 2016; Rozov et al., 2016); PlasFlow (<https://github.com/smaegol/PlasFlow>), that employs a neural network approach to identify plasmid sequences in metagenomics data (Krawczyk et al., 2018); PLSDB (<https://ccb-microbe.cs.uni-saarland.de/plsdb/>), a complete bacterial plasmids database (Galata et al., 2018); Plasmid Atlas (<https://github.com/tiagofilipe12/pATLAS>), a web interface to browse for plasmids and their associated genes (Jesus et al., 2019) and PlasmidSeeker (<https://github.com/bioinfo-ut/PlasmidSeeker>), a k-mer based tool for the identification of known plasmids from sequencing reads (Roosaare et al., 2018). Future advances in plasmid metagenomics and optical mapping of intact plasmids combined with sequencing-based analysis will enhance our knowledge of the plasmidome present in different environments and to improve plasmid assembly, respectively (Orlek et al., 2017; Bikkarolla et al., 2019).

Reducing antibiotic use alone is insufficient to tackle AR since plasmids are often transferred at high rates in the absence of antibiotic selection (Lopatkin et al., 2017). Recently, novel strategies combining conjugation inhibition and promoting plasmid loss to reduce the global burden of AR were proposed (Buckner et al., 2018).

8. Genomic islands and integrative and conjugative elements

GIs comprise a cluster of genes that had been acquired by a bacterial genome by HGT and contributed to diversification and host adaptation (Bellanger et al., 2014; Kung et al., 2010; Langille et al., 2010). These broad definition may also comprise other MGEs, such as ICEs and prophages (Langille et al., 2010). GIs frequently display a sporadic distribution along the genome, a large size, an association with phage integrase genes and a sequence composition that is significantly distinct from that of the host. These elements also tend to target tRNA genes and are normally flanked by direct repeats (Bellanger et al., 2014; Kung et al., 2010; Langille et al., 2010). However, these features are not specific to these elements, which makes GIs prediction a difficult task. In fact, highly expressed genes frequently present a different sequence composition than the rest of the genome, yielding false-positive predictions of GIs. Also, foreign DNA tends to undergo amelioration in the host genome, limiting the detection of ancient GIs and leading to false-negative predictions (Karlin, 2001; Langille et al., 2010). The transfer of GIs from a donor to the recipient species with a similar sequence composition will also result in false-negatives. An additional difficulty is the fact that several GIs are not inserted next to tRNA genes and not all are flanked by direct repeats (Vernikos and Parkhill, 2008). As such, comparative genomic analysis and phylogeny-based methods should be engaged with sequence composition to perform a more robust

prediction (Langille et al., 2010). Comparative genomics relies on the availability of related sequenced genomes. Moreover, if the genomes are too closely related, GIs that were inserted before the genomes diverged may not be predicted. Using at least one genome that has recently diverged, will lead to more accurate predictions (Langille et al., 2010).

In *P. aeruginosa*, the spread of AR via GIs is greatly underappreciated (Partridge et al., 2018b; Stokes et al., 2012). Few reports have explored the role of these elements for the transfer of CEGs (Botelho et al., 2018c; Chowdhury et al., 2017; Di Pilato et al., 2015; Hong et al., 2016; Martinez et al., 2012; Roy Chowdhury et al., 2016). Chowdhury et al. reported that GIs 1 and 2 are frequently targeted by Tn6060-like and Tn6163-like carbapenemase-harboring transposons, respectively (Chowdhury et al., 2017; Roy Chowdhury et al., 2016).

P. aeruginosa genomic islands that carry genes encoding for virulence factors are collectively known as pathogenicity islands (PAPI) and help to promote the pathogenic promiscuity of specific strains. For instance, several GIs were reported in *P. aeruginosa* LESB58 strain from a CF patient (Jani et al., 2016; Winstanley et al., 2009). Two GIs (PAPI-1 and PAPI-2) were reported in a *P. aeruginosa* PA14 highly virulent strain from a burn patient (Harrison et al., 2010; He et al., 2004). These elements present a mosaic structure and may contribute individually and synergistically to virulence (He et al., 2004). PAPI-2 harbors the *exoU* gene and its chaperone, *spcU*. ExoU is a phospholipase effector cytotoxin that, alongside ExoY, ExoT and ExoS, is secreted by the *P. aeruginosa* type-III secretion system (Dortet et al., 2018; Juan et al., 2017a; Sawa et al., 2016). ExoY and ExoT are identified in the majority of strains, while ExoS and ExoU are mutually exclusive. In fact, *exoS* + *exoU*- genotype is typically associated with an invasive phenotype, while the *exoS* - *exoU* + genotype is related to highly cytotoxic strains such as PA14 and frequently linked to chronic infections (Diaz and Hauser, 2010; Juan et al., 2017a). ExoU is a potent virulence determinant in animal models and is associated with poor clinical outcomes in human patients. The *exoU* gene was also identified on GIs related to PAPI-2 (ExoU islands A, B and C) (Botelho et al., 2018a; Kulasekara et al., 2006).

Currently, *in silico* analysis of GIs may be accomplished by IslandViewer 4 (<http://www.pathogenomics.sfu.ca/islandviewer/>), an integrated interface of four different GI prediction methods: IslandPick, IslandPath-DIMOB, SIGI-HMM, and Islander (Bertelli et al., 2017).

ICEs are self-transmissible mosaic and modular MGEs that combine features of i) transposons and phages, since ICEs can also integrate into and excise from the chromosome; and ii) plasmids, since ICEs can also exist as circular extrachromosomal elements and can be transferred by




Fig. 3. The HGT of an ICE. Besides being vertically transmitted to daughter cells during cell division, ICEs can be transferred to other bacteria by conjugation. ICEs are normally integrated in the chromosome and under certain conditions can be excised and form a plasmid-like circular intermediate (Step 1). In the presence of recipient cells, donor cells can undergo a conjugative transfer event of a single DNA strand by an ICE-encoded T4SS channel (green cylinder) through rolling circle replication (Step 2). DNA replication in the recipient bacterium generates the complementary strand to synthesize the double stranded, circular form of the ICE. Recombination events result in integration of the ICE into the chromosome of each cell (Step 3).

Box 2**The mosaic structure of ICEs.**

Understanding the life cycles of phages, plasmids and transposons is mandatory to correctly inform our views on ICEs. The regulatory and maintenance mechanisms are similar to those of phages and plasmids; integration/excision to phages and transposons; and conjugative transfer to plasmids (Burrus, 2017; Carraro et al., 2015; Cury et al., 2017; Delavat et al., 2017; Guglielmini et al., 2011; Johnson and Grossman, 2015; Kung et al., 2010; Toleman and Walsh, 2011; Wozniak and Waldor, 2010). In fact, a growing body of evidence is challenging the line separating plasmids and ICEs (Delavat et al., 2017; Guglielmini et al., 2011; Johnson and Grossman, 2015). Besides the aforementioned common features, several ICEs are also capable of autonomous rolling-cycle replication, which may be critical to facilitate ICE maintenance (Johnson and Grossman, 2015). Additionally, the relaxase and the T4SS encoded by ICEs resembles that of conjugative plasmids (Burrus, 2017; Carraro et al., 2015; Cury et al., 2017; Delavat et al., 2017; Guglielmini et al., 2011; Johnson and Grossman, 2015; Kung et al., 2010; Toleman and Walsh, 2011; Wozniak and Waldor, 2010). Like mobilizable plasmids, these GIs may not harbor the complete machinery for self-conjugation and may use the T4SS apparatus from a co-resident ICE or from a conjugative plasmid to be successfully transferred in *trans*. In this case, these GIs are named integrative and mobilizable elements (Guédon et al., 2017). Guglielmini et al. constructed a phylogenetic tree of VirB4, a highly conserved ATPase from the T4SS apparatus of different conjugative plasmids and ICEs, and formulated the hypothesis of interchangeable conjugation modules along the evolutionary history of these elements (Guglielmini et al., 2011). VirB4 proteins encoded by ICEs are, however, more related to those encoded by other ICEs and the same happened for plasmids. A close interplay between these elements in the deepest clades of the phylogenetic tree was observed, suggesting that plasmids may behave like ICEs and/or *vice versa*. A search of more than 1000 genomes revealed that ICEs are present in most bacterial clades and may be more prevalent than conjugative plasmids in all clades (Guglielmini et al., 2011). These researchers also found that mobilizable plasmids and integrative and mobilizable elements outnumber conjugative plasmids and ICEs, suggesting a wide utilization of T4SS in *trans*. The majority of ICEs and integrative and mobilizable elements analyzed were found in only one copy per genome (Bellanger et al., 2014). Some ICEs may have become fixed into the chromosome due to degeneration of its phage integrase genes and/or conjugative elements. ICEs are unlikely to share just one common ancestor. Since a wide variety of functional modules has been identified, the hypothesis that different ICE groups arose independently and evolved through multiple recombination events (with specific modules being acquired at different times and from different sources), seems more robust (Kung et al., 2010; Wozniak and Waldor, 2010).

conjugation (Fig. 3 and Box 2) (Burrus, 2017; Carraro et al., 2015; Cury et al., 2017; Delavat et al., 2017; Guglielmini et al., 2011; Johnson and Grossman, 2015; Kung et al., 2010; Toleman and Walsh, 2011; Wozniak and Waldor, 2010). These elements replicate as part of the host genome and are vertically inherited with the chromosome, remaining quiescent and with most of their mobility genes repressed (Burrus, 2017; Carraro et al., 2015; Carraro and Burrus, 2015; Delavat et al., 2017). ICEs are a subset of GIs (since they may also present a different GC content than that of the host chromosome, may harbor phage-related genes and may be flanked by direct and/or inverted repeats) which encode a self-conjugative transfer and integration modules (Langille et al., 2010; Toleman and Walsh, 2011). Four mating-pair formation (MPF) classes cover the T4SS among Proteobacteria: MPF_T, MPF_G, MPF_F and MPF_I (Guglielmini et al., 2013). The first is widely disseminated among conjugative plasmids and ICEs, while MPF_F is more prevalent in plasmids of γ -Proteobacteria and MPF_G is found essentially on ICEs. MPF_I is rarely identified (Guglielmini et al., 2013).

Conjugative transposons are also defined as ICEs. In fact, the first ICE to be discovered was Tn916 (back then defined as a conjugative transposon) from *Enterococcus faecalis* (Flannagan and Clewell, 1991). Since Tn916-like ICEs are major drivers of HGT of AR genes (Roberts and Mullany, 2011), strategies to block their transposition by interfering with the transposase-DNA complex architecture were recently proposed as new possibilities to overcome AR (Rubio-Cosials et al., 2018).

ICEs are encompassed by integration/excision, conjugation, maintenance/regulation and accessory modules, the latter being associated with virulence, catabolic functions and/or AR and displaying little relationship between them (Burrus, 2017; Carraro et al., 2015; Cury et al., 2017; Delavat et al., 2017; Guglielmini et al., 2011; Johnson and Grossman, 2015; Kung et al., 2010; Toleman and Walsh, 2011; Wozniak and Waldor, 2010). These accessory or cargo genes are inserted in hotspot regions without disrupting ICE function. Interchangeable functional modules play a major role in shaping the evolution of ICEs (Burrus, 2017; Carraro et al., 2015; Cury et al., 2017; Delavat et al., 2017; Guglielmini et al., 2011; Johnson and Grossman, 2015; Kung et al., 2010; Toleman and Walsh, 2011; Wozniak and Waldor, 2010). Frequently, genes encoding for similar functions are clustered together. Interestingly, a recent study reported that separate chromosomal clusters may be regrouped as ‘tripartite’ ICE and be later transferred

(Haskett et al., 2016).

As most GIs, ICEs frequently target a single insertion site, which is often a tRNA gene (Botelho et al., 2018c; Burrus, 2017; Carraro et al., 2015; Cury et al., 2017; Delavat et al., 2017; Guglielmini et al., 2011; Johnson and Grossman, 2015; Kung et al., 2010; Toleman and Walsh, 2011; Wozniak and Waldor, 2010). Even though multiple copies of these genes may be present among different hosts, ICEs tend to target only one of these loci, most likely due to specific interactions between the attachments sites of the ICE (*attP*, based on the phage nomenclature) and of the corresponding bacterial host (*attB*). However, promiscuous ICE can insert in different sites and so can be found in several locations of some bacterial genomes. This insertion is mediated by site-specific recombination promoted by a phage-like integrase, most frequently belonging to the tyrosine recombinase family. These enzymes may also be involved in the excision process (Burrus, 2017; Carraro et al., 2015; Cury et al., 2017; Delavat et al., 2017; Guglielmini et al., 2011; Johnson and Grossman, 2015; Kung et al., 2010; Toleman and Walsh, 2011; Wozniak and Waldor, 2010). The IntB13 tyrosine recombinase from ICE_{clc} is an example of a P4-family integrase with unusual long length that mediates both integration and excision of the ICE (Gaillard et al., 2006; Ravatn et al., 1998).

Most ICEs identified among *P. aeruginosa* fall into two families: pKLC102-like and *clc*-related ICEs (Kung et al., 2010), a distinction that is not always consensual. For example, Klockgether et al. preferred to consider these elements as members of the same family with a common ancestry, taking into account the conserved function and synteny of the backbone genes, sharing an amino acid sequence identity of at least 20% (Klockgether et al., 2011). Main differences between these families are related to the integration site and the type of integrase: ICEs related to pKLC102 (such as PAGI-4, PAGI-5 and aforementioned ExoU island A and pathogenicity islands PAPI-1 and PAPI-2) frequently integrate into the 3' end of tRNA^{Lys} genes and present XerC/XerD-like integrases, while ICE_{clc}-like elements (such as PAGI-2, PAGI-3 and LESGI-3) are more prone to target the 3' end of a cluster of tRNA^{Gly} genes and frequently harbor a bacteriophage P4-integrase (Botelho et al., 2018c; Kung et al., 2010; Larbig et al., 2002; Toleman and Walsh, 2011). Also, pKLC102-like ICEs typically share a set of syntenic conserved core genes, while ICE_{clc}-like elements tend to exhibit a bipartite structure, with a conserved portion next to the integration site and unique cargo genes in the remaining ICE sequence (Kung et al., 2010). Interestingly,

Fischer et al. found that ICEs from the pKLC102 and ICE_{cl} families were frequently identified among several representatives of the major *P. aeruginosa* clones C and PA14 and were the main contributors for genomic diversity (Fischer et al., 2016). The ICE Tn4371 family also represents a large group of ICEs with a common backbone and which are widely distributed, such as in *P. aeruginosa* PA7 and 2192 (Ryan et al., 2009; Toleman and Walsh, 2011).

To date, few reports of carbapenemases were associated with an ICE location in *P. aeruginosa*, such as a *bla*_{GES-5}-harboring GI2 identified in Australia, a *bla*_{GES-6}-bearing ICE retrieved from a representative of the high-risk clone ST235, a *bla*_{SPM-1} inserted into ICE_{Tn4371}6061 and reported in Brazil and a *bla*_{NDM-1}-harboring ICE from Singapore and Poland (Botelho et al., 2018a; Chowdhury et al., 2017; Ding et al., 2018; Fonseca et al., 2015; Urbanowicz et al., 2019). Curiously, these reports are quite recent, leading us to speculate that the advent of WGS and bioinformatics will help to accurately identify more ICEs and eventually assess the real contribution for the spread of CEGs. In fact, a recent *in silico* analysis revealed that ICEs may play a vital role in disseminating CEGs among pseudomonads (Botelho et al., 2018c).

ICEberg 2.0 (<http://db-mmml.sjtu.edu.cn/ICEberg/>) is a web-based resource that provides significant information about ICE identified in Gram-negative and Gram-positive bacteria (M. Liu et al., 2018). ICEberg also provides a tool to predict ICEs in bacterial genome sequences, named ICEfinder. The CONJscan (https://github.com/gem-pasteur/Macsfinder_models) is a tool that scans a set of protein sequences for T4SS and relaxases of both ICE and plasmids using hidden Markov models (Abby et al., 2016; Guglielmini et al., 2014).

9. To be or not to be: the fitness cost and maintenance of antibiotic resistance

Fitness is the capacity of a genotype or individual to survive and reproduce (Andersson and Hughes, 2010; San Millan and MacLean, 2017; Vogwill and MacLean, 2015). The classic paradigm is that AR determinants endeavor a fitness cost for the bacterial host, in terms of reduced growth, competition and/or infectivity (Andersson and Hughes, 2010; Baltrus, 2013; Hall et al., 2017; Loftie-Eaton et al., 2017; Melnyk et al., 2015; Rodriguez-Beltran et al., 2018; San Millan et al.,

2018; Vogwill and MacLean, 2015), even though some important exceptions were reported (Böttger et al., 1998; Sander et al., 2002). It is also assumed that evolution of resistance occurs by only a few mutations of large effect (Andersson and Hughes, 2010; Vogwill and MacLean, 2015). Despite conferring high levels of resistance, mutants with a high fitness cost are less likely to outcompete susceptible strains in a given population. On the contrary, low-level mutations can be fixed if associated with a low fitness cost (Andersson and Hughes, 2010). Paradoxically, the weaker the selection for resistance (for example, by non-lethal antibiotic concentrations) the higher the likelihood for persistence of mutants with a low fitness cost and mutator phenotypes (Andersson and Hughes, 2014). Actually, a growing body of evidence suggests that high levels of AR evolve due to combinations of multiple mutations that exert a minimal epistatic cost to the host (Trindade et al., 2009; Vogwill and MacLean, 2015; Ward et al., 2009; Wistrand-Yuen et al., 2018).

The long-term fate of resistant mutants is largely influenced by the evolution of compensatory mechanisms, which may restore fitness without compromising resistance to antibiotics (Box 3) (Andersson and Hughes, 2010). Indeed, Melnik et al. tracked the development of ciprofloxacin resistance in *P. aeruginosa* under constant and fluctuating antibiotic administration, and found that high-fitness, resistant strains (broadly adapted generalists, as defined by Kassen et al. (Kassen, 2002)) evolved quickly under fluctuating treatments and that second-site mutations were responsible for compensating the fitness cost associated with AR (Melnik et al., 2017). As such, the authors suggested that these therapies generate persistent AR by selecting for the evolution of cost-free resistance of *P. aeruginosa* strains. Pacheco et al. studied the independent overexpression of the four most important efflux pumps involved in AR (MexAB-OprM, MexCD-OprJ, MexEF-OprN and MexXY) by *P. aeruginosa* mutants (Olivares Pacheco et al., 2017). These researchers verified that the lack of fitness cost observed among these mutants was associated with a metabolic rewiring as a compensation derived from overexpressing these pumps. This non-mutational mechanism includes an increased expression of the nitrate respiratory chain under aerobic conditions (Olivares Pacheco et al., 2017).

Genetic elements such as plasmids and ICE frequently harbor AR and/or virulence genes that may provide a selective advantage for the

Box 3

Evolution and fitness cost of HGT.

HGT plays a decisive role in AR spreading among different Gram-negative and Gram-positive bacteria. A survey conducted by Kloeisges et al. suggests that nearly 75% of the genes from a given genome have been horizontally transferred at least once during evolution (Kloeisges et al., 2011). HGT is more frequent between closely related species from the same taxonomic group, having similar genomes and an analogous GC content (Andam and Gogarten, 2011; Hooper et al., 2009; Kloeisges et al., 2011). As so, the pangenome also represents the pool of genes that may be available through HGT to any close relative of the same taxon (Soucy et al., 2015). Another study found bi-directional associations among the similarity in tRNA pools of bacteria and the number of HGT events happening between them (Tuller et al., 2011). Both situations make quantification of HGT quite challenging (Andam and Gogarten, 2011; Polz et al., 2013). Although HGT among distantly related species can still occur, there seems to exist a biological barrier for gene acquisition from donors with different genomic GC content. Species such as *Aeromonas salmonicida* act as a natural mixing pot which harbors *Pseudomonas* and enteric types of plasmids and thus allow the exchange of genetic material (Olsen and Wright, 1976). Most genes subjected to HGT are responsible for metabolic functions, while the transfer of genes related to information processing (such as replication, transcription and translation) is rare (Cohen et al., 2011). Also, HGT was shown to play a more important role than gene duplication for the expansion of protein families among prokaryotes (Treangen and Rocha, 2011). HGT may also depend on the transfer mechanism (Popa et al., 2011; Popa and Dagan, 2011). As a consequence of HGT, prokaryotic genome evolution models should be viewed less as a bifurcated tree, and more as network models, which allow the depiction of several events such as recombination, gene fusion and HGT (Harrison and Brockhurst, 2012; Kloeisges et al., 2011; Polz et al., 2013; Popa et al., 2011; Soucy et al., 2015; Treangen and Rocha, 2011).

Once foreign DNA has been integrated into the host, it might need to adapt within the genome in order to resist purifying selection (Moran et al., 2009). In order to survive, the genes usually need to incur in a low fitness cost to the host and/or to provide a selective advantage to themselves or to the recipient strain (Baltrus, 2013). Their expression will largely be governed by the presence of promoters and/or regulatory mechanisms on the vicinity of its insertion loci. Nonetheless, comparative genomic analysis unveiled that many genes acquired by HGT appear to have neutral effects for the host (Gogarten and Townsend, 2005; Soucy et al., 2015). For example, introns are self-splicing elements that provide a nearly neutral effect, and can persist since the success in invading hosts compensates for the fitness cost to the host associated with the additional DNA, RNA and protein synthesis. Neutral acquisitions can later be subjected to novel combinations, allowing its domestication over time if a beneficial phenotype is guaranteed. If imported genes remain neutral, purifying selection will most likely lead to its loss (Gogarten and Townsend, 2005; Soucy et al., 2015).

host. Acquisition of a useful gene pool by a host genome could counterbalance the fitness cost associated with maintaining these elements (Broaders et al., 2013; Rankin et al., 2011). Even though every HGT may exert a fitness cost, and even if these costs are offset in cases of benefits for the host, these events leave lasting genomic signatures that shape evolution of the MGE to fit within the host genome and/or the host to accommodate the new guest (Baltrus, 2013).

Besides selection on beneficial traits, efficient replication, segregation, re-infection by conjugation, growth inhibition or killing of plasmid-free cells by toxin-antitoxin systems and a low cost have an important role in maintaining plasmids harboring AR genes in a given population (Loftie-Eaton et al., 2017, 2016; Lopatkin et al., 2017; Sommer et al., 2017). To accommodate this MGE, compensatory mutations may occur in the plasmid, the host or in both (Harrison et al., 2015; Harrison and Brockhurst, 2012; Kottara et al., 2018; Loftie-Eaton et al., 2017; San Millan et al., 2014b; Rodriguez-Beltran et al., 2018; San Millan et al., 2016, 2015b; San Millan and MacLean, 2017; Stevenson et al., 2018; Zwanzig et al., 2019). Given the abundance of non-transmissible plasmids (Smillie et al., 2010), San Millan et al. showed that epistatic interactions between co-resident plasmids unable to be transferred by conjugation, decreases the cost associated with harboring several plasmids in *P. aeruginosa* and promotes persistence (San Millan et al., 2014a). Loftie-Eaton et al. uncovered a plasmid-host epistasis mechanism promoting plasmid stability in the coevolved host through acquisition of a transposon-encoded putative toxin-antitoxin and co-integrate resolution system, alongside host mutations (Loftie-Eaton et al., 2016). The researchers also found that this transposition expanded plasmid-host range. Another study reported how compensatory adaptation and positive selection interact to assure persistence of non-conjugative plasmid (San Millan et al., 2014b). Harrison et al. explored the compensatory evolution mechanisms responsible for ameliorating the cost of maintaining the plasmid, in both parasitic (cost prevail over benefits, a situation where plasmids are frequently lost due to purifying selection) and mutualistic (benefits take the lead and selection favors the capture of beneficial genes into the chromosome and subsequent loss of the plasmid backbone) treatments (Harrison et al., 2015). Such plasmid-plasmid and plasmid-host interactions expose the multiple solutions for co-evolution, and so to better understand HGT one should use co-evolutionary instead of a simple evolutionary approach (Harrison and Brockhurst, 2012). However, co-evolution may not always lead to amelioration of plasmid burden (Smith, 2011). Most plasmids carry several AR genes, increasing the probability of co-selection and thereby decreasing the effect of purifying selection on a high fitness cost resistance gene. Worryingly, this effect could dissociate the evolution of resistance to a given antibiotic from the use of the same agent (Sommer et al., 2017; Sundqvist et al., 2010; Yen and Papin, 2017).

However, evolving resistance by plasmid acquisition exerts a higher cost than resistance by chromosomal mutations? The answer is not simple. Apparently, the acquisition of a MGE by the host frequently imposes a fitness, with a high cost. However, AR mutations in essential and highly conserved genes are likely to be more costly (Aleksun and Levy, 2007). A recent meta-analysis approach revealed that evolving resistance by acquisition of plasmids was associated with a smaller cost, which could help to explain why these elements are so widely disseminated (Vogwill and MacLean, 2015). Also, the host may re-acquire a plasmid that was carried during its evolutionary past, and so compensatory mutations (within the host's chromosome or the plasmid) can offset the cost of the recently acquired plasmid (Loftie-Eaton et al., 2017; Vogwill and MacLean, 2015). This host-plasmid coevolution could also help to explain why acquisition of the same plasmids by different hosts exerts a different cost (Starikova et al., 2013).

ICEs appear to have a bistable lifestyle that determines the change between vertical and horizontal transmission (Delavat et al., 2017). In the ICE integrated state, core functions are downregulated in order to exert a low fitness cost on the host and to ensure the likelihood of

vertical transmission, while beneficial genes carried by the ICEs may pose a selective advantage to the host; the excision and subsequent horizontal transfer state can be activated at low frequencies upon external triggers in specific growth conditions (Burrus, 2017; Delavat et al., 2017, 2016). Activation occurs at a low level most likely to avoid harming the host and to guarantee sufficient HGT and population survival. So far, this theory has only been observed for ICE_{cl} (Delavat et al., 2016; Gaillard et al., 2008; Miyazaki et al., 2018), and single cell observations should be extended to other ICE systems. Future studies should also address the mechanisms producing and maintaining this dual lifestyle and the contribution of selective forces driving ICE-host co-evolution.

10. Population structure of *P. aeruginosa* and the impact of high-risk clones

P. aeruginosa has a non-clonal epidemic population structure, punctuated by specific sequence types (ST) (Jolley et al., 2018; Pirnay et al., 2009). Its population structure is comparable to that from *Neisseria meningitidis*, a superficially clonal structure with high recombination frequency, in which rare successful epidemic lineages tend to arise. There seems to be a consensus that clinical and environmental isolates are indistinguishable and that there is no association between a given clone and a specific habitat (Piray et al., 2009). Hospital transmission can, however, increase the prevalence of particularly adapted clones (Oliver et al., 2015).

Some STs are worldwide disseminated and frequently linked to outbreaks and to the dissemination of carbapenemases. These STs have been designated high-risk clones, of which major examples are ST235, ST111 and ST175 in *P. aeruginosa* (Kos et al., 2015; Oliver et al., 2015; Sánchez-Diener et al., 2017; Woodford et al., 2011). The bla_{VIM-2} was the most frequently encountered CEG among ST235 and ST111 (Oliver et al., 2015; Turton et al., 2015). Albeit being responsible for the spread of AR through HGT, these STs have also been associated with mutational resistance. In fact, the mechanisms responsible for the multidrug resistant phenotypes frequently observed in ST175 are mostly determined by specific mutations on *oprD*, *ampR*, *mexZ*, *gyrA* and *parC* (Cabot et al., 2016, 2012; López-Causapé et al., 2018).

Among the few major high-risk clones, ST235 is the most widely disseminated (Chowdhury et al., 2017; Kos et al., 2015; Maatallah et al., 2011; Miyoshi-Akiyama et al., 2017; Oliver et al., 2015; Roy Chowdhury et al., 2016; Treepong et al., 2018; van Belkum et al., 2015). Nearly 100 horizontally-acquired resistance elements have been reported among ST235 isolates (Chowdhury et al., 2017; Oliver et al., 2015; Roy Chowdhury et al., 2016; Treepong et al., 2018). The *dprA* gene was identified by Treepong et al. as a specific marker of this lineage (Treepong et al., 2018). Due to the reported contribution of DprA in homologous recombination and acquisition of foreign DNA, its presence in ST235 lineage may help to explain the increased ability to acquire AR elements. Class A and B carbapenemases were also frequently associated with ST111 isolates (Castanheira et al., 2014; Chowdhury et al., 2017; Kos et al., 2015; Oliver et al., 2015; Turton et al., 2015; van Belkum et al., 2015).

The geographical distribution of the high-risk clones and the diversity of AR elements suggest that the spread of these STs is global and the acquisition of the resistance genes is mainly local (Oliver et al., 2015; Turton et al., 2015). Previous studies suggest that environmental species may constitute an important reservoir for the dissemination of clinically relevant carbapenemases, which are vertically amplified upon transfer to *P. aeruginosa* high-risk clones (Jiang et al., 2017; Juan et al., 2010; Scotta et al., 2011). The high prevalence of these elements among high-risk clones may be partially explained by the genetic capitalism theory, given that a widely disseminated ST should have a greater probability of acquiring new AR genes and to be further selected and amplified due to the high antibiotic pressure in the hospital environment (Baquero, 2004). Other theories support that the high-risk clones

have a naturally increased ability to acquire foreign DNA, since these STs appear to have lost the CRISPR (clustered regularly interspaced short palindromic repeats)-Cas (CRISPR associated proteins) system, which act as an adaptive immune system in prokaryotic cells and protects them from invasion by bacteriophages and plasmids (Bondy-Denomy and Davidson, 2014; Chen et al., 2018; England et al., 2018; Li et al., 2016; Turton et al., 2015; van Belkum et al., 2015).

Mulet et al. studied the biological markers that may be responsible for the success of *P. aeruginosa* high-risk clones (Mulet et al., 2013). The authors found that these STs were defective in motility and pigment production, presented a reduced *in vitro* fitness in competition experiments and showed increased levels of biofilm formation and spontaneous mutant frequencies. Interestingly, these traits resemble those resulting from adaptation to chronic infections (Mulet et al., 2013).

Treeping et al. linked the dissemination of ST235 with the selective pressure exerted by fluoroquinolones (Treeping et al., 2018). A similar observation for ST175 was previously reported by Cabot et al., since all strains tested presented the same set of quinolone resistance-determining region mutations (Cabot et al., 2012). Curiously, as above-mentioned, some mutations did not involve biological costs, as those targeting *gyrA* and *gyrB* genes (Andersson and Hughes, 2010). Conferring low levels of AR without incurring in fitness cost can help to explain the high frequency of gyrase mutations in several clinical isolates (Wong and Kassen, 2011). Future studies performing comparative genomic analysis of high-risk clones and non-epidemic multi-drug resistant strains will help to shed a new light on the traits responsible for the success of these STs. Moreover, the existence of specific features that promote the development of *de novo* mutations and/or the acquisition of AR genes by HGT still needs to be further explored (Oliver et al., 2015).

11. Conclusions and future directions

An intriguing question is why the class I integrons tend to be integrated into the chromosome rather than inserted in plasmids. Conducting an *in silico* analysis of CEGs deposited in public databases and analyzing specific traits may help to explain this preference. It would also be interesting to explore the molecular basis for why chromosomal mutations are more important for resistance to carbapenems than acquisition of MGEs harboring AR genes. One possible explanation may be the burden of HGT to the host, and combination of multiple mutations that exert a minimal epistatic cost to the host may yield high levels of AR *per se* (Kassen, 2002; Trindade et al., 2009; Vogwill and MacLean, 2015; Ward et al., 2009).

High-risk clones appear to be more capable of acquiring and/or maintaining AR genes when comparing with other clones. However, the underlying basis for this remains a mystery. Previous reports suggested that the absence, modification or downregulation of proper defense systems against foreign DNA (such as CRISPR-Cas, toxin-antitoxin and phage-exclusion) in high-risk clones would lead to its increased invasion by MGEs (Bondy-Denomy and Davidson, 2014; Kos et al., 2015; Turton et al., 2015; van Belkum et al., 2015). These observations should be extended to a wider pool of clones in order to address if these systems are indeed absent from all representatives of this subset of *P. aeruginosa* lineages. Another possible explanation for this riddle could be that high-risk clones may share stable lineage-specific GIs that are highly capable of acquiring AR gene cassettes by the class I integrons presented in these platforms. Performing comparative genomic studies to identify the presence/absence of specific genes in high-risk clones and in other *P. aeruginosa* strains may help to address the reason why these clones are so successful. It would also be interesting to expand the work conducted by Khaledi et al. and perform a RNA-seq analysis to determine whether high-risk clones exhibit a different gene expression profile (Bruchmann et al., 2015; Khaledi et al., 2016). Another question that needs to be addressed is the nature of high-risk clones (Woodford et al., 2011): are these clones a minor pre-existing variant that recently

succeeded, with resistance as a key driver, and/or a previously successful antibiotic-susceptible variant that underwent clonal expansion due to the acquisition of AR genes?

In the future, the approach taken to study HGT and the spread of AR genes in *P. aeruginosa* needs to be rethought, and we believe a greater emphasis should be placed on the contribution of GIs and more specifically ICEs. With the advent of WGS, the real contribution of these MGEs for the dissemination of CEGs will be correctly addressed. WGS analysis, together with epidemiological data, may also help to give further insights into the differences between members of high-risk clones and may be able to identify the routes of transmission of *P. aeruginosa* between patients and different sources. More work is needed to identify the main hospital reservoirs of MDR *P. aeruginosa*, which may include plumbing systems, ventilators and/or long-term colonization of patients (Livingston et al., 2018). It is vital to have an in-depth understanding of these MDR organisms in order to prevent infections and design control strategies to limit their future spread. The genomic era and integrated molecular epidemiology approaches may help outbreak investigations, genome-based prediction of AR and long-term persistence surveillance of MDR *P. aeruginosa* in real-time (Boolchandani et al., 2019; S. Baker et al., 2018; Besser et al., 2018; Davis et al., 2015; Gardy and Loman, 2017; Quainoo et al., 2017; Quick et al., 2014; Rossen et al., 2018; Su et al., 2018; Turton et al., 2015).

Acknowledgements

This work was supported by the Applied Molecular Biosciences Unit-UCIBIO which is financed by national funds from FCT/MCTES (UID/Multi/04378/2019). JB and FG were supported by grants from Fundação para a Ciência e a Tecnologia (SFRH/BD/104095/2014 and SFRH/BPD/95556/2013, respectively).

References

- Abby, S.S., Cury, J., Guglielmini, J., Néron, B., Touchon, M., Rocha, E.P.C., 2016. Identification of protein secretion systems in bacterial genomes. *Sci. Rep.* 6, 23080. <https://doi.org/10.1038/srep23080>.
- Alekshun, M.N., Levy, S.B., 2007. Molecular mechanisms of antibacterial multidrug resistance. *Cell* 128, 1037–1050. <https://doi.org/10.1016/j.cell.2007.03.004>.
- Andam, C.P., Gogarten, J.P., 2011. Biased gene transfer in microbial evolution. *Nat. Rev. Microbiol.* 9, 543–555. <https://doi.org/10.1038/nrmicro2593>.
- Andersen, S.B., Ghoul, M., Marvig, R.L., Lee, Z.-B., Molin, S., Johansen, H.K., Griffin, A.S., 2018. Privatisation rescues function following loss of cooperation. *Elife* 7, e38594. <https://doi.org/10.7554/eLife.38594>.
- Andersson, D.I., Hughes, D., 2014. Microbiological effects of sublethal levels of antibiotics. *Nat. Rev. Microbiol.* 12, 465–478. <https://doi.org/10.1038/nrmicro3270>.
- Andersson, D.I., Hughes, D., 2010. Antibiotic resistance and its cost: is it possible to reverse resistance? *Nat. Rev. Microbiol.* 8, 260–271. <https://doi.org/10.1038/nrmicro2319>.
- Antipov, D., Hartwick, N., Shen, M., Raiko, M., Lapidus, A., Pevzner, P.A., 2016. plasmidSPAdes: assembling plasmids from whole genome sequencing data. *Bioinformatics* 32, 3380–3387. <https://doi.org/10.1093/bioinformatics/btw493>.
- Arredondo-Alonso, S., Willems, R.J., van Schaik, W., Schürch, A.C., 2017. On the (im)possibility of reconstructing plasmids from whole-genome short-read sequencing data. *Microb. Genom.* 3, e000128. <https://doi.org/10.1099/mgen.0.000128>.
- Baker, S., Thomson, N., Weill, F.-X., Holt, K.E., 2018. Genomic insights into the emergence and spread of antimicrobial-resistant bacterial pathogens. *Science* 360, 733–738. <https://doi.org/10.1126/science.aar3777>.
- Baker, S.J., Payne, D.J., Rappuoli, R., De Gregorio, E., 2018. Technologies to address antimicrobial resistance. *Proc. Natl. Acad. Sci. U. S. A.* 115, 12887–12895. <https://doi.org/10.1073/pnas.1717160115>.
- Balasubramanian, D., Kumari, H., Mathee, K., 2014. *Pseudomonas aeruginosa* AmpR: an acute-chronic switch regulator. *Pathog. Dis.* 73, 1–14. <https://doi.org/10.1111/2049-632X.12208>.
- Balasubramanian, D., Schnepfer, L., Kumari, H., Mathee, K., 2013. A dynamic and intricate regulatory network determines *Pseudomonas aeruginosa* virulence. *Nucleic Acids Res.* 41, 1–20. <https://doi.org/10.1093/nar/gks1039>.
- Baltrus, D.A., 2013. Exploring the costs of horizontal gene transfer. *Trends Ecol. Evol.* 28, 489–495. <https://doi.org/10.1016/j.tree.2013.04.002>.
- Baquero, F., 2004. From pieces to patterns: evolutionary engineering in bacterial pathogens. *Nat. Rev. Microbiol.* 2, 510–518. <https://doi.org/10.1038/nrmicro909>.
- Barbosa, C., Beardmore, R., Schulenburg, H., Jansen, G., 2018. Antibiotic combination efficacy (ACE) networks for a *Pseudomonas aeruginosa* model. *PLOS Biol.* 16, e2004356. <https://doi.org/10.1371/journal.pbio.2004356>.
- Barbosa, C., Trebosc, V., Kemmer, C., Rosenstiel, P., Beardmore, R., Schulenburg, H.,

- Jansen, G., 2017. Alternative evolutionary paths to bacterial antibiotic resistance cause distinct collateral effects. *Mol. Biol. Evol.* 34, 2229–2244. <https://doi.org/10.1093/molbev/msx158>.
- Barnes, M.D., Taracila, M.A., Rutter, J.D., Bethel, C.R., Galdadas, I., Hujer, A.M., Caselli, E., Prati, F., Dekker, J.P., Papp-Wallace, K.M., Haider, S., Bonomo, R.A., 2018. Deciphering the evolution of cephalosporin resistance to ceftolozane-tazobactam in *Pseudomonas aeruginosa*. *MBio* 9, e02085-18. <https://doi.org/10.1128/mbio.02085-18>.
- Basseti, M., Vena, A., Croxatto, A., Righi, E., Guery, B., 2018. How to manage *Pseudomonas aeruginosa* infections. *Drugs Context* 7, 212527. <https://doi.org/10.7573/dic.212527>.
- Baxter, J.C., Funnell, B.E., 2014. Plasmid partition mechanisms. *Microbiol. Spectr.* 2. <https://doi.org/10.1128/microbiolspec.PLAS-0023-2014>.
- Bell, G., MacLean, C., 2018. The search for “evolution-proof” antibiotics. *Trends Microbiol.* 26, 471–483. <https://doi.org/10.1016/j.tim.2017.11.005>.
- Bellanger, X., Payot, S., Leblond-Bourget, N., Guédon, G., 2014. Conjugative and mobilizable genomic islands in bacteria: evolution and diversity. *FEMS Microbiol. Rev.* 38, 720–760. <https://doi.org/10.1111/1574-6976.12058>.
- Belotti, P.T., Thabet, L., Laffargue, A., André, C., Coulange-Mayonnove, L., Arpin, C., Messadi, A., M'Zali, F., Quentin, C., Dubois, V., 2015. Description of an original integron encompassing *bla*_{VIM-2}, *qnrVCI* and genes encoding bacterial group II intron proteins in *Pseudomonas aeruginosa*. *J. Antimicrob. Chemother.* 70, 2237–2240. <https://doi.org/10.1093/jac/dkv103>.
- Benveniste, R., Davies, J., 1973. Aminoglycoside antibiotic-inactivating enzymes in actinomycetes similar to those present in clinical isolates of antibiotic-resistant bacteria. *Proc. Natl. Acad. Sci. U. S. A.* 70, 2276–2280. <https://doi.org/10.1073/pnas.70.8.2276>.
- Berglund, F., Marathe, N.P., Österlund, T., Bengtsson-Palme, J., Kotsakis, S., Flach, C.-F., Larsson, D.G.J., Kristiansson, E., 2017. Identification of 76 novel B1 metallo- β -lactamases through large-scale screening of genomic and metagenomic data. *Microbiome* 5, 134. <https://doi.org/10.1186/s40168-017-0353-8>.
- Berrazeg, M., Jeannot, K., Ntsogo Enguéné, V.Y., Broutin, I., Loeffert, S., Fournier, D., Plésiat, P., 2015. Mutations in β -lactamase AmpC increase resistance of *Pseudomonas aeruginosa* isolates to antipseudomonal cephalosporins. *Antimicrob. Agents Chemother.* 59, 6248–6255. <https://doi.org/10.1128/AAC.00825-15>.
- Bertelli, C., Laird, M.R., Williams, K.P., Lau, B.Y., Hoad, G., Winsor, G.L., Brinkman, F.S., Brinkman, F.S.L., 2017. IslandViewer 4: expanded prediction of genomic islands for larger-scale datasets. *Nucleic Acids Res.* 45, W30–W35. <https://doi.org/10.1093/nar/gkx343>.
- Besser, J., Carleton, H.A., Gerner-Smith, P., Lindsey, R.L., Trees, E., 2018. Next-generation sequencing technologies and their application to the study and control of bacterial infections. *Clin. Microbiol. Infect.* 24, 335–341. <https://doi.org/10.1016/j.cmi.2017.10.013>.
- Bikkarolla, S.K., Nordberg, V., Rajer, F., Müller, V., Kabir, M.H., KKS, Dvyrnas, A., Ambjörnsson, T., Giske, C.G., Navér, L., Sandegren, L., Westerlund, F., 2019. Optical DNA mapping combined with cas 9-targeted resistance gene identification for rapid tracking of resistance plasmids in a neonatal intensive care unit outbreak. *MBio* 10, e00347-19. <https://doi.org/10.1128/mbio.00347-19>.
- Blair, J.M.A., Webber, M.A., Baylay, A.J., Ogbolu, D.O., Piddock, L.J.V., 2015. Molecular mechanisms of antibiotic resistance. *Nat. Rev. Microbiol.* 13, 42–51. <https://doi.org/10.1038/nrmicro3380>.
- Boissy, R., Ahmed, A., Janto, B., Earl, J., Hall, B.G., Hogg, J.S., Pusch, G.D., Hiller, L.N., Powell, E., Hayes, J., Yu, S., Kathju, S., Stoodley, P., Post, J.C., Ehrlich, G.D., Hu, F.Z., 2011. Comparative supragenomic analyses among the pathogens *Staphylococcus aureus*, *Streptococcus pneumoniae*, and *Haemophilus influenzae* using a modification of the finite supragenome model. *BMC Genom.* 12, 187. <https://doi.org/10.1186/1471-2164-12-187>.
- Bondy-Denomy, J., Davidson, A.R., 2014. To acquire or resist: the complex biological effects of CRISPR-Cas systems. *Trends Microbiol.* 22, 218–225. <https://doi.org/10.1016/j.tim.2014.01.007>.
- Bonnin, R.A., Bogaerts, P., Girlich, D., Huang, T.-D., Dortet, L., Glupczynski, Y., Naas, T., 2018. Molecular characterization of OXA-198 carbapenemase-producing *Pseudomonas aeruginosa* clinical isolates. *Antimicrob. Agents Chemother.* 62, e02496–e02517. <https://doi.org/10.1128/AAC.02496-17>.
- Bonnin, R.A., Poiré, L., Nordmann, P., Eikmeyer, F.G., Wibberg, D., Pühler, A., Schlüter, A., 2013. Complete sequence of broad-host-range plasmid pNOR-2000 harbouring the metallo- β -lactamase gene *bla*_{VIM-2} from *Pseudomonas aeruginosa*. *J. Antimicrob. Chemother.* 68, 1060–1065. <https://doi.org/10.1093/jac/dks526>.
- Boolchandani, M., D'Souza, A.W., Dantas, G., 2019. Sequencing-based methods and resources to study antimicrobial resistance. *Nat. Rev. Genet.* 20, 356–370. <https://doi.org/10.1038/s41576-019-0108-4>.
- Botelho, J., Grosso, F., Peixe, L., 2018a. Unravelling the genome of a *Pseudomonas aeruginosa* isolate belonging to the high-risk clone ST235 reveals an integrative conjugative element housing a *bla*_{GES-6} carbapenemase. *J. Antimicrob. Chemother.* 73, 77–83. <https://doi.org/10.1093/jac/dkx337>.
- Botelho, J., Grosso, F., Peixe, L., 2017a. Characterization of the pJB12 plasmid from *Pseudomonas aeruginosa* reveals Tn6352, a novel putative transposon associated with mobilization of the *bla*_{VIM-2}-harboring *Int58* integron. *Antimicrob. Agents Chemother.* 61, e02532–e02616. <https://doi.org/10.1128/AAC.02532-16>.
- Botelho, J., Grosso, F., Quinteira, S., Brilhante, M., Ramos, H., Peixe, L., 2018b. Two decades of *bla*_{VIM-2}-producing *Pseudomonas aeruginosa* dissemination: an interplay between mobile genetic elements and successful clones. *J. Antimicrob. Chemother.* 73, 873–882. <https://doi.org/10.1093/jac/dkx517>.
- Botelho, J., Grosso, F., Quinteira, S., Mabrouk, A., Peixe, L., 2017b. The complete nucleotide sequence of an IncP-2 megaplasmid unveils a mosaic architecture comprising a putative novel *bla*_{VIM-2}-harbouring transposon in *Pseudomonas aeruginosa*. *J. Antimicrob. Chemother.* 72, 2225–2229. <https://doi.org/10.1093/jac/dkx143>.
- Botelho, J., Grosso, F., Sousa, C., Peixe, L., 2015. Characterization of a new genetic environment associated with GES-6 carbapenemase from a *Pseudomonas aeruginosa* isolate belonging to the high-risk clone ST235. *J. Antimicrob. Chemother.* 70, 615–617. <https://doi.org/10.1093/jac/dku391>.
- Botelho, J., Roberts, A.P., León-Sampedro, R., Grosso, F., Peixe, L., 2018c. Carbapenemases on the move: it's good to be on ICEs. *Mob. DNA* 9, 37. <https://doi.org/10.1186/s13100-018-0141-4>.
- Böttger, E.C., Springer, B., Pletschette, M., Sander, P., 1998. Fitness of antibiotic-resistant microorganisms and compensatory mutations. *Nat. Med.* 4, 1343–1344. <https://doi.org/10.1038/3906>.
- Boucher, H.W., Talbot, G.H., Bradley, J.S., Edwards, J.E., Gilbert, D., Rice, L.B., Scheld, M., Spellberg, B., Bartlett, J., 2009. Bad bugs, no drugs: no ESKAPE! an update from the Infectious Diseases Society of America. *Clin. Infect. Dis.* 48, 1–12. <https://doi.org/10.1086/595011>.
- Bouheraoua, N., Poiré, L., Bourreau, B., Bonnin, R., Laroche, L., Naas, T., Nordmann, P., 2018. Integrase-mediated recombination of the *bel-1* gene cassette encoding the extended-spectrum β -lactamase BEL-1. *Antimicrob. Agents Chemother.* 62, e00030-18. <https://doi.org/10.1128/AAC.00030-18>.
- Brauner, A., Fridman, O., Gefen, O., Balaban, N.Q., 2016. Distinguishing between resistance, tolerance and persistence to antibiotic treatment. *Nat. Rev. Microbiol.* 14, 320–330. <https://doi.org/10.1038/nrmicro.2016.34>.
- Breidenstein, E.B.M., de la Fuente-Núñez, C., Hancock, R.E.W., 2011. *Pseudomonas aeruginosa*: all roads lead to resistance. *Trends Microbiol.* 19, 419–426. <https://doi.org/10.1016/j.tim.2011.04.005>.
- Broaders, E., Gahan, C.G.M., Marchesi, J.R., 2013. Mobile genetic elements of the human gastrointestinal tract. *Gut. Microb.* 4, 271–280. <https://doi.org/10.4161/gmic.24627>.
- Brochado, A.R., Telzerow, A., Bobonis, J., Banzhaf, M., Mateus, A., Selkrig, J., Huth, E., Bassler, S., Zamarreño Beas, J., Zietek, M., Ng, N., Foerster, S., Ezraty, B., Py, B., Barras, F., Savitski, M.M., Bork, P., Göttig, S., Typas, A., 2018. Species-specific activity of antibacterial drug combinations. *Nature* 559, 259–263. <https://doi.org/10.1038/s41586-018-0278-9>.
- Bruchmann, S., Dötsch, A., Nouri, B., Chaberny, I.F., Häussler, S., 2013. Quantitative contributions of target alteration and decreased drug accumulation to *Pseudomonas aeruginosa* fluoroquinolone resistance. *Antimicrob. Agents Chemother.* 57, 1361–1368. <https://doi.org/10.1128/AAC.01581-12>.
- Bruchmann, S., Muthukumarasamy, U., Pohl, S., Preusse, M., Bielecka, A., Nicolai, T., Hamann, I., Hillert, R., Kola, A., Gastmeier, P., Eckweiler, D., Häussler, S., 2015. Deep transcriptome profiling of clinical *Klebsiella pneumoniae* isolates reveals strain and sequence type-specific adaptation. *Environ. Microbiol.* 17, 4690–4710. <https://doi.org/10.1111/1462-2920.13016>.
- Buckner, M.M.C., Ciusa, M.L., Piddock, L.J.V., 2018. Strategies to combat antimicrobial resistance: anti-plasmid and plasmid curing. *FEMS Microbiol. Rev.* 42, 781–804. <https://doi.org/10.1093/femsre/fuy031>.
- Burrows, L.L., 2018. The therapeutic pipeline for *Pseudomonas aeruginosa* infections. *ACS Infect. Dis.* 4, 1041–1047. <https://doi.org/10.1021/acsinfecdis.8b00112>.
- Burrus, V., 2017. Mechanisms of stabilization of integrative and conjugative elements. *Curr. Opin. Microbiol.* 38, 44–50. <https://doi.org/10.1016/J.MIB.2017.03.014>.
- Cabot, G., Florit-Mendoza, L., Sánchez-Diener, I., Zamorano, L., Oliver, A., 2018. Deciphering β -lactamase-independent β -lactam resistance evolution trajectories in *Pseudomonas aeruginosa*. *J. Antimicrob. Chemother.* 73, 3322–3331. <https://doi.org/10.1093/jac/dky364>.
- Cabot, G., López-Causapé, C., Ocampo-Sosa, A.A., Sommer, L.M., Domínguez, M.Á., Zamorano, L., Juan, C., Tubau, F., Rodríguez, C., Moyà, B., Peña, C., Martínez-Martínez, L., Plesiat, P., Oliver, A., Pomar, V., Gurgui, M., Almirante, B., Sorde, R., Larrosa, N., Aguilar, M., Martín, C., Granados, A., Fontanals, D., Calbo, E., Rodríguez-Baño, J., De Cueto, M., Navarro, M.D., Rodríguez-López, F., Torre-Cisneros, J., Casal, M., Lara, R., Natera, C., Rivero, A., Murillas, J., 2016. Deciphering the resistome of the widespread *Pseudomonas aeruginosa* sequence type 175 international high-risk clone through whole-genome sequencing. *Antimicrob. Agents Chemother.* 60, 7415–7423. <https://doi.org/10.1128/AAC.01720-16>.
- Cabot, G., Ocampo-Sosa, A.A., Domínguez, M.Á., Gago, J.F., Juan, C., Tubau, F., Rodríguez, C., Moyà, B., Peña, C., Martínez-Martínez, L., Oliver, A., Spanish Network for Research in Infectious Diseases (REIPI), 2012. Genetic markers of widespread extensively drug-resistant *Pseudomonas aeruginosa* high-risk clones. *Antimicrob. Agents Chemother.* 56, 6349–6357. <https://doi.org/10.1128/AAC.01388-12>.
- Cabot, G., Ocampo-Sosa, A.A., Tubau, F., Macía, M.D., Rodríguez, C., Moyà, B., Zamorano, L., Suárez, C., Peña, C., Martínez-Martínez, L., Oliver, A., Spanish Network for Research in Infectious Diseases (REIPI), 2011. Overexpression of AmpC and efflux pumps in *Pseudomonas aeruginosa* isolates from bloodstream infections: prevalence and impact on resistance in a Spanish multicenter study. *Antimicrob. Agents Chemother.* 55, 1906–1911. <https://doi.org/10.1128/AAC.01645-10>.
- Cantón, R., Ruiz-Garbajosa, P., 2011. Co-resistance: an opportunity for the bacteria and resistance genes. *Curr. Opin. Pharmacol.* 11, 477–485. <https://doi.org/10.1016/J.COPH.2011.07.007>.
- Carraro, N., Burrus, V., 2015. The dualistic nature of integrative and conjugative elements. *Mob. Genet. Elem.* 5, 98–102. <https://doi.org/10.1080/2159256X.2015.1102796>.
- Carraro, N., Poulin, D., Burrus, V., 2015. Replication and active partition of integrative

- and conjugative elements (ICEs) of the SXT/R391 family: the line between ICEs and conjugative plasmids is getting thinner. *PLOS Genet.* 11, e1005298. <https://doi.org/10.1371/journal.pgen.1005298>.
- Cassini, A., Högberg, L.D., Plachouras, D., Quattrocchi, A., Hoxha, A., Simonsen, G.S., Colomb-Cotinat, M., Kretzschmar, M.E., Devleeschauwer, B., Cecchini, M., Ouakrim, D.A., Oliveira, T.C., Struelens, M.J., Suetens, C., Monnet, D.L., Burden of AMR Collaborative Group, R., Mertens, K., Struyf, T., Catry, B., Latour, K., Ivanov, I.N., Dobrev, E.G., Andrašević, A.T., Soprek, S., Budimir, A., Paphitou, N., Žemlicková, H., Olsen, S.S., Sönksen, U.W., Martin, P., Ivanova, M., Lyytikäinen, O., Jalava, J., Coignard, B., Eckmanns, T., Sin, M.A., Haller, S., Daikos, G.L., Gikas, A., Tsiodras, S., Kontopoulou, F., Tóth, Á., Hajdu, Á., Guðlaugsson, Ó., Kristinnsson, K.G., Murchan, S., Burns, K., Pezzotti, P., Gagliotti, C., Dumpis, U., Liuimienė, A., Perrin, M., Borg, M.A., Greeff, S.C., de Monen, J.C., Koek, M.B., Elström, P., Zabicka, D., Deptula, A., Hryniewicz, W., Canica, M., Nogueira, P.J., Fernandes, P.A., Manageiro, V., Popescu, G.A., Serban, R.I., Schréterová, E., Litvová, S., Štefkovicová, M., Kolman, J., Klavs, I., Korošec, A., Aracil, B., Asensio, A., Pérez-Vázquez, M., Billström, H., Larsson, S., Reilly, J.S., Johnson, A., Hopkins, S., 2018. Attributable deaths and disability-adjusted life-years caused by infections with antibiotic-resistant bacteria in the EU and the European Economic Area in 2015: a population-level modelling analysis. *Lancet Infect. Dis.* 19, 56–66. [https://doi.org/10.1016/S1473-3099\(18\)30605-4](https://doi.org/10.1016/S1473-3099(18)30605-4).
- Castanheira, M., Deshpande, L.M., Costello, A., Davies, T.A., Jones, R.N., 2014. Epidemiology and carbapenem resistance mechanisms of carbapenem-non-susceptible *Pseudomonas aeruginosa* collected during 2009–11 in 14 European and Mediterranean countries. *J. Antimicrob. Chemother.* 69, 1804–1814. <https://doi.org/10.1093/jac/dku048>.
- CDC. Antibiotic Use in the United States, 2017. Progress and Opportunities. US Department of Health and Human Services, CDC, Atlanta, GA. <https://www.cdc.gov/antibiotic-use/stewardship-report/index.html>.
- Chambonier, G., Roux, L., Redelberger, D., Fadel, F., Filloux, A., Sivanesson, M., de Bentzmann, S., Bordin, C., 2016. The hybrid histidine kinase lads forms a multi-component signal transduction system with the GacS/GacA two-component system in *Pseudomonas aeruginosa*. *PLOS Genet.* 12, e1006032. <https://doi.org/10.1371/journal.pgen.1006032>.
- Chandler, C.E., Horspool, A.M., Hill, P.J., Wozniak, D.J., Schertzer, J.W., Rasko, D.A., Ernst, R.K., 2018. Genomic and phenotypic diversity among ten laboratory isolates of *Pseudomonas aeruginosa* PAO1. *J. Bacteriol.* 201, e00595–18. <https://doi.org/10.1128/JB.00595-18>.
- Chatterjee, A., Modarai, M., Naylor, N.R., Boyd, S.E., Atun, R., Barlow, J., Holmes, A.H., Johnson, A., Robotham, J.V., 2018. Quantifying drivers of antibiotic resistance in humans: a systematic review. *Lancet Infect. Dis.* 18, e368–e378. [https://doi.org/10.1016/S1473-3099\(18\)30296-2](https://doi.org/10.1016/S1473-3099(18)30296-2).
- Chávez-Jacobo, V.M., Hernández-Ramírez, K.C., Romo-Rodríguez, P., Pérez-Gallardo, R.V., Campos-García, J., Félix Gutiérrez-Corona, J., García-Merinos, J.P., Meza-Carmen, V., Silva-Sánchez, J., Ramírez-Díaz, M.I., 2018. CrpP is a novel ciprofloxacin-modifying enzyme encoded by the *Pseudomonas aeruginosa* pUM505 plasmid. *Antimicrob. Agents Chemother.* 62, e02629–e02717. <https://doi.org/10.1128/AAC.02629-17>.
- Chen, R., Déziel, E., Groleau, M.-C., Schaefer, A.L., Greenberg, E.P., 2019. Social cheating in a *Pseudomonas aeruginosa* quorum-sensing variant. *Proc. Natl. Acad. Sci. U. S. A.* 116, 7021–7026. <https://doi.org/10.1073/pnas.1819801116>.
- Chen, W., Zhang, Ya, Zhang, Yifei, Pi, Y., Gu, T., Song, L., Wang, Y., Ji, Q., 2018. CRISPR/Cas9-based genome editing in *Pseudomonas aeruginosa* and cytidine deaminase-mediated base editing in *Pseudomonas* species. *iScience* 6, 222–231. <https://doi.org/10.1016/j.isci.2018.07.024>.
- Chevalier, S., Bouffartigues, E., Bodilis, J., Maillot, O., Lesouhaitier, O., Feuillolay, M.G.J., Orange, N., Dufour, A., Cornelis, P., 2017. Structure, function and regulation of *Pseudomonas aeruginosa* porins. *FEMS Microbiol. Rev.* 41, 698–722. <https://doi.org/10.1093/femsre/fux020>.
- Chowdhury, P.R., Scott, M.J., Djordjevic, S.P., 2017. Genomic islands 1 and 2 carry multiple antibiotic resistance genes in *Pseudomonas aeruginosa* ST235, ST253, ST111 and ST175 and are globally dispersed. *J. Antimicrob. Chemother.* 72, 620–622. <https://doi.org/10.1093/jac/dkw471>.
- Clemente, J.C., Pehrsson, E.C., Blaser, M.J., Sandhu, K., Gao, Z., Wang, B., Magris, M., Hidalgo, G., Contreras, M., Noya-Alarcon, O., Lander, O., McDonald, J., Cox, M., Walter, J., Oh, P.L., Ruiz, J.F., Rodriguez, S., Shen, N., Song, S.J., Metcalf, J., Knight, R., Dantas, G., Dominguez-Bello, M.G., 2015. The microbiome of uncontacted Amerindians. *Sci. Adv.* 1, e1500183–e1500183. <https://doi.org/10.1126/sciadv.1500183>.
- Cohen, O., Gophna, U., Pupko, T., 2011. The complexity hypothesis revisited: connectivity rather than function constitutes a barrier to horizontal gene transfer. *Mol. Biol. Evol.* 28, 1481–1489. <https://doi.org/10.1093/molbev/msq333>.
- Collignon, P., Beggs, J.J., Walsh, T.R., Gandra, S., Laxminarayan, R., 2018. Anthropological and socioeconomic factors contributing to global antimicrobial resistance: a univariate and multivariable analysis. *Lancet Planet. Heal.* 2, e398–e405. [https://doi.org/10.1016/S2542-5196\(18\)30186-4](https://doi.org/10.1016/S2542-5196(18)30186-4).
- Cornaglia, G., Giamarellou, H., Rossolini, G.M., 2011. Metallo- β -lactamases: a last frontier for β -lactams? *Lancet Infect. Dis.* 11, 381–393. [https://doi.org/10.1016/S1473-3099\(11\)70056-1](https://doi.org/10.1016/S1473-3099(11)70056-1).
- Cornforth, D.M., Dees, J.L., Ibberson, C.B., Huse, H.K., Mathiesen, I.H., Kirketerp-Møller, K., Wolcott, R.D., Rumbaugh, K.P., Bjarnsholt, T., Whiteley, M., 2018. *Pseudomonas aeruginosa* transcriptome during human infection. *Proc. Natl. Acad. Sci. U. S. A.* 115, E5125–E5134. <https://doi.org/10.1073/pnas.1717525115>.
- Costello, S.E., Deshpande, L.M., Davis, A.P., Mendes, R.E., Castanheira, M., 2018. Aminoglycoside-modifying enzymes and 16S ribosomal RNA methyltransferases-encoding genes among a global collection of gram-negative isolates. *J. Glob. Antimicrob. Resist.* 16, 278–285. <https://doi.org/10.1016/J.JGAR.2018.10.020>.
- Crofts, T.S., Gasparrini, A.J., Dantas, G., 2017. Next-generation approaches to understand and combat the antibiotic resistome. *Nat. Rev. Microbiol.* 15, 422–434. <https://doi.org/10.1038/nrmicro.2017.28>.
- Cury, J., Jové, T., Touchon, M., Néron, B., Rocha, E.P., 2016. Identification and analysis of integrons and cassette arrays in bacterial genomes. *Nucleic Acids Res.* 44, 4539–4550. <https://doi.org/10.1093/nar/gkw319>.
- Cury, J., Touchon, M., Rocha, E.P.C., 2017. Integrative and conjugative elements and their hosts: composition, distribution and organization. *Nucleic Acids Res.* 45, 8943–8956. <https://doi.org/10.1093/nar/gkx607>.
- Cuzon, G., Naas, T., Nordmann, P., 2011. Functional characterization of Tn4401, a Tn3-based transposon involved in blaKPC gene mobilization. *Antimicrob. Agents Chemother.* 55, 5370–5373. <https://doi.org/10.1128/AAC.05202-11>.
- D'Costa, V.M., King, C.E., Kalan, L., Morar, M., Sung, W.W.L., Schwarz, C., Froese, D., Zazula, G., Calmels, F., Debruyne, R., Golding, G.B., Poinar, H.N., Wright, G.D., 2011. Antibiotic resistance is ancient. *Nature* 477, 457–461. <https://doi.org/10.1038/nature10388>.
- D'Costa, V.M., McGrann, K.M., Hughes, D.W., Wright, G.D., 2006. Sampling the antibiotic resistome. *Science* 311, 374–377. <https://doi.org/10.1126/science.1120800>.
- Dai, X., Zhou, D., Xiong, W., Feng, J., Luo, W., Luo, G., Wang, H., Sun, F., Zhou, X., 2016. The IncP-6 plasmid p10265-KPC from *Pseudomonas aeruginosa* carries a Novel Δ ISEc33-Associated blaKPC-2 gene cluster. *Front. Microbiol.* 7, 310. <https://doi.org/10.3389/fmicb.2016.00310>.
- Dantas, G., Sommer, M.O.A., Oluwasegun, R.D., Church, G.M., 2008. Bacteria subsisting on antibiotics. *Science* 320, 100–103. <https://doi.org/10.1126/science.1155157>.
- Darch, S.E., Simoska, O., Fitzpatrick, M., Barraza, J.P., Stevenson, K.J., Bonnez, R.T., Shear, J.B., Whiteley, M., 2018. Spatial determinants of quorum signaling in a *Pseudomonas aeruginosa* infection model. *Proc. Natl. Acad. Sci. U. S. A.* 115, 4779–4784. <https://doi.org/10.1073/pnas.1719317115>.
- Davies, J., Davies, D., 2010. Origins and evolution of antibiotic resistance. *Microbiol. Mol. Biol. Rev.* 74, 417–433. <https://doi.org/10.1128/MMBR.00016-10>.
- Davis, R.J., Jensen, S.O., Van Hal, S., Espedido, B., Gordon, A., Farhat, R., Chan, R., 2015. Whole genome sequencing in real-time investigation and management of a *Pseudomonas aeruginosa* outbreak on a Neonatal Intensive Care Unit. *Infect. Control Hosp. Epidemiol.* 36, 1058–1064. <https://doi.org/10.1017/ice.2015.133>.
- de Kraker, M.E.A., Stewardson, A.J., Harbarth, S., 2016. Will 10 million people die a year due to antimicrobial resistance by 2050? *PLoS Med.* 13, e1002184. <https://doi.org/10.1371/journal.pmed.1002184>.
- Delavat, F., Mitri, S., Pelet, S., van der Meer, J.R., 2016. Highly variable individual donor cell fates characterize robust horizontal gene transfer of an integrative and conjugative element. *Proc. Natl. Acad. Sci. U. S. A.* 113, E3375–E3383. <https://doi.org/10.1073/pnas.1604479113>.
- Delavat, F., Miyazaki, R., Carraro, N., Pradervand, N., van der Meer, J.R., 2017. The hidden life of integrative and conjugative elements. *FEMS Microbiol. Rev.* 41, 512–537. <https://doi.org/10.1093/femsre/fux008>.
- Di Pilato, V., Pollini, S., Rossolini, G.M., 2015. Tn6249, a new Tn6162 transposon derivative carrying a double-integron platform and involved with acquisition of the bla_{VIM-1} metallo- β -lactamase gene in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 59, 1583–1587. <https://doi.org/10.1128/AAC.04047-14>.
- Díaz-Orejas, R., Espinosa, M., Yeo, C.C., 2017. The importance of the expendable: toxin-antitoxin genes in plasmids and chromosomes. *Front. Microbiol.* 8, 1479. <https://doi.org/10.3389/fmicb.2017.01479>.
- Diaz, M.H., Hauser, A.R., 2010. *Pseudomonas aeruginosa* cytotoxin ExoU is injected into phagocytic cells during acute pneumonia. *Infect. Immun.* 78, 1447–1456. <https://doi.org/10.1128/IAI.01134-09>.
- diCenzo, G.C., Finan, T.M., 2017. The divided bacterial genome: structure, function, and evolution. *Microbiol. Mol. Biol. Rev.* 81. <https://doi.org/10.1128/MMBR.00019-17>.
- Diene, S.M., Rolain, J.-M., 2014. Carbapenemase genes and genetic platforms in Gram-negative bacilli: *Enterobacteriaceae*, *Pseudomonas* and *Acinetobacter* species. *Clin. Microbiol. Infect.* 20, 831–838. <https://doi.org/10.1111/1469-0691.12655>.
- Ding, Y., Teo, J.W.P., Drautz-Moses, D.I., Schuster, S.C., Givskov, M., Yang, L., 2018. Acquisition of resistance to carbapenem and macrolide-mediated quorum sensing inhibition by *Pseudomonas aeruginosa* via ICE_{Tn43716385}. *Commun. Biol.* 1, 57. <https://doi.org/10.1038/s42003-018-0064-0>.
- Domingues, S., da Silva, G.J., Nielsen, K.M., 2012. Integrons. *Mob. Genet. Elem.* 2, 211–223. <https://doi.org/10.4161/mge.22967>.
- Dortet, L., Lombardi, C., Cretin, F., Dessen, A., Filloux, A., 2018. Pore-forming activity of the *Pseudomonas aeruginosa* type III secretion system translocon alters the host epigenome. *Nat. Microbiol.* 3, 378–386. <https://doi.org/10.1038/s41564-018-0109-7>.
- Döbelmann, B., Willmann, M., Steglich, M., Bunk, B., Nübel, U., Peter, S., Neher, R.A., 2017. Rapid and consistent evolution of colistin resistance in extensively drug-resistant *Pseudomonas aeruginosa* during morbidostat culture. *Antimicrob. Agents Chemother.* 61. <https://doi.org/10.1128/AAC.00043-17>.
- Dötsch, A., Schniederjans, M., Khaledi, A., Hornischer, K., Schulz, S., Bielecka, A., Eckweiler, D., Pohl, S., Häussler, S., 2015. The *Pseudomonas aeruginosa* transcriptional landscape is shaped by environmental heterogeneity and genetic variation. *MBio* 6, e00749. <https://doi.org/10.1128/mbio.00749-15>.
- Du, D., Wang-Kan, X., Neuberger, A., van Veen, H.W., Pos, K.M., Piddock, L.J.V., Luisi, B.F., 2018. Multidrug efflux pumps: structure, function and regulation. *Nat. Rev. Microbiol.* 16, 523–539. <https://doi.org/10.1038/s41579-018-0048-6>.
- Dubois, V., Arpin, C., Noury, P., Quentin, C., 2002. Clinical strain of *Pseudomonas aeruginosa* carrying a bla(TEM-21) gene located on a chromosomal interrupted TnA type

- transposon. *Antimicrob. Agents Chemother.* 46, 3624–3626. <https://doi.org/10.1128/AAC.46.11.3624-3626.2002>.
- Dunphy, L.J., Yen, P., Papin, J.A., 2019. Integrated experimental and computational analyses reveal differential metabolic functionality in antibiotic-resistant *Pseudomonas aeruginosa*. *Cell Syst.* 8, 3–14.e3. <https://doi.org/10.1016/j.cels.2018.12.002>.
- Durão, P., Balbontín, R., Gordo, I., 2018. Evolutionary mechanisms shaping the maintenance of antibiotic resistance. *Trends Microbiol.* 26, 677–691. <https://doi.org/10.1016/j.tim.2018.01.005>.
- EFSA, 2013. Scientific opinion on carbapenem resistance in food animal ecosystems. *EFSA J.* 11, 3501. <https://doi.org/10.2903/j.efsa.2013.3501>.
- El Garch, F., Bogaerts, P., Bebrone, C., Galleni, M., Glupczynski, Y., 2011. OXA-198, an acquired carbapenem-hydrolyzing class D beta-lactamase from *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 55, 4828–4833. <https://doi.org/10.1128/AAC.00522-11>.
- England, W.E., Kim, T., Whitaker, R.J., 2018. Metapopulation structure of CRISPR-Cas immunity in *Pseudomonas aeruginosa* and its viruses. *mSystems* 3, e00075-18. <https://doi.org/10.1128/mSystems.00075-18>.
- Estrela, S., Brown, S.P., 2018. Community interactions and spatial structure shape selection on antibiotic resistant lineages. *PLOS Comput. Biol.* 14, e1006179. <https://doi.org/10.1371/journal.pcbi.1006179>.
- European Centre for Disease Prevention and Control (ECDC), 2018a. Healthcare-associated Infections Acquired in Intensive Care Units – Annual Epidemiological Report for 2016. . <https://ecdc.europa.eu/en/publications-data/healthcare-associated-infections-intensive-care-units-annual-epidemiological-0>.
- European Centre for Disease Prevention and Control (ECDC), 2018b. Surveillance of antimicrobial resistance in Europe – annual report of the european antimicrobial resistance. *Surveill. Netw.* 2017. <https://doi.org/10.2900/230516>.
- European Centre for Disease Prevention and Control (ECDC), 2018c. Annual Epidemiological Report for 2016 – Antimicrobial Consumption. . <https://ecdc.europa.eu/en/publications-data/antimicrobial-consumption-annual-epidemiological-report-2016>.
- European Centre for Disease Prevention and Control (ECDC), 2017. Antimicrobial resistance surveillance in Europe 2016 [WWW Document]. *Eur. Cent. Dis. Prev. Control*. URL <https://ecdc.europa.eu/en/publications-data/antimicrobial-resistance-surveillance-europe-2016> (accessed 21.07.19).
- European Centre for Disease Prevention Control (ECDC), 2013. Point prevalence survey reports: European long-term care facilities [WWW Document]. *Eur. Cent. Dis. Prev. Control*. URL <https://ecdc.europa.eu/en/infectious-diseases-public-health/healthcare-associated-infections-long-term-care-facilities-0> (accessed 21.07.19).
- European Centre for Disease Prevention Control (ECDC), 2014. Point prevalence survey report: Acute care hospitals [WWW Document]. *Eur. Cent. Dis. Prev. Control*. URL <https://ecdc.europa.eu/en/healthcare-associated-infections-acute-care-hospitals/surveillance-disease-data/report> (accessed 21.07.19).
- Evans, B.A., Amyes, S.G.B., 2014. OXA β -lactamases. *Clin. Microbiol. Rev.* 27, 241–263. <https://doi.org/10.1128/CMR.00117-13>.
- Fajardo, A., Hernando-Amado, S., Oliver, A., Ball, G., Filloux, A., Martínez, J.L., 2014. Characterization of a novel Zn²⁺-dependent intrinsic imipenemase from *Pseudomonas aeruginosa*. *J. Antimicrob. Chemother.* 69, 2972–2978. <https://doi.org/10.1093/jac/dku267>.
- Feng, W., Zhou, D., Wang, Q., Luo, W., Zhang, D., Sun, Q., Tong, Y., Chen, W., Sun, F., Xia, P., 2016. Dissemination of IMP-4-encoding pIMP-HZI-related plasmids among *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* in a Chinese teaching hospital. *Sci. Rep.* 6, 33419. <https://doi.org/10.1038/srep33419>.
- Fernández, L., Hancock, R.E.W., 2012. Adaptive and mutational resistance: role of porins and efflux pumps in drug resistance. *Clin. Microbiol. Rev.* 25, 661–681. <https://doi.org/10.1128/CMR.00043-12>.
- Ferreira, J.L., Gao, F.Z., Rossmann, F.M., Nans, A., Brenzinger, S., Hosseini, R., Wilson, A., Briegel, A., Thormann, K.M., Rosenthal, P.B., Beeby, M., 2019. γ -Proteobacteria eject their polar flagella under nutrient depletion, retaining flagellar motor relic structures. *PLOS Biol.* 17, e3000165. <https://doi.org/10.1371/journal.pbio.3000165>.
- Fischer, S., Klockgether, J., Morán Losada, P., Chouvarine, P., Cramer, N., Davenport, C.F., Dethlefsen, S., Dorda, M., Goemann, A., Hilker, R., Mielke, S., Schönfelder, T., Suerbaum, S., Türk, O., Woltemate, S., Wiehlmann, L., Tümmeler, B., 2016. Intracellular genome diversity of the major *Pseudomonas aeruginosa* clones C and PA14. *Environ. Microbiol. Rep.* 8, 227–234. <https://doi.org/10.1111/1758-2229.12372>.
- Flannagan, S.E., Clewell, D.B., 1991. Conjugative transfer of Tn916 in *Enterococcus faecalis*: trans activation of homologous transposons. *J. Bacteriol.* 173, 7136–7141. <https://doi.org/10.1128/jb.173.22.7136-7141.1991>.
- Fonseca, E.L., Marin, M.A., Encinas, F., Vicente, A.C.P., 2015. Full characterization of the integrative and conjugative element carrying the metallo- β -lactamase *bla*_{SPM-1} and bicyclomycin *bcr1* resistance genes found in the pandemic *Pseudomonas aeruginosa* clone SP/ST277. *J. Antimicrob. Chemother.* 70, 2547–2550. <https://doi.org/10.1093/jac/dkv152>.
- Forsberg, K.J., Patel, S., Gibson, M.K., Lauber, C.L., Knight, R., Fierer, N., Dantas, G., 2014. Bacterial phylogeny structures soil resistomes across habitats. *Nature* 509, 612–616. <https://doi.org/10.1038/nature13377>.
- Forsberg, K.J., Reyes, A., Wang, B., Selleck, E.M., Sommer, M.O.A., Dantas, G., 2012. The shared antibiotic resistome of soil bacteria and human pathogens. *Science* 337, 1107–1111. <https://doi.org/10.1126/science.1220761>.
- Freschi, L., Bertelli, C., Jeukens, J., Moore, M.P., Kukavica-Ibrulj, I., Emond-Rheault, J.-G., Hamel, J., Fothergill, J.L., Tucker, N.P., McClean, S., Klockgether, J., de Soya, A., Brinkman, F.S.L., Levesque, R.C., Winstanley, C., 2018a. Genomic characterisation of an international *Pseudomonas aeruginosa* reference panel indicates that the two major groups draw upon distinct mobile gene pools. *FEMS Microbiol. Lett.* 365, fny120. <https://doi.org/10.1093/femsle/fny120>.
- Freschi, L., Jeukens, J., Kukavica-Ibrulj, I., Boyle, B., Dupont, M.-J., Laroche, J., Larose, S., Maaroufi, H., Fothergill, J.L., Moore, M., Winsor, G.L., Aaron, S.D., Barbeau, J., Bell, S.C., Burns, J.L., Camara, M., Cantin, A., Charette, S.J., Dewar, K., Déziel, É., Grimwood, K., Hancock, R.E.W., Harrison, J.J., Heeb, S., Jelsbak, L., Jia, B., Kenna, D.T., Kidd, T.J., Klockgether, J., Lam, J.S., Lamont, L.L., Lewenza, S., Loman, N., Malouin, F., Manos, J., McArthur, A.G., McKeown, J., Milot, J., Naghra, H., Nguyen, D., Pereira, S.K., Perron, G.G., Pirnay, J.-P., Rainey, P.B., Rousseau, S., Santos, P.M., Stephenson, A., Taylor, V., Turton, J.F., Wagelchner, N., Williams, P., Thrane, S.W., Wright, G.D., Brinkman, F.S.L., Tucker, N.P., Tümmeler, B., Winstanley, C., Levesque, R.C., 2015. Clinical utilization of genomics data produced by the international *Pseudomonas aeruginosa* consortium. *Front. Microbiol.* 6, 1036. <https://doi.org/10.3389/fmicb.2015.01036>.
- Freschi, L., Vincent, A.T., Jeukens, J., Emond-Rheault, J.-G., Kukavica-Ibrulj, I., Dupont, M.-J., Charette, S.J., Boyle, B., Levesque, R.C., 2018b. The *Pseudomonas aeruginosa* pan-genome provides new insights on its population structure, horizontal gene transfer and pathogenicity. *Genome Biol. Evol.* 11, 109–120. <https://doi.org/10.1093/gbe/evy259>.
- Frost, I., Smith, W.P.J., Mitri, S., Millan, A.S., Davit, Y., Osborne, J.M., Pitt-Francis, J.M., MacLean, R.C., Foster, K.R., 2018. Cooperation, competition and antibiotic resistance in bacterial colonies. *ISME J.* 12, 1582–1593. <https://doi.org/10.1038/s41396-018-0090-4>.
- Gaillard, M., Pernet, N., Vogne, C., Hagenbüchle, O., van der Meer, J.R., 2008. Host and invader impact of transfer of the clc genomic island into *Pseudomonas aeruginosa* PAO1. *Proc. Natl. Acad. Sci. U. S. A.* 105, 7058–7063. <https://doi.org/10.1073/pnas.0801269105>.
- Gaillard, M., Vallaeys, T., Vorhölter, F.J., Minoia, M., Werlen, C., Senthil, V., Pühler, A., van der Meer, J.R., 2006. The clc element of *Pseudomonas* sp. strain B13, a genomic island with various catabolic properties. *J. Bacteriol.* 188, 1999–2013. <https://doi.org/10.1128/JB.188.5.1999-2013.2006>.
- Galata, V., Fehlmann, T., Backes, C., Keller, A., 2018. PLSDB: a resource of complete bacterial plasmids. *Nucleic Acids Res.* 47, D195–D202. <https://doi.org/10.1093/nar/gky1050>.
- Gao, B., Gallagher, T., Zhang, Y., Elbadawi-Sidhu, M., Lai, Z., Fiehn, O., Whiteson, K.L., 2018. Tracking polymicrobial metabolism in cystic fibrosis airways: *Pseudomonas aeruginosa* metabolism and physiology are influenced by rothia mucilaginosus-derived metabolites. *mSphere* 3, e00151-18. <https://doi.org/10.1128/mSphere.00151-18>.
- Garcillán-Barcia, M.P., Alvarado, A., de la Cruz, F., 2011. Identification of bacterial plasmids based on mobility and plasmid population biology. *FEMS Microbiol. Rev.* 35, 936–956. <https://doi.org/10.1111/j.1574-6976.2011.00291.x>.
- Garday, J.L., Loman, N.J., 2017. Towards a genomics-informed, real-time, global pathogen surveillance system. *Nat. Rev. Genet.* 19, 9–20. <https://doi.org/10.1038/nrg.2017.88>.
- Garza-Ramos, U., Barrios, H., Reyna-Flores, F., Tamayo-Legorreta, E., Catalan-Najera, J.C., Morfin-Otero, R., Rodríguez-Noriega, E., Volkow, P., Cornejo-Juarez, P., González, A., Gaytan-Martínez, J., del Rocio González-Martínez, M., Vazquez-Farías, M., Silva-Sánchez, J., 2015. Widespread of ESB- and carbapenemase GES-type genes on carbapenem-resistant *Pseudomonas aeruginosa* clinical isolates: a multicenter study in Mexican hospitals. *Diagn. Microbiol. Infect. Dis.* 81, 135–137. <https://doi.org/10.1016/j.diagmicrobio.2014.09.029>.
- Gellatly, S.L., Hancock, R.E.W., 2013. *Pseudomonas aeruginosa*: new insights into pathogenesis and host defenses. *Pathog. Dis.* 67, 159–173. <https://doi.org/10.1111/2049-632X.12033>.
- Ghaly, T.M., Gillings, M.R., 2018. Mobile DNAs as ecologically and evolutionarily independent units of life. *Trends Microbiol.* 26, 904–912. <https://doi.org/10.1016/j.tim.2018.05.008>.
- Giani, T., Arena, F., Pollini, S., Di Pilato, V., D'Andrea, M.M., Henrici De Angelis, L., Bassetti, M., Rossolini, G.M., Vismara, C., Luzzaro, F., Cavallo, R., Dusi, P.A., Pagani, E., Sarti, M., Farina, C., Rigoli, R., Scarparo, C., Pecile, P., Cusi, M.G., Mencacci, A., Manso, E., Spanu, T., Labonia, M., Tassi, V., Amato, G., Stefani, S., Giraldi, C., Rassu, M., 2018. Italian nationwide survey on *Pseudomonas aeruginosa* from invasive infections: activity of ceftolozane/tazobactam and comparators, and molecular epidemiology of carbapenemase producers. *J. Antimicrob. Chemother.* 73, 664–671. <https://doi.org/10.1093/jac/dkx453>.
- Gifford, D.R., Furió, V., Pakpou, A., Vogwill, T., Oliver, A., MacLean, R.C., 2018. Identifying and exploiting genes that potentiate the evolution of antibiotic resistance. *Nat. Ecol. Evol.* 2, 1033–1039. <https://doi.org/10.1038/s41559-018-0547-x>.
- Gillings, M.R., Paulsen, I.T., Tetu, S.G., 2017. Genomics and the evolution of antibiotic resistance. *Ann. N. Y. Acad. Sci.* 1388, 92–107. <https://doi.org/10.1111/nyas.13268>.
- Girlich, D., Naas, T., Nordmann, P., 2004. Biochemical characterization of the naturally occurring oxacillinase OXA-50 of *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 48, 2043–2048. <https://doi.org/10.1128/AAC.48.6.2043-2048.2004>.
- Gogarten, J.P., Townsend, J.P., 2005. Horizontal gene transfer, genome innovation and evolution. *Nat. Rev. Microbiol.* 3, 679–687. <https://doi.org/10.1038/nrmicro1204>.
- Gonzalez, M.R., Ducret, V., Fleuchot, B., Jafari, P., Raffoul, W., Applegate, L.A., Que, Y.-A., Perron, K., 2018. Transcriptome analysis of *Pseudomonas aeruginosa* cultured in human burn wound exudates. *Front. Cell. Infect. Microbiol.* 8, 39. <https://doi.org/10.3389/fcimb.2018.00039>.
- Good, B.H., McDonald, M.J., Barrick, J.E., Lenski, R.E., Desai, M.M., 2017. The dynamics of molecular evolution over 60,000 generations. *Nature* 551, 45–50. <https://doi.org/10.1038/nature24287>.
- Granato, E.T., Ziegenhain, C., Marvig, R.L., Kümmerli, R., 2018. Low spatial structure and selection against secreted virulence factors attenuates pathogenicity in *Pseudomonas aeruginosa*. *ISME J.* 12, 2907–2918. <https://doi.org/10.1038/s41396-018-0231-9>.
- Guédon, G., Libante, V., Coluzzi, C., Payot, S., Leblond-Bourget, N., 2017. The obscure world of integrative and mobilizable elements, highly widespread elements that pirate bacterial conjugative systems. *Genes (Basel)* 8, 337. <https://doi.org/10.3390/genes8080337>.

- genes8110337.
- Guglielmini, J., de la Cruz, F., Rocha, E.P.C., 2013. Evolution of conjugation and Type IV secretion systems. *Mol. Biol. Evol.* 30, 315–331. <https://doi.org/10.1093/molbev/mst221>.
- Guglielmini, J., Néron, B., Abby, S.S., Garcillán-Barcia, M.P., de la Cruz, F., Rocha, E.P.C., 2014. Key components of the eight classes of type IV secretion systems involved in bacterial conjugation or protein secretion. *Nucleic Acids Res.* 42, 5715–5727. <https://doi.org/10.1093/nar/gku194>.
- Guglielmini, J., Quintais, L., Garcillán-Barcia, M.P., de la Cruz, F., Rocha, E.P.C., 2011. The repertoire of ICE in prokaryotes underscores the unity, diversity, and ubiquity of conjugation. *PLoS Genet.* 7, e1002222. <https://doi.org/10.1371/journal.pgen.1002222>.
- Halary, S., Leigh, J.W., Cheaib, B., Lopez, P., Baptiste, E., 2010. Network analyses structure genetic diversity in independent genetic worlds. *Proc. Natl. Acad. Sci. U. S. A.* 107, 127–132. <https://doi.org/10.1073/pnas.0908978107>.
- Hall, J.P.J., Brockhurst, M.A., Harrison, E., 2017. Sampling the mobile gene pool: innovation via horizontal gene transfer in bacteria. *Philos. Trans. R. Soc. B: Biol. Sci.* 372, 20160424. <https://doi.org/10.1098/rstb.2016.0424>.
- Han, M.-L., Zhu, Y., Creek, D.J., Lin, Y.-W., Gutu, A.D., Hertzog, P., Purcell, T., Shen, H.-H., Moskowitz, S.M., Velkov, T., Li, J., 2019. Comparative metabolomics and transcriptomics reveal multiple pathways associated with polymyxin killing in *Pseudomonas aeruginosa*. *mSystems* 4. <https://doi.org/10.1128/mSystems.00149-18>.
- Hancock, R.E., 1998. Resistance mechanisms in *Pseudomonas aeruginosa* and other non-fermentative gram-negative bacteria. *Clin. Infect. Dis.* 27, S93–S99. <https://doi.org/10.1086/514909>.
- Harrison, E., Brockhurst, M.A., 2012. Plasmid-mediated horizontal gene transfer is a coevolutionary process. *Trends Microbiol.* 20, 262–267. <https://doi.org/10.1016/j.tim.2012.04.003>.
- Harrison, E., Guymer, D., Spiers, A.J., Paterson, S., Brockhurst, M.A., 2015. Parallel compensatory evolution stabilizes plasmids across the parasitism-mutualism continuum. *Curr. Biol.* 25, 2034–2039. <https://doi.org/10.1016/j.cub.2015.06.024>.
- Harrison, E.M., Carter, M.E.K., Luck, S., Ou, H.-Y., He, X., Deng, Z., O'Callaghan, C., Kadioglu, A., Rajakumar, K., 2010. Pathogenicity islands PAPI-1 and PAPI-2 contribute individually and synergistically to the virulence of *Pseudomonas aeruginosa* strain PA14. *Infect. Immun.* 78, 1437–1446. <https://doi.org/10.1128/IAI.00621-09>.
- Haskett, T.L., Terpolilli, J.J., Bekuma, A., O'Hara, G.W., Sullivan, J.T., Wang, P., Ronson, C.W., Ramsay, J.P., 2016. Assembly and transfer of tripartite integrative and conjugative genetic elements. *Proc. Natl. Acad. Sci. U. S. A.* 113, 12268–12273. <https://doi.org/10.1073/pnas.1613358113>.
- He, J., Baldini, R.L., Déziel, E., Saucier, M., Zhang, Q., Liberati, N.T., Lee, D., Urbach, J., Goodman, H.M., Rahme, L.G., 2004. The broad host range pathogen *Pseudomonas aeruginosa* strain PA14 carries two pathogenicity islands harboring plant and animal virulence genes. *Proc. Natl. Acad. Sci. U. S. A.* 101, 2530–2535. <https://doi.org/10.1073/PNAS.0304622101>.
- Hilker, R., Munder, A., Klockgether, J., Losada, P.M., Chouvarine, P., Cramer, N., Davenport, C.F., Dethlefsen, S., Fischer, S., Peng, H., Schönfelder, T., Türk, O., Wiehmann, L., Wöbeling, F., Gulbins, E., Goemann, A., Tümmeler, B., 2015. Interclonal gradient of virulence in the *P. pseudomonas aeruginosa* pangenome from disease and environment. *Environ. Microbiol.* 17, 29–46. <https://doi.org/10.1111/1462-2920.12606>.
- Holmes, A.H., Moore, L.S.P., Sundsfjord, A., Steinbakk, M., Regmi, S., Karkey, A., Guerin, P.J., Piddock, L.J.V., 2016. Understanding the mechanisms and drivers of antimicrobial resistance. *Lancet* 387, 176–187. [https://doi.org/10.1016/S0140-6736\(15\)00473-0](https://doi.org/10.1016/S0140-6736(15)00473-0).
- Hong, J.S., Yoon, E.J., Lee, H., Jeong, S.H., Lee, K., 2016. Clonal dissemination of *Pseudomonas aeruginosa* sequence type 235 isolates carrying blaIMP-6 and emergence of blaGES-24 and blaIMP-10 on novel genomic islands PAGI-15 and -16 in South Korea. *Antimicrob. Agents Chemother.* 60, 7216–7223. <https://doi.org/10.1128/AAC.01601-16>.
- Hooper, S.D., Mavromatis, K., Kyrpides, N.C., 2009. Microbial co-habitation and lateral gene transfer: what transposases can tell us. *Genome Biol.* 10, R45. <https://doi.org/10.1186/gb-2009-10-4-r45>.
- Hopkins, K.L., Findlay, J., Mustafa, N., Pike, R., Parsons, H., Wright, L., Woodford, N., Meunier, D., 2016. SPM-1 metallo- β -lactamase-producing *Pseudomonas aeruginosa* ST277 in the UK. *J. Med. Microbiol.* 65, 696–697. <https://doi.org/10.1099/jmm.0.000269>.
- Hornischer, K., Khaledi, A., Pohl, S., Schniederjans, M., Pezoldt, L., Casilag, F., Muthukumarasamy, U., Bruchmann, S., Thöming, J., Kordes, A., Häussler, S., 2018. BACTOME—a reference database to explore the sequence- and gene expression-variation landscape of *Pseudomonas aeruginosa* clinical isolates. *Nucleic Acids Res.* 47, D716–D720. <https://doi.org/10.1093/nar/gky895>.
- Hülter, N., Ilhan, J., Wein, T., Kadibalban, A.S., Hammerschmidt, K., Dagan, T., 2017. An evolutionary perspective on plasmid lifestyle modes. *Curr. Opin. Microbiol.* 38, 74–80. <https://doi.org/10.1016/j.mib.2017.05.001>.
- Ilhan, J., Kupczok, A., Woehle, C., Wein, T., Hülter, N.F., Rosenstiel, P., Landan, G., Mizrahi, I., Dagan, T., 2018. Segregational drift and the interplay between plasmid copy number and evolvability. *Mol. Biol. Evol.* 36, 472–486. <https://doi.org/10.1093/molbev/msy225>.
- Imamovic, L., Ellabaan, M.M.H., Dantas Machado, A.M., Citterio, L., Wulff, T., Molin, S., Krogh Johansen, H., Sommer, M.O.A., 2018. Drug-driven phenotypic convergence supports rational treatment strategies of chronic infections. *Cell* 172, 121–134.e14. <https://doi.org/10.1016/j.cell.2017.12.012>.
- Ito, R., Mustapha, M.M., Tomich, A.D., Callaghan, J.D., McElheny, C.L., Mettett, R.T., Shanks, R.M.Q., Sluis-Cremer, N., Doi, Y., 2017a. Widespread fosfomycin resistance in gram-negative bacteria attributable to the chromosomal fosA gene. *MBio* 8, e00749–e00817. <https://doi.org/10.1128/MBIO.00749-17>.
- Ito, R., Tomich, A.D., McElheny, C.L., Mettett, R.T., Sluis-Cremer, N., Doi, Y., 2017b. Inhibition of fosfomycin resistance protein fosA by phosphonofomate (foscarnet) in multidrug-resistant gram-negative pathogens. *Antimicrob. Agents Chemother.* 61, e01424–17. <https://doi.org/10.1128/AAC.01424-17>.
- Jackson, J., Chen, C., Buising, K., 2013. Aminoglycosides: how should we use them in the 21st century? *Curr. Opin. Infect. Dis.* 26, 516–525. <https://doi.org/10.1097/QCO.000000000000012>.
- Jani, M., Mathee, K., Azad, R.K., 2016. Identification of novel genomic islands in Liverpool epidemic strain of *Pseudomonas aeruginosa* using segmentation and clustering. *Front. Microbiol.* 7, 1210. <https://doi.org/10.3389/fmicb.2016.01210>.
- Jeanot, K., Danassie, M., Triponney, P., Bour, M., Gueudet, T., Beyrouthy, R., Bonnet, R., Plésiat, P., 2018. A novel IncQ plasmid carrying gene blaCTX-M-3 in *Pseudomonas aeruginosa*. *J. Antimicrob. Chemother.* 74, 823–825. <https://doi.org/10.1093/jac/dky454>.
- Jensen, P.A., Zhu, Z., van Opijnen, T., 2017. Antibiotics disrupt coordination between transcriptional and phenotypic stress responses in pathogenic bacteria. *Cell Rep.* 20, 1705–1716. <https://doi.org/10.1016/j.celrep.2017.07.062>.
- Jesus, T.F., Ribeiro-Gonçalves, B., Silva, D.N., Bortolao, V., Ramirez, M., Carriço, J.A., 2019. Plasmid ATLAS: plasmid visual analytics and identification in high-throughput sequencing data. *Nucleic Acids Res.* 47, D188–D194. <https://doi.org/10.1093/nar/gky107>.
- Jiang, X., Ellabaan, M.M.H., Charusanti, P., Munck, C., Blin, K., Tong, Y., Weber, T., Sommer, M.O.A., Lee, S.Y., 2017. Dissemination of antibiotic resistance genes from antibiotic producers to pathogens. *Nat. Commun.* 8, 15784. <https://doi.org/10.1038/ncomms15784>.
- Jochumsen, N., Marvig, R.L., Damkjaer, S., Jensen, R.L., Paulander, W., Molin, S., Jelsbak, L., Folkesson, A., 2016. The evolution of antimicrobial peptide resistance in *Pseudomonas aeruginosa* is shaped by strong epistatic interactions. *Nat. Commun.* 7, 13002. <https://doi.org/10.1038/ncomms13002>.
- Johnson, C.M., Grossman, A.D., 2015. Integrative and conjugative elements (ICEs): what they do and how they work. *Annu. Rev. Genet.* 49, 577–601. <https://doi.org/10.1146/annurev-genet-112414-055018>.
- Jolley, K.A., Bray, J.E., Maiden, M.C.J., 2018. Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res.* 3, 124. <https://doi.org/10.12688/wellcomeopenres.14826.1>.
- Juan, C., Moya, B., Perez, J.L., Oliver, A., 2006. Stepwise upregulation of the *Pseudomonas aeruginosa* chromosomal cephalosporinase conferring high-level-lactam resistance involves three AmpD homologues. *Antimicrob. Agents Chemother.* 50, 1780–1787. <https://doi.org/10.1128/AAC.50.5.1780-1787.2006>.
- Juan, C., Peña, C., Oliver, A., 2017a. Host and pathogen biomarkers for severe *Pseudomonas aeruginosa* infections. *J. Infect. Dis.* 215, S44–S51. <https://doi.org/10.1093/infdis/jiw299>.
- Juan, C., Torrens, G., González-Nicolau, M., Oliver, A., 2017b. Diversity and regulation of intrinsic β -lactamases from non-fermenting and other Gram-negative opportunistic pathogens. *FEMS Microbiol. Rev.* 41, 781–815. <https://doi.org/10.1093/femsre/fux043>.
- Juan, C., Zamorano, L., Mena, A., Alberti, S., Perez, J.L., Oliver, A., 2010. Metallo- β -lactamase-producing *Pseudomonas putida* as a reservoir of multidrug resistance elements that can be transferred to successful *Pseudomonas aeruginosa* clones. *J. Antimicrob. Chemother.* 65, 474–478. <https://doi.org/10.1093/jac/dkp491>.
- Jutkina, J., Marathe, N.P., Flach, C.-F., Larsson, D.G.J., 2018. Antibiotics and common antibacterial biocides stimulate horizontal transfer of resistance at low concentrations. *Sci. Total Environ.* 616–617, 172–178. <https://doi.org/10.1016/j.scitotenv.2017.10.312>.
- Karkman, A., Do, T.T., Walsh, F., Virta, M.P.J., 2018. Antibiotic-resistance genes in waste water. *Trends Microbiol.* 26, 220–228. <https://doi.org/10.1016/j.tim.2017.09.005>.
- Karlin, S., 2001. Detecting anomalous gene clusters and pathogenicity islands in diverse bacterial genomes. *Trends Microbiol.* 9, 335–343. [https://doi.org/10.1016/S0966-842X\(01\)02079-0](https://doi.org/10.1016/S0966-842X(01)02079-0).
- Kassen, R., 2002. The experimental evolution of specialists, generalists, and the maintenance of diversity. *J. Evol. Biol.* 15, 173–190. <https://doi.org/10.1046/j.1420-9101.2002.00377.x>.
- Khaledi, A., Schniederjans, M., Pohl, S., Rainer, R., Bodenhofer, U., Xia, B., Klawonn, F., Bruchmann, S., Preusse, M., Eckweiler, D., Dötsch, A., Häussler, S., 2016. Transcriptome profiling of antimicrobial resistance in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 60, 4722–4733. <https://doi.org/10.1128/AAC.00075-16>.
- Klein, E.Y., Van Boeckel, T.P., Martinez, E.M., Pant, S., Gandra, S., Levin, S.A., Goossens, H., Laxminarayan, R., 2018. Global increase and geographic convergence in antibiotic consumption between 2000 and 2015. *Proc. Natl. Acad. Sci. U. S. A.* 115, E3463–E3470. <https://doi.org/10.1073/pnas.1717295115>.
- Klockgether, J., Cramer, N., Wiehmann, L., Davenport, C.F., Tümmeler, B., 2011. *Pseudomonas aeruginosa* genomic structure and diversity. *Front. Microbiol.* 2, 150. <https://doi.org/10.3389/fmicb.2011.00150>.
- Kloesges, T., Popa, O., Martin, W., Dagan, T., 2011. Networks of gene sharing among 329 proteobacterial genomes reveal differences in lateral gene transfer frequency at different phylogenetic depths. *Mol. Biol. Evol.* 28, 1057–1074. <https://doi.org/10.1093/molbev/msq297>.
- Knapp, C.W., Doling, J., Ehler, P.A.I., Graham, D.W., 2010. Evidence of increasing antibiotic resistance gene abundances in archived soils since 1940. *Environ. Sci.*

- Technol. 44, 580–587. <https://doi.org/10.1021/es901221x>.
- Knopp, M., Andersson, D.I., 2018. Predictable phenotypes of antibiotic resistance mutations. *MBio* 9, e00770–e00818. <https://doi.org/10.1128/mBio.00770-18>.
- Knöppel, A., Näsvall, J., Andersson, D.I., 2017. Evolution of antibiotic resistance without antibiotic exposure. *Antimicrob. Agents Chemother.* 61, e01495–17. <https://doi.org/10.1128/AAC.01495-17>.
- Koepfen, K., Hampton, T.H., Jarek, M., Scharfe, M., Gerber, S.A., Mielcarz, D.W., Demers, E.G., Dolben, E.L., Hammond, J.H., Hogan, D.A., Stanton, B.A., 2016. A novel mechanism of host–pathogen interaction through sRNA in bacterial outer membrane vesicles. *PLOS Pathog.* 12, e1005672. <https://doi.org/10.1371/journal.ppat.1005672>.
- Kollaran, A.M., Joge, S., Kotian, H.S., Badal, D., Prakash, D., Mishra, A., Varma, M., Singh, V., 2019. Context-specific requirement of forty-four two-component loci in *Pseudomonas aeruginosa* swarming. *iScience* 13, 305–317. <https://doi.org/10.1016/j.isci.2019.02.028>.
- Kos, V.N., Déraspe, M., McLaughlin, R.E., Whiteaker, J.D., Roy, P.H., Alm, R.A., Corbeil, J., Gardner, H., 2015. The resistome of *Pseudomonas aeruginosa* in relationship to phenotypic susceptibility. *Antimicrob. Agents Chemother.* 59, 427–436. <https://doi.org/10.1128/AAC.03954-14>.
- Kottara, A., Hall, J.P.J., Harrison, E., Brockhurst, M.A., 2018. Variable plasmid fitness effects and mobile genetic element dynamics across *Pseudomonas* species. *FEMS Microbiol. Ecol.* 94, fix172. <https://doi.org/10.1093/femsec/fix172>.
- Krawczyk, P.S., Lipinski, L., Dziembowski, A., 2018. PlasFlow: predicting plasmid sequences in metagenomic data using genome signatures. *Nucleic Acids Res.* 46, e35. <https://doi.org/10.1093/nar/gkx1321>.
- Kulasekara, B.R., Kulasekara, H.D., Wolfgang, M.C., Stevens, L., Frank, D.W., Lory, S., 2006. Acquisition and evolution of the *exoU* locus in *Pseudomonas aeruginosa*. *J. Bacteriol.* 188, 4037–4050. <https://doi.org/10.1128/JB.02000-05>.
- Kung, V.L., Ozer, E.A., Hauser, A.R., 2010. The accessory genome of *Pseudomonas aeruginosa*. *Microbiol. Mol. Biol. Rev.* 74, 621–641. <https://doi.org/10.1128/MMBR.00027-10>.
- Langille, M.G.I., Hsiao, W.W.L., Brinkman, F.S.L., 2010. Detecting genomic islands using bioinformatics approaches. *Nat. Rev. Microbiol.* 8, 373–382. <https://doi.org/10.1038/nrmicro2350>.
- Lanza, V.F., de Toro, M., Garcillán-Barcia, M.P., Mora, A., Blanco, J., Coque, T.M., de la Cruz, F., 2014. Plasmid flux in *Escherichia coli* ST131 sublineages, analyzed by Plasmid Constellation Network (PLACNET), a new method for plasmid reconstruction from whole genome sequences. *PLoS Genet.* 10, e1004766. <https://doi.org/10.1371/journal.pgen.1004766>.
- Larbig, K.D., Christmann, A., Johann, A., Klockgether, J., Hartsch, T., Merkl, R., Wiehlmann, L., Fritz, H.-J., Tümmeler, B., 2002. Gene islands integrated into tRNA(Gly) genes confer genome diversity on a *Pseudomonas aeruginosa* clone. *J. Bacteriol.* 184, 6665–6680. <https://doi.org/10.1128/jb.184.23.6665-6680.2002>.
- Lassek, C., Berger, A., Zühlke, D., Wittmann, C., Riedel, K., 2016. Proteome and carbon flux analysis of *Pseudomonas aeruginosa* clinical isolates from different infection sites. *Proteomics* 16, 1381–1385. <https://doi.org/10.1002/pmic.201500228>.
- Laudy, A.E., Róg, P., Smolińska-Król, K., Ćmiel, M., Słoczyńska, A., Patzer, J., Dzierżanowska, D., Wolinowska, R., Starościak, B., Tyski, S., 2017. Prevalence of ESBL-producing *Pseudomonas aeruginosa* isolates in Warsaw, Poland, detected by various phenotypic and genotypic methods. *PLoS One* 12, e0180121. <https://doi.org/10.1371/journal.pone.0180121>.
- Lee, J.-Y., Park, Y.K., Chung, E.S., Na, I.Y., Ko, K.S., 2016. Evolved resistance to colistin and its loss due to genetic reversion in *Pseudomonas aeruginosa*. *Sci. Rep.* 6, 25543. <https://doi.org/10.1038/srep25543>.
- Lee, J., Zhang, L., 2015. The hierarchy quorum sensing network in *Pseudomonas aeruginosa*. *Protein Cell* 6, 26–41. <https://doi.org/10.1007/s13238-014-0100-x>.
- Lehtinen, S., Blanquart, F., Croucher, N.J., Turner, P., Lipsitch, M., Fraser, C., 2017. Evolution of antibiotic resistance is linked to any genetic mechanism affecting bacterial duration of carriage. *Proc. Natl. Acad. Sci. U. S. A.* 114, 1075–1080. <https://doi.org/10.1073/pnas.1617849114>.
- Levin-Reisman, I., Ronin, I., Gefen, O., Braniss, I., Shoshani, N., Balaban, N.Q., 2017. Antibiotic tolerance facilitates the evolution of resistance. *Science* 355, 826–830. <https://doi.org/10.1126/science.aaj2191>.
- Li, H., Toleman, M.A., Bennett, P.M., Jones, R.N., Walsh, T.R., 2008. Complete Sequence of p07-406, a 24,179-base-pair plasmid harboring the blaVIM-7 metallo-beta-lactamase gene in a *Pseudomonas aeruginosa* isolate from the United States. *Antimicrob. Agents Chemother.* 52, 3099–3105. <https://doi.org/10.1128/AAC.01093-07>.
- Li, R., Fang, L., Tan, S., Yu, M., Li, X., He, S., Wei, Y., Li, G., Jiang, J., Wu, M., 2016. Type I CRISPR-Cas targets endogenous genes and regulates virulence to evade mammalian host immunity. *Cell Res.* 26, 1273–1287. <https://doi.org/10.1038/cr.2016.135>.
- Li, X.-Z., Plésiat, P., Nikaido, H., 2015. The challenge of efflux-mediated antibiotic resistance in Gram-negative bacteria. *Clin. Microbiol. Rev.* 28, 337–418. <https://doi.org/10.1128/CMR.00117-14>.
- Limoli, D.H., Whitfield, G.B., Kitao, T., Ivey, M.L., Davis, M.R., Grahl, N., Hogan, D.A., Rahme, L.G., Howell, P.L., O'Toole, G.A., Goldberg, J.B., 2017. *Pseudomonas aeruginosa* alginate overproduction promotes coexistence with *Staphylococcus aureus* in a model of cystic fibrosis respiratory infection. *MBio* 8, e00186–e00217. <https://doi.org/10.1128/mBio.00186-17>.
- Ling, L.L., Schneider, T., Peoples, A.J., Sporing, A.L., Engels, I., Conlon, B.P., Mueller, A., Schäberle, T.F., Hughes, D.E., Epstein, S., Jones, M., Lazarides, L., Steadman, V.A., Cohen, D.R., Felix, C.R., Fetterman, K.A., Millett, W.P., Nitti, A.G., Zullo, A.M., Chen, C., Lewis, K., 2015. A new antibiotic kills pathogens without detectable resistance. *Nature* 517, 455–459. <https://doi.org/10.1038/nature14098>.
- Lister, P.D., Wolter, D.J., Hanson, N.D., 2009. Antibacterial-resistant *Pseudomonas aeruginosa*: clinical impact and complex regulation of chromosomally encoded resistance mechanisms. *Clin. Microbiol. Rev.* 22, 582–610. <https://doi.org/10.1128/CMR.00040-09>.
- Liu, J., Yang, L., Chen, D., Peters, B.M., Li, L., Li, B., Xu, Z., Shirliff, M.E., 2018. Complete sequence of pBM413, a novel multidrug resistance megaplasmid carrying qnrVC6 and blaIMP-45 from *Pseudomonas aeruginosa*. *Int. J. Antimicrob. Agents* 51, 145–150. <https://doi.org/10.1016/j.ijantimicag.2017.09.008>.
- Liu, M., Li, X., Xie, Y., Bi, D., Sun, J., Li, J., Tai, C., Deng, Z., Ou, H.-Y., 2018. ICEberg 2.0: an updated database of bacterial integrative and conjugative elements. *Nucleic Acids Res.* 47, D660–D665. <https://doi.org/10.1093/nar/gky1123>.
- Livemore, D.M., 2018. The 2018 Garrod Lecture: preparing for the Black Swans of resistance. *J. Antimicrob. Chemother.* 73, 2907–2915. <https://doi.org/10.1093/jac/dky265>.
- Livemore, D.M., 2001. Of *Pseudomonas*, porins, pumps and carbapenems. *J. Antimicrob. Chemother.* 47, 247–250. <https://doi.org/10.1093/jac/47.3.247>.
- Livemore, D.M., 1992. Interplay of impermeability and chromosomal beta-lactamase activity in imipenem-resistant *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 36, 2046–2048. <https://doi.org/10.1128/aac.36.9.2046>.
- Livingston, S., Cadnum, J.L., Gestrich, S., Jenson, A.L., Donskey, C.J., 2018. Efficacy of automated disinfection with ozonated water in reducing sink drainage system colonization with *Pseudomonas* species and *Candida auris*. *Infect. Control Hosp. Epidemiol.* 39, 1497–1498. <https://doi.org/10.1017/ice.2018.176>.
- Lofte-Eaton, W., Bashford, K., Quinn, H., Dong, K., Millstein, J., Hunter, S., Thomason, M.K., Merrikk, H., Ponciano, J.M., Top, E.M., 2017. Compensatory mutations improve general permissiveness to antibiotic resistance plasmids. *Nat. Ecol. Evol.* 1, 1354–1363. <https://doi.org/10.1038/s41559-017-0243-2>.
- Lofte-Eaton, W., Yano, H., Burleigh, S., Simmons, R.S., Hughes, J.M., Rogers, L.M., Hunter, S.S., Settles, M.L., Forney, L.J., Ponciano, J.M., Top, E.M., 2016. Evolutionary paths that expand plasmid host-range: implications for spread of antibiotic resistance. *Mol. Biol. Evol.* 33, 885–897. <https://doi.org/10.1093/molbev/msv339>.
- Lopatkin, A.J., Meredith, H.R., Srimani, J.K., Pfeiffer, C., Durrett, R., You, L., 2017. Persistence and reversal of plasmid-mediated antibiotic resistance. *Nat. Commun.* 8, 1689. <https://doi.org/10.1038/s41467-017-01532-1>.
- López-Causapé, C., Cabot, G., del Barrio-Tofiño, E., Oliver, A., 2018. The versatile mutational resistome of *Pseudomonas aeruginosa*. *Front. Microbiol.* 9, 685. <https://doi.org/10.3389/fmicb.2018.00685>.
- Lynch, J., Zhanel, G., Clark, N., 2017. Emergence of antimicrobial resistance among *Pseudomonas aeruginosa*: implications for therapy. *Semin. Respir. Crit. Care Med.* 38, 326–345. <https://doi.org/10.1055/s-0037-1602583>.
- Maatallah, M., Cheriaa, J., Backrouf, A., Iversen, A., Grundmann, H., Do, T., Lanotte, P., Mastouri, M., Elghmati, M.S., Rojo, F., Mejd, S., Giske, C.G., 2011. Population structure of *Pseudomonas aeruginosa* from five mediterranean countries: evidence for frequent recombination and epidemic occurrence of CC235. *PLoS One* 6, e25617. <https://doi.org/10.1371/journal.pone.0025617>.
- Magiorakos, A.-P., Srinivasan, A., Carey, R.B., Carmeli, Y., Falagas, M.E., Giske, C.G., Harbarth, S., Hindler, J.F., Kahlmeter, G., Olsson-Liljequist, B., Paterson, D.L., Rice, L.B., Stelling, J., Struelens, M.J., Vatopoulos, A., Weber, J.T., Monnet, D.L., 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin. Microbiol. Infect.* 18, 268–281. <https://doi.org/10.1111/j.1469-0691.2011.03570.x>.
- Maier, L., Pruteanu, M., Kuhn, M., Zeller, G., Telzerow, A., Anderson, E.E., Brochado, A.R., Fernandez, K.C., Dose, H., Mori, H., Patil, K.R., Bork, P., Typas, A., 2018. Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature* 555, 623–628. <https://doi.org/10.1038/nature25979>.
- Martinez, E., Marquez, C., Ingold, A., Merlino, J., Djordjevic, S.P., Stokes, H.W., Chowdhury, P.R., 2012. Diverse mobilized class 1 integrons are common in the chromosomes of pathogenic *Pseudomonas aeruginosa* clinical isolates. *Antimicrob. Agents Chemother.* 56, 2169–2172. <https://doi.org/10.1128/AAC.06048-11>.
- Martinez, J.L., Sánchez, M.B., Martínez-Solano, L., Hernández, A., Garmendia, L., Fajardo, A., Alvarez-Ortega, C., 2009. Functional role of bacterial multidrug efflux pumps in microbial natural ecosystems. *FEMS Microbiol. Rev.* 33, 430–449. <https://doi.org/10.1111/j.1574-6976.2008.00157.x>.
- Mathee, K., Narasimhan, G., Valdes, C., Qiu, X., Matewisch, J.M., Koehrsen, M., Rokas, A., Yandava, C.N., Engels, R., Zeng, E., Olavarietta, R., Doud, M., Smith, R.S., Montgomery, P., White, J.R., Godfrey, P.A., Kodira, C., Birren, B., Galagan, J.E., Lory, S., 2008. Dynamics of *Pseudomonas aeruginosa* genome evolution. *Proc. Natl. Acad. Sci. U. S. A.* 105, 3100–3105. <https://doi.org/10.1073/pnas.0711982105>.
- McGeary, R.P., Tan, D.T., Schenk, G., 2017. Progress toward inhibitors of metallo-beta-lactamases. *Future Med. Chem.* 9, 673–691. <https://doi.org/10.4155/fmc-2017-0007>.
- McInerney, J.O., McNally, A., O'Connell, M.J., 2017. Why prokaryotes have pangenomes. *Nat. Microbiol.* 2, 17040. <https://doi.org/10.1038/nmicrobiol.2017.40>.
- Melnik, A.H., McCloskey, N., Hinz, A.J., Dettman, J., Kassen, R., 2017. Evolution of cost-free resistance under fluctuating drug selection in *Pseudomonas aeruginosa*. *mSphere* 2, e00158-17. <https://doi.org/10.1128/mSphere.00158-17>.
- Melnik, A.H., Wong, A., Kassen, R., 2015. The fitness costs of antibiotic resistance mutations. *Evol. Appl.* 8, 273–283. <https://doi.org/10.1111/eva.12196>.
- Meredith, H.R., Andreani, V., Ma, H.R., Lopatkin, A.J., Lee, A.J., Anderson, D.J., Batt, G., You, L., 2018. Applying ecological resistance and resilience to dissect bacterial antibiotic responses. *Sci. Adv.* 4, eaau1873. <https://doi.org/10.1126/sciadv.aau1873>.

- Meunier, D., Doumith, M., Findlay, J., Mustafa, N., Mallard, K., Anson, J., Panagea, S., Pike, R., Wright, L., Woodford, N., Hopkins, K.L., 2016. Carbapenem resistance mediated by bla_{OXA-181} in *Pseudomonas aeruginosa*. J. Antimicrob. Chemother. 71, 2056–2057. <https://doi.org/10.1093/jac/dkw087>.
- Mikkelsen, H., Sivanesson, M., Filloux, A., 2011. Key two-component regulatory systems that control biofilm formation in *Pseudomonas aeruginosa*. Environ. Microbiol. 13, 1666–1681. <https://doi.org/10.1111/j.1462-2920.2011.02495.x>.
- Miyazaki, R., Yano, H., Senthilo, V., van der Meer, J.R., 2018. Physiological and transcriptome changes induced by *Pseudomonas putida* acquisition of an integrative and conjugative element. Sci. Rep. 8, 5550. <https://doi.org/10.1038/s41598-018-23858-6>.
- Miyoshi-Akiyama, T., Tada, T., Ohmagari, N., Viet Hung, N., Tharavichitkul, P., Pokhrel, B.M., Gniadkowski, M., Shimajima, M., Kirikae, T., 2017. Emergence and spread of epidemic multidrug-resistant *Pseudomonas aeruginosa*. Genome Biol. Evol. 9, 3238–3245. <https://doi.org/10.1093/gbe/evx243>.
- Moradali, M.F., Ghods, S., Rehm, B.H.A., 2017. *Pseudomonas aeruginosa* lifestyle: a paradigm for adaptation, survival, and persistence. Front. Cell. Infect. Microbiol. 7, 39. <https://doi.org/10.3389/fcimb.2017.00039>.
- Moran, N.A., 2002. Microbial minimalism: genome reduction in bacterial pathogens. Cell 108, 583–586. [https://doi.org/10.1016/s0092-8674\(02\)00665-7](https://doi.org/10.1016/s0092-8674(02)00665-7).
- Moran, N.A., McLaughlin, H.J., Sorek, R., 2009. The dynamics and time scale of ongoing genomic erosion in symbiotic bacteria. Science 323, 379–382. <https://doi.org/10.1126/science.1167140>.
- Moura, A., Soares, M., Pereira, C., Leitao, N., Henriques, I., Correia, A., 2009. INTEGRALL: a database and search engine for integrons, integrases and gene cassettes. Bioinformatics 25, 1096–1098. <https://doi.org/10.1093/bioinformatics/btp105>.
- Moya, B., Dötsch, A., Juan, C., Blázquez, J., Zamorano, L., Haussler, S., Oliver, A., 2009. β -Lactam resistance response triggered by inactivation of a nonessential penicillin-binding protein. PLoS Pathog. 5, e1000353. <https://doi.org/10.1371/journal.ppat.1000353>.
- Mulet, X., Cabot, G., Ocampo-Sosa, A.A., Domínguez, M.A., Zamorano, L., Juan, C., Tubau, F., Rodríguez, C., Moyà, B., Peña, C., Martínez-Martínez, L., Oliver, A., Spanish Network for Research in Infectious Diseases (REIPI), 2013. Biological markers of *Pseudomonas aeruginosa* epidemic high-risk clones. Antimicrob. Agents Chemother. 57, 5527–5535. <https://doi.org/10.1128/AAC.01481-13>.
- Munck, C., Albertsen, M., Telke, A., Ellabaan, M., Nielsen, P.H., Sommer, M.O.A., 2015. Limited dissemination of the wastewater treatment plant core resistome. Nat. Commun. 6, 8452. <https://doi.org/10.1038/ncomms9452>.
- Naas, T., Bonnin, R.A., Cuzon, G., Villegas, M.-V., Nordmann, P., 2013. Complete sequence of two KPC-harboring plasmids from *Pseudomonas aeruginosa*. J. Antimicrob. Chemother. 68, 1757–1762. <https://doi.org/10.1093/jac/dkt094>.
- Nadal Jimenez, P., Koch, G., Thompson, J.A., Xavier, K.B., Cool, R.H., Quax, W.J., 2012. The multiple signaling systems regulating virulence in *Pseudomonas aeruginosa*. Microbiol. Mol. Biol. Rev. 76, 46–65. <https://doi.org/10.1128/MMBR.05007-11>.
- Nascimento, A.P.B., Ortiz, M.F., Martins, W.M.B.S., Moraes, G.L., Fehlberg, L.C.C., Almeida, L.G.P., Ciapina, L.P., Gales, A.C., Vasconcelos, A.T.R., 2016. Intracolon genome stability of the metallo- β -lactamase SPM-1-producing *Pseudomonas aeruginosa* ST277, an endemic clone disseminated in Brazilian hospitals. Front. Microbiol. 7, 1946. <https://doi.org/10.3389/fmicb.2016.01946>.
- Nguyen, L., Garcia, J., Gruenberg, K., MacDougall, C., 2018. Multidrug-resistant *Pseudomonas* infections: hard to treat, but hope on the horizon? Curr. Infect. Dis. Rep. 20, 23. <https://doi.org/10.1007/s11908-018-0629-6>.
- Nicolas, E., Lambin, M., Dandoy, D., Galloy, C., Nguyen, N., Oger, C.A., Hallet, B., 2015. The Tn3-family of replicative transposons. Microbiol. Spectr. 3, MDNA3-0060-2014. <https://doi.org/10.1128/microbiolspec.MDNA3-0060-2014>.
- Núñez-Núñez, M., Navarro, M.D., Palomo, V., Rajendran, N.B., del Toro, M.D., Voss, A., Sharland, M., Sifakis, F., Tacconelli, E., Rodríguez-Baño, J., Burkert, F., Carrara, E., von Cube, M., Drgona, L., Gilchrist, K., Goossens, H., Harbarth, S., Hequet, D., Jafri, H., Kahlmeter, G., Kuster, S., Luxemburger, C., McCarthy, M., Niks, M., Oualim, A., Poljak, M., Sandulescu, O., Schweiger, A., Vuong, C., Wiegand, I., Widmer, A., Witschi, A.T., Zanetti, G., Zingg, W., 2018. The methodology of surveillance for antimicrobial resistance and healthcare-associated infections in Europe (SUSPIRE): a systematic review of publicly available information. Clin. Microbiol. Infect. 24, 105–109. <https://doi.org/10.1016/j.cmi.2017.07.014>.
- O'Neill, J., 2016. Tackling drug-resistant infections globally: final report and recommendations. Rev. Antimicrob. Resist. https://amr-review.org/sites/default/files/160525_Final%20paper_with%20cover.pdf.
- Olivares Pacheco, J., Alvarez-Ortega, C., Alcalde Rico, M., Martínez, J.L., 2017. Metabolic compensation of fitness costs is a general outcome for antibiotic-resistant *Pseudomonas aeruginosa* mutants overexpressing efflux pumps. MBio 8. <https://doi.org/10.1128/mBio.00500-17>. e00500-17.
- Oliveira, P.H., Touchon, M., Cury, J., Rocha, E.P.C., 2017. The chromosomal organization of horizontal gene transfer in bacteria. Nat. Commun. 8, 841. <https://doi.org/10.1038/s41467-017-00808-w>.
- Oliver, A., Mulet, X., López-Causapé, C., Juan, C., 2015. The increasing threat of *Pseudomonas aeruginosa* high-risk clones. Drug Resist. Updat. 21–22, 41–59. <https://doi.org/10.1016/j.DRUP.2015.08.002>.
- Olsen, R.H., Wright, C.D., 1976. Interaction of *Pseudomonas* and *Enterobacteriaceae* plasmids in *Aeromonas salmonicida*. J. Bacteriol. 128, 228–234. <https://j.b.asm.org/content/128/1/228.long>.
- Orlek, A., Stoesser, N., Anjum, M.F., Doumith, M., Ellington, M.J., Peto, T., Crook, D., Woodford, N., Walker, A.S., Phan, H., Sheppard, A.E., 2017. Plasmid classification in an era of whole-genome sequencing: application in studies of antibiotic resistance epidemiology. Front. Microbiol. 8, 182. <https://doi.org/10.3389/fmicb.2017.00182>.
- Pang, Z., Raudonis, R., Glick, B.R., Lin, T.-J., Cheng, Z., 2019. Antibiotic resistance in *Pseudomonas aeruginosa*: mechanisms and alternative therapeutic strategies. Biotechnol. Adv. 37, 177–192. <https://doi.org/10.1016/j.BIOTECHADV.2018.11.013>.
- Papagiannitsis, C.C., Medvecky, M., Chudejova, K., Skalova, A., Rotova, V., Spanelova, P., Jakubu, V., Zemlickova, H., Hrabak, J., Czech Participants of the European Antimicrobial Resistance Surveillance Network, 2017. Molecular characterization of carbapenemase-producing *Pseudomonas aeruginosa* of Czech origin and evidence for clonal spread of extensively resistant sequence Type 357 expressing IMP-7 metallo- β -lactamase. Antimicrob. Agents Chemother. 61, e01811-17. <https://doi.org/10.1128/AAC.01811-17>.
- Parkins, M.D., Somayaji, R., Waters, V.J., 2018. Epidemiology, biology, and impact of clonal *Pseudomonas aeruginosa* infections in cystic fibrosis. Clin. Microbiol. Rev. 31, e00019-18. <https://doi.org/10.1128/CMR.00019-18>.
- Pärnänen, K.M.M., Narciso-da-Rocha, C., Kneis, D., Berendonk, T.U., Cacace, D., Do, T.T., Elpers, C., Fatta-Kassinos, D., Henriques, I., Jaeger, T., Karkman, A., Martínez, J.L., Michael, S.G., Michael-Kordatou, I., O'Sullivan, K., Rodríguez-Mozaz, S., Schwartz, T., Sheng, H., Sorum, H., Stedtfeld, R.D., Tiedje, J.M., Giustina, S.V., Della, Walsh, F., Vaz-Moreira, I., Virta, M., Manaia, C.M., 2019. Antibiotic resistance in European wastewater treatment plants mirrors the pattern of clinical antibiotic resistance prevalence. Sci. Adv. 5, eaau9124. <https://doi.org/10.1126/sciadv.aau9124>.
- Partridge, S.R., 2011. Analysis of antibiotic resistance regions in Gram-negative bacteria. FEMS Microbiol. Rev. 35, 820–855. <https://doi.org/10.1111/j.1574-6976.2011.00277.x>.
- Partridge, S.R., Brown, H.J., Stokes, H.W., Hall, R.M., 2001. Transposons Tn1696 and Tn21 and their integrons In4 and In2 have independent origins. Antimicrob. Agents Chemother. 45, 1263–1270. <https://doi.org/10.1128/AAC.45.4.1263-1270.2001>.
- Partridge, S.R., Di Pilato, V., Doi, Y., Feldgarden, M., Haft, D.H., Klimke, W., Kumar-Singh, S., Liu, J.-H., Malhotra-Kumar, S., Prasad, A., Rossolini, G.M., Schwarz, S., Shen, J., Walsh, T., Wang, Y., Xavier, B.B., 2018a. Proposal for assignment of allele numbers for mobile colistin resistance (mcr) genes. J. Antimicrob. Chemother. 73, 2625–2630. <https://doi.org/10.1093/jac/dky262>.
- Partridge, S.R., Kwong, S.M., Firth, N., Jensen, S.O., 2018b. Mobile genetic elements associated with antimicrobial resistance. Clin. Microbiol. Rev. 31, e00088-17. <https://doi.org/10.1128/CMR.00088-17>.
- Partridge, S.R., Tsafnat, G., 2018. Automated annotation of mobile antibiotic resistance in Gram-negative bacteria: the Multiple Antibiotic Resistance Annotator (MARA) and database. J. Antimicrob. Chemother. 73, 883–890. <https://doi.org/10.1093/jac/dkx513>.
- Peña-Miller, R., Rodríguez-González, R., MacLean, R.C., San Millán, A., 2015. Evaluating the effect of horizontal transmission on the stability of plasmids under different selection regimes. Mob. Genet. Elem. 5, 1–5. <https://doi.org/10.1080/2159256X.2015.1045115>.
- Petrovski, S., Blackmore, D.W., Jackson, K.L., Stanisich, V.A., 2011. Mercury(II)-resistance transposons Tn502 and Tn512, from *Pseudomonas* clinical strains, are structurally different members of the Tn5053 family. Plasmid 65, 58–64. <https://doi.org/10.1016/j.plasmid.2010.08.003>.
- Pfennigwerth, N., Lange, F., Belmar Campos, C., Hentschke, M., Gattermann, S.G., Kaase, M., 2017. Genetic and biochemical characterization of HMB-1, a novel subclass B1 metallo- β -lactamase found in a *Pseudomonas aeruginosa* clinical isolate. J. Antimicrob. Chemother. 72, dkx554. <https://doi.org/10.1093/jac/dkx554>.
- Picao, R.C., Poirer, L., Gales, A.C., Nordmann, P., 2009. Diversity of β -lactamases produced by ceftazidime-resistant *Pseudomonas aeruginosa* isolates causing bloodstream infections in Brazil. Antimicrob. Agents Chemother. 53, 3908–3913. <https://doi.org/10.1128/AAC.00453-09>.
- Pirnay, J.-P., Bilocq, F., Pot, B., Cornelis, P., Zizi, M., Van Eldere, J., Deschaght, P., Vanechoutte, M., Jennes, S., Pitt, T., De Vos, D., 2009. *Pseudomonas aeruginosa* population structure revisited. PLoS One 4, e7740. <https://doi.org/10.1371/journal.pone.0007740>.
- Poirer, L., Rodríguez-Martínez, J.-M., Al Naiemi, N., Debets-Ossenkopp, Y.J., Nordmann, P., 2010. Characterization of DIM-1, an integron-encoded metallo-beta-lactamase from a *Pseudomonas stutzeri* clinical isolate in the Netherlands. Antimicrob. Agents Chemother. 54, 2420–2424. <https://doi.org/10.1128/AAC.01456-09>.
- Political Declaration of the High-Level Meeting of the General Assembly on Antimicrobial Resistance: draft resolution/submitted by the President of the General Assembly. <https://digitallibrary.un.org/record/842813?ln=en>.
- Polz, M.F., Alm, E.J., Hanage, W.P., 2013. Horizontal gene transfer and the evolution of bacterial and archaeal population structure. Trends Genet. 29, 170–175. <https://doi.org/10.1016/j.tig.2012.12.006>.
- Popa, O., Dagan, T., 2011. Trends and barriers to lateral gene transfer in prokaryotes. Curr. Opin. Microbiol. 14, 615–623. <https://doi.org/10.1016/j.mib.2011.07.027>.
- Popa, O., Hazkani-Covo, E., Landan, G., Martin, W., Dagan, T., 2011. Directed networks reveal genomic barriers and DNA repair bypasses to lateral gene transfer among prokaryotes. Genome Res. 21, 599–609. <https://doi.org/10.1101/gr.115592.110>.
- Quainoo, S., Coolen, J.P.M., van Hijum, S.A.F.T., Huynen, M.A., Melchers, W.J.G., van Schaik, W., Wertheim, H.F.L., 2017. Whole-genome sequencing of bacterial pathogens: the future of nosocomial outbreak analysis. Clin. Microbiol. Rev. 30, 1015–1063. <https://doi.org/10.1128/CMR.00016-17>.
- Queenan, A.M., Bush, K., 2007. Carbapenemases: the versatile beta-lactamases. Clin. Microbiol. Rev. 20, 440–458. <https://doi.org/10.1128/CMR.00001-07>.
- Quick, J., Cumley, N., Wearn, C.M., Niebel, M., Constantinidou, C., Thomas, C.M., Pallen, J.

- M.J., Moiemien, N.S., Bamford, A., Oppenheim, B., Loman, N.J., 2014. Seeking the source of *Pseudomonas aeruginosa* infections in a recently opened hospital: an observational study using whole-genome sequencing. *BMJ Open* 4, e006278. <https://doi.org/10.1136/bmjopen-2014-006278>.
- Ragheb, M.N., Thomason, M.K., Hsu, C., Nugent, P., Gage, J., Samadpour, A.N., Kariisa, A., Merrikkh, C.N., Miller, S.I., Sherman, D.R., Merrikkh, H., 2018. Inhibiting the evolution of antibiotic resistance. *Mol. Cell* 73, 157–165.e5. <https://doi.org/10.1016/j.molcel.2018.10.015>.
- Ramsay, J.P., Firth, N., 2017. Diverse mobilization strategies facilitate transfer of non-conjugative mobile genetic elements. *Curr. Opin. Microbiol.* 38, 1–9. <https://doi.org/10.1016/j.mib.2017.03.003>.
- Rankin, D.J., Rocha, E.P.C., Brown, S.P., 2011. What traits are carried on mobile genetic elements and why? *Heredity* (Edinb) 106, 1–10. <https://doi.org/10.1038/hdy.2010.24>.
- Ravatt, R., Studer, S., Springael, D., Zehnder, A.J., van der Meer, J.R., 1998. Chromosomal integration, tandem amplification, and deamplification in *Pseudomonas putida* F1 of a 105-kilobase genetic element containing the chlorocatechol-degradative genes from *Pseudomonas* sp. Strain B13. *J. Bacteriol.* 180, 4360–4369.
- Roberts, A.P., Chandler, M., Courvalin, P., Guédon, G., Mullany, P., Pembroke, T., Rood, J.I., Smith, C.J., Summers, A.O., Tsuda, M., Berg, D.E., 2008. Revised nomenclature for transposable genetic elements. *Plasmid* 60, 167–173. <https://doi.org/10.1016/j.plasmid.2008.08.001>.
- Roberts, A.P., Mullany, P., 2011. Tn916-like genetic elements: a diverse group of modular mobile elements conferring antibiotic resistance. *FEMS Microbiol. Rev.* 35, 856–871. <https://doi.org/10.1111/j.1574-6976.2011.00283.x>.
- Rodríguez-Beltrán, J., Hernández-Beltrán, J.C.R., DelaFuente, J., Escudero, J.A., Fuentes-Hernández, A., MacLean, R.C., Peña-Miller, R., San Millán, A., 2018. Multicopy plasmids allow bacteria to escape from fitness trade-offs during evolutionary innovation. *Nat. Ecol. Evol.* 2, 873–881. <https://doi.org/10.1038/s41559-018-0529-z>.
- Rodríguez-Martínez, J.-M., Poirel, L., Nordmann, P., 2009. Extended-spectrum cephalosporins in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 53, 1766–1771. <https://doi.org/10.1128/AAC.01410-08>.
- Roemhild, R., Gokhale, C.S., Dirksen, P., Blake, C., Rosenstiel, P., Traulsen, A., Andersson, D.I., Schulenburg, H., 2018. Cellular hysteresis as a principle to maximize the efficacy of antibiotic therapy. *Proc. Natl. Acad. Sci. U. S. A.* 115, 9767–9772. <https://doi.org/10.1073/pnas.1810004115>.
- Roosaare, M., Puustusmaa, M., Möls, M., Vaher, M., Remm, M., 2018. PlasmidSeeker: identification of known plasmids from bacterial whole genome sequencing reads. *PeerJ* 6, e4588. <https://doi.org/10.7717/peerj.4588>.
- Rosenthal, V.D., Al-Abdely, H.M., El-Kholi, A.A., AlKhawaja, S.A.A., Leblebicioglu, H., Mehta, Y., Rai, V., Hung, N.V., Kanji, S.S., Salama, M.F., Salgado-Yeppez, E., Elahi, N., Morfin Otero, R., Apisarnthanarak, A., De Carvalho, B.M., Ider, B.E., Fisher, D., Buenafior, M.C.S.G., Petrov, M.M., Quesada-Mora, A.M., Zand, F., Gurskis, V., Anguseva, T., Ikram, A., Aguilar de Moros, D., Duszyńska, W., Mejia, N., Horhat, F.G., Belskiy, V., Mijoljevic, V., Di Silvestre, G., Furova, K., Ramos-Ortiz, Gloria, Y., Gamar Elanbya, M.O., Satari, H.I., Gupta, U., Dendane, T., Raka, L., Guanche-Garcell, H., Hu, B., Padgett, D., Jayatilake, K., Ben Jaballah, N., Apostolopoulou, E., Prudencio Leon, W.E., Sepulveda-Chavez, A., Telechea, H.M., Trotter, A., Alvarez-Moreno, C., Kushner-Davalos, L., Desse, J.E., Maurizi, D.M., Montanini, A.M., Chaparro, G.J., Stagnaro, J.P., Romani, A., Bianchi, A.C., Álvarez, G., Palaoro, A., Bernan, M., Cabrera-Montesino, R., Domínguez, C., Rodríguez, C.G., Silva, C.G., Bogdanowicz, E., Riera, F.O., Benchetrit, G., Perez, I., Vimercati, J., Marcos, L.S., Ramasco, L., Caridi, M., Oyola, M.C., Rodríguez, M.C., Spadaro, M.L., Olivieri, M.S., Saul, P., Juarez, P.D., Pérez, R.H., Botta, P., Quintana, D.Q., Ríos, A.M., Stagnaro, J.P., Chediack, V., Chilon, W., Alsayegh, A.I., Yaseen, F.H., Hani, L.F., Sower, S.F., Magray, T.A., Medeiros, E.A., Alves De Oliveira, A., Romario-Mendes, A., Fernandes-Valente, C., Santos, C., Escudeiro, D., Azevedo-Ferreira Lima, D., Azevedo-Pereira, D., Onzi-Siliprandi, E.M., Serpa-Maia, F., Aguiar-Leitao, F., Assuncao-Ponte, G., Dos Anjos-Lima, J., Olszewski, J., Harten Pinto Coelho, K., Alves De Lima, L.F., Mendonça, M., Maciel-Canuto Amaral, M.L., Tenorio, M.T., Gerah, S., Andrade-Oliveira-Reis, M., Moreira, M., Ximenes-Rocha Batista, M., Campos-Uchoa, R.S., Rocha-Vasconcelos Carneiro, R., Amaral De Moraes, R., Do Nascimento, S.C., Moreira-Matos, T., Lima-De Barros Araujo, T.M., De Jesus Pinheiro-Bandeira, T., Machado-Silva, V.L., Santos Monteiro, W.M., Hristozova, E., Kostadinov, E.D., Angelova, K., Velinova, V.A., Dicheva, Y., Guo, X., Ye, G., Li, R., Song, L., Liu, K., Liu, T., Song, G., Wang, C., Yang, X., Yu, H., Yang, Y., Martínez, A., Vargas-García, A.R., Lagares-Guzmán, A., González, A.P., Linares, C., Ávila-Acosta, C., Santofimio, D., Yepes-Gomez, D., Marin-Tobar, D.A., Mazo-Elorza, D.P., Chapeta-Parada, E.G., Camacho-Moreno, G., Roncancio-Vill, G.E., Valderrama-Marquez, I.A., Ruiz-Gallardo, J.E., Ospina-Martínez, J.O., Osorio, J., Marín-Urbe, J.I., López, J.C., Gualtero, S., Rojas, J.R., Gomez-Nieto, K., Rincon, L.Y.M., Meneses-Ovallos, L., Canas-Giraldo, L.M., Burgos-Florez, L.D., Amaral-Almeida Costa, M., Rodriguez, M., Barahona-Guzmán, N., Mancera-Paez, O., Rios-Arana, P.A., Ortega, R., Romero-Torres, S.L., Pulido-Leon, S.M., Valderrama, S., Moreno-Mejia, V.M., Raigoza-Martinez, W., Villamil-Gomez, W., Pardo-Lopez, Y.A., Argüello-Ruiz, A., Solano-Chinchilla, A., Muñoz-Gutierrez, G.A., Calvo-Hernández, I., Maroto-Vargas, L., Zuniga, M.A., Valverde-Hernandez, M., Chavarria-Ugalde, O., Herrera, B., Díaz, C., Bovera, M.M., Cevallos, C., Pelaez, C., Jara, E., Delgado, V., Coello-Gordon, E.E., Picoita, F., Guerrero-Toapant, F.M., Valencia, F., Santacruz, G., Gonzalez, H., Pazzino, L.N., Garcia, M.F., Arboleda, M., Lascano, N., Alquina, N., Ramírez, V., Yousef, R.H.A., Moustafa, A.E.M., Ahmed, A., Elansary, A.M., Ali, A.M., Hasanin, A., Messih, A.A., Ramadan, A., El Awady, B.A., Hassan, D.M., Abd El Aziz, D., Hamza, H., Agha, H.M., Ghazi, I.A., Elkhalil, J., Fattah, M.A., Elanany, M., Mansour, M., Haleim, M.M.A., Fouda, R., El-Sherif, R.H., Bekeit, S., Bayani, V., Elkholi, Y.S., Abdelhamid, Y.M., Salah, Z., Rivera, D.M., Chawla, A., Manked, A.N., Azim, A., Mubarak, A., Thakur, A., Dharan, A.V., Patil, A., Sasidharan, A., Bilolikar, A.K., Anirban Karmakar, A., Mathew, A.M., Kulkarni, A., Agarwal, A.,
- Sriram, A., Dwivedy, A., Dasgupta, A., Bhakta, A., Suganya, A.R., Pojary, A., Mani, A.K., Sakle, A., Abraham, B.K., Padmini, B., Ramachandran, B., Ray, B., Pati, B.K., Chaudhury, B.N., Mishra, B.M., Biswas, S., Saibala, M.B., Jawadwala, B.Q., Rodrigues, C., Modi, C., Patel, C., Khanna, D.K., Devaprasad, D., Divekar, D., Aggarwal, D.G., Divatia, J.V., Zala, D., Pathrose, E., Abubakar, F., Chacko, F., Gehlot, G.S., Khanna, G., Sale, H.K., Roy, I., Shelaonkar, J., Sorabjee, J., Eappen, J., Mathew, J., Pal, J., Varma, K., Joshi, K.L., Sandhu, K., Kelkar, R., Ranganathan, L., Pushparaj, L., Lavate, M., Latha, M., Suryawanshi, M., Bhattacharyya, M., Kavathekar, M., Agarwal, M.K., Patel, M., Shah, M., Sivakumar, M.N., Kharbanda, M., Bej, M., Potdar, M., Chakravarthy, M., Karpagam, M., Myatra, S.N., Gita, N., Rao, N.P., Sen, N., Ramakrishnan, N., Jaggi, N., Anani, N., Pawar, N.K., Modi, N., Pandya, N., Mohanty, N., Thakkar, P., Joshi, P., Sahoo, P.K., Nair, P.K., Kumar, P.S., Patil, P., Mukherjee, P., Mathur, P., Shah, P., Sukanya, R., Arjun, R., Chawla, R., Gopalakrishnan, R., Venkataraman, R., Raut, S., Krupanandan, R., Tejani, R., Misra, R., Debroy, R., Saranya, S., Narayanan, S., Mishra, S., Saseedharan, S., Sengupta, S., Patnaik, S.K., Sinha, S., Blessymole, S., Rohra, S., Rajagopal, S., Mukherjee, S., Sengupta, S., John, S., Bhattacharya, S., Singh, S., Sohanlal, T., Vadi, S., Dalal, S.S., Todi, S.K., Kumar, S., Kansal, S., Misra, S., Bhattacharyya, S., Nirkhawale, S., Purkayastha, S.K., Mukherjee, S., Singh, S., Sahu, S., Sharma, S., Kumar, S., Basu, S., Shetty, S., Shah, S., Singhal, T., Francis, T., Anand, T., Venkateshwar, V., Thomas, V., Kothari, V., Velupandi, Kantroo, V., Sitohang, G., Kadarsh, R., Sanaei, A., Maghsudi, B., Sabetian, G., Masjedi, M., Alebouyeh, M., Sherafat, S.J., Mohamed, Y.K., Al Khamis, A., Alsaadi, A.S., Al-Jarir, A.A., Mutwalli, A.H., Rillorta, A.S.K., Thomas, A., Kelany, A., Manao, A., Alamri, D.M., Santiago, E.B., Cruzpero, E.P., Sawan, F.A., Al Qasmah, F.A., Alabdaly, H., Al-Dossary, H.A., Ahmed, H., Roshdi, H., Al-Akhami, H.Y., Hanafi, H., Ammari, H., Hani, H.M.A., Asiri, I.A.M.A., Mendoza, J.A., Philipose, J., Selga, J.O., Kehkashan, Ghalilah, K.M., Redito, L.S., Joshi, L., Al-Alawi, M., Al-Gethamy, M.M., Madco, M., Manuel, M.G., Girvan, M., Aldalaton, M., De Guzman, M., Alkhamaly, M., Masfar, M., Karrar, M.A.A., Al Azmi, M.M., Qusai, M.L., Torres, M.M., Al-Abdullah, N., Tawfic, N.A., Elsayed, N., Abdulkhalik, N.S., Bugis, N.A., Ariola, N.C., Gad, N., Alghosn, N., Tashkandi, N., Zharani, N.A., De Vera, P.A., Krishnan, R., Al Shahr, R.H., Jaha, R.N.A., Thomas, R., Cresencia, R.L., Penuliri, R., Lozada, R.V., Al Qahtani, S., Twfik, S., Al Faraj, S.H., El-Sherbiny, S., Ali, S.J.B., Briones, S., Bukhari, S.Z., Alotaibi, T.S.A., Gopal, U., Nair, U., Abdulatif, W.A., Hussain, W.M., Demotica, W.M., Spahija, G., Bafti, N., Gashi, A., Omar, A.A., Mohamed, A.M., Rebello, F.M., Almousa, H.H., Abdo, N.M., George, S.M., Khamis, S., Thomas, S., Ahmad Zaatar, A., Anwar Al Souheil, A., Ayash, H., Zeid, I., Tannous, J., Zahreddine, N.K., Ahmadi, R., Mahfouz, T., Kardas, T., Tanzi, V., Kanafani, Z., Hammoud, Z., Dagys, A., Grinkevičiute, D., Kevalas, R., Kondratas, T., Petrovska, M., Popovska, K., Mitrev, Z., Miteva, Z.B., Jankovska, K., Gurovska, S.T., Gan, C.S., Othman, A.A., Yusof, A.M., Abidin, A.S.Z., Aziz, F.A., Weng, F.K., Zainol, H., Bakar, K.B.A., Lum, L.C.S., Mansor, M., Zaman, M.K., Jamaluddin, M.F.H., Hasan, M.S., Rahman, R.A., Zaini, R.H.M., Zhazali, R., Sri Ponnampala, S.S.L., Chuah, S.L., Shukeri, W.F.W.M., Hassan, W.N.W., Yusoff, W.N.W., Mat, W.R.W., Curenio-Diaz, M.A., Aguirre-Avalos, G., Flores-Alvarado, A., Cerero-Gudino, A., Zamores-Pedroza, A., Cano-Munoz, B., Hernandez-Chena, B.E., Carreon-Martinez, C.C., Coronado-Magana, H., Corona-Jimenez, F., Rodriguez-Noriega, E., Alcala-Martinez, E., Gonzalez-Diaz, E., Guerra-Infante, F.M., Arteaga-Troncoso, G., Martinez-Falcon, G., Leon-Garnica, G., Delgado-Aguirre, H.A., Perez-Gomez, H.R., Sosa-Gonzalez, I.E., Galindo-Ortega, J.A., Ayala-Gaytan, J.J., Rodriguez-Pacheco, J., Zamorano-Flores, L., Lopez-Pulgarin, J.A., Miranda-Navales, M.G., Ramirez, M., Lopez-Hurtado, M., Lozano, M., Gomez, M.E., Sanchez-Castuera, M.E., Kasten-Monges, M., Gonzalez-Martinez, M., Sanchez-Vargas, M., Culebro-Burguet, M.C., Altuzar-Figueroa, M.A., Mijangos-Mendez, J.C., Ramirez, O.G., Espinosa, O.S., De Leon-Escobedo, R., Salas-Flores, R., Ruiz-Rendon, R., Petersen-Morfin, S., Aguirre-Diaz, S.A., Esparza-Ahumada, S., Vega-Gonzalez, S., Gaona-Flores, V., Monroy-Colin, V.A., Cruz-Rivera, Z., Bat-Erdene, A., Narankhuu, B., Chojiamts, B., Tuvdenyam, B., Batkhuu, B., Chuluunchimeg, K.H., Enkhsetseg, D., Batjargal, G., Bayasgalan, G., Dorj, M., Mendsaikhan, N., Baatar, O., Suverderene, P., Baigalmaa, S., Khajidmaa, T., Begzjav, T., Tsuyanga, Ariyasuren, Z., Zeggwag, A.A., Berechid, K., Abidi, K., Madani, N., Aboual, R., Koirala, A., Giri, R., Sainju, S., Acharya, S.P., Ahmed, A., Raza, A., Parveen, A., Sultan, F., Khan, M., Paul, N., Daud, N., Yusuf, S., Nizamuddin, S., Garcia-Mayorca, E., Castaño, E.G., Moreno-Castillo, J.L., Ballinas-Aquino, J.M., Lara, L., Vargas, M., Rojas-Bonilla, M.I., Ramos, S.J., Mapp, T., De Iturrazo, V., La Hoz Vergara, C.E., Linares-Calderon, C.F., Moreno, D., Ramirez, E., Ramirez Wong, F.M., Montenegro-Orrego, G.M., Sandoval-Castillo, H.R., Pichilingue-Chagray, J., Mueras-Quevedo, J., Aibar-Yaranga, K.F., Castillo-Bravo, L.I., Santivanez-Monge, L.M., Mayorga-Espichan, M.J., Rosario-Tueros, M.L.G., Changano-Rodriguez, M.V., Salazar-Ramirez, N.E., Marquez-Mondalga, V.A., Tajanlangit, A.L.N., Tamayo, A.S., Llamas, C.M.J.P., Labro, E., Dy, A.P., Fortin, J.D., Bergosa, L.D., Salvio, L.G., Bermudez, V., Sg-Buenafior, M.C., Trajano, M.F., Mendoza, M.T., Javellana, O.P., Maglente, R.R., Arreza-Galapia, Y., Navoa-Ng, J.A., Kubler, A., Barteczko-Grajek, B., Dragan, B., Zurawska, M., Mikaszewska-Sokolewicz, M., Zielinska, M., Ramos-Ortiz, G.Y., Florin-Rogobete, A., Vlad, C.D., Muntean, D., Sandesc, D., Papurica, M., Licker, M., Bedreag, O.H., Popescu, R., Grecu, S.D., Dumitrascu, V., Molkov, A., Galishevskiy, D., Furman, M., Simic, A., Lekic, D., Ristic, G., Eremija, J., Kojovic, J., Nikolic, L., Bjelovic, M., Lesnakova, A., Hlinkova, S., Gamar-Elanbya, M.O., Supa, N., Prasan, P., Pimathai, R., Wanitanukool, S., Somabutr, S., Ben-Jaballah, N., Borgi, A., Bouziri, A., Dilek, A., Oncul, A., Kaya, A., Demiroz, A.P., Gunduz, A., Ozgultekin, A., Inan, A., Yalcin, A.N., Ramazanoglu, A., Engin, A., Willke, A., Mecro, B.C., Aygun, C., Bulut, C., Uzun, C., Becerik, C., Hatipoglu, C.A., Guclu, C.Y., Ozdemir, D., Yildizdas, D., Ugurcan, D., Azak, E., Guclu, E., Yilmaz, E.M., Sebnem-Erdinc, F., Sirmatel, F., Ulger, F., Sari, F., Kizilates, F., Usluer, G., Ceylan, G., Ersoz, G., Kaya, G., Ertem, G.T., Senol, G., Agin, H., Cadakad, H., Yilmaz, H., Sungurtekin, H., Zengin, H., Turgut, H., Ozgunes, I., Devrim, I., Erdem, I., Işcanlı, I.G.E., Bakir, M.M., Geyik, M.F., Oral, M., Meric, M., Cengiz, M., Ozelcik, M., Altindis, M., Sunbul, M., Elaldi, N., Kuyucu, N.,

- Unal, N., Oztoprak, N., Yasar, N., Erben, N., Bayram, N., Dursun, O., Karabay, O., Coskun, O., Horoz, O.O., Turhan, O., Sandal, O.S., Tekin, R., Esen, S., Erdogan, S.Y., Unal, S., Karacorlu, S., Sen, S., Sen, S., Sacar, S., Yazar, V., Oruc, Y., Sahip, Y., Kaya, Z., Philip, A., Elhoufi, A., Alrahma, H., Sached, E., Perez, F., Empaire, G.D., Vidal, H., Montes-Bravo, L., Guzman Siriti, M.E., Orozco, N., Navarrete, N., Ruiz, Y., De Anez, Z.D.-G., Van Trang, D.T., Minh, D.Q., Co, D.X., Anh, D.P.P., Thu, L.T.A., Tuyet, L.T.D., Nguyen, L.T.T., Chau, N.uy, Binh, N.G., Tien, N.P., Anh, N.Q., Hang, P.T., Hanh, T.T.M., Hang, T.T.T., Thu, T.A., Thoa, V.T.H., 2016. International Nosocomial Infection Control Consortium report, data summary of 50 countries for 2010–2015: device-associated module. *Am. J. Infect. Control* 44, 1495–1504. <https://doi.org/10.1016/j.ajic.2016.08.007>.
- Rossen, J.W.A., Friedrich, A.W., Moran-Gilad, J., ESCMID Study Group for Genomic and Molecular Diagnostics (ESGMD), 2018. Practical issues in implementing whole-genome-sequencing in routine diagnostic microbiology. *Clin. Microbiol. Infect.* 24, 355–360. <https://doi.org/10.1016/j.cmi.2017.11.001>.
- Rossi, E., Falcone, M., Molin, S., Johansen, H.K., 2018. High-resolution in situ transcriptomics of *Pseudomonas aeruginosa* unveils genotype independent patho-phenotypes in cystic fibrosis lungs. *Nat. Commun.* 9, 3459. <https://doi.org/10.1038/s41467-018-05944-5>.
- Roy Chowdhury, P., Merlino, J., Labbate, M., Cheong, E.Y.-L., Gottlieb, T., Stokes, H.W., 2009. Tn6060, a transposon from a genomic island in a *Pseudomonas aeruginosa* clinical isolate that includes two class 1 integrons. *Antimicrob. Agents Chemother.* 53, 5294–5296. <https://doi.org/10.1128/AAC.00687-09>.
- Roy Chowdhury, P., Scott, M., Worden, P., Huntington, P., Hudson, B., Karagiannis, T., Charles, I.G., Djordjevic, S.P., 2016. Genomic islands 1 and 2 play key roles in the evolution of extensively drug-resistant ST235 isolates of *Pseudomonas aeruginosa*. *Open Biol.* 6. <https://doi.org/10.1098/rsob.150175>.
- Rozov, R., Brown Kav, A., Bogumil, D., Shterzer, N., Halperin, E., Mizrahi, I., Shamir, R., Valencia, A., 2016. Recycler: an algorithm for detecting plasmids from de novo assembly graphs. *Bioinformatics* 33, btw651. <https://doi.org/10.1093/bioinformatics/btw651>.
- Rozwandowicz, M., Brouwer, M.S.M., Fischer, J., Wagenaar, J.A., Gonzalez-Zorn, B., Guerra, B., Mevius, D.J., Hordijk, J., 2018. Plasmids carrying antimicrobial resistance genes in *Enterobacteriaceae*. *J. Antimicrob. Chemother.* 73, 1121–1137. <https://doi.org/10.1093/jac/dkx488>.
- Rubio-Cosials, A., Schulz, E.C., Lamberts, L., Smyshlyayev, G., Rojas-Cordova, C., Forslund, K., Karaca, E., Bebel, A., Bork, P., Barabas, O., 2018. Transposase-DNA complex structures reveal mechanisms for conjugative transposition of antibiotic resistance. *Cell* 173, 208–220.e20. <https://doi.org/10.1016/j.cell.2018.02.032>.
- Ryan, M.P., Pembroke, J.T., Adley, C.C., 2009. Novel Tn4371-ICE like element in *Ralstonia pickettii* and Genome mining for comparative elements. *BMC Microbiol.* 9, 242. <https://doi.org/10.1186/1471-2180-9-242>.
- Salabi, A.E., Toleman, M.A., Weeks, J., Bruderer, T., Frei, R., Walsh, T.R., 2010. First report of the metallo-beta-lactamase SPM-1 in Europe. *Antimicrob. Agents Chemother.* 54, 582. <https://doi.org/10.1128/AAC.00719-09>.
- Samuelsen, O., Buarø, L., Toleman, M.A., Giske, C.G., Hermansen, N.O., Walsh, T.R., Sundsfjord, A., 2009. The first metallo-beta-lactamase identified in Norway is associated with a TnIC-like transposon in a *Pseudomonas aeruginosa* isolate of sequence type 233 imported from Ghana. *Antimicrob. Agents Chemother.* 53, 331–332. <https://doi.org/10.1128/AAC.00785-08>.
- San Millan, A., 2018. Evolution of plasmid-mediated antibiotic resistance in the clinical context. *Trends Microbiol.* 26, 978–985. <https://doi.org/10.1016/j.tim.2018.06.007>.
- San Millan, A., Escudero, J.A., Gifford, D.R., Mazel, D., MacLean, R.C., 2016. Multicopy plasmids potentiate the evolution of antibiotic resistance in bacteria. *Nat. Ecol. Evol.* 1, 0010. <https://doi.org/10.1038/s41559-016-0010>.
- San Millan, A., Heilbron, K., MacLean, R.C., 2014a. Positive epistasis between co-infecting plasmids promotes plasmid survival in bacterial populations. *ISME J.* 8, 601–612. <https://doi.org/10.1038/ismej.2013.182>.
- San Millan, A., Peña-Miller, R., Toll-Riera, M., Halbert, Z.V., McLean, A.R., Cooper, B.S., MacLean, R.C., 2014b. Positive selection and compensatory adaptation interact to stabilize non-transmissible plasmids. *Nat. Commun.* 5, 5208. <https://doi.org/10.1038/ncomms6208>.
- San Millan, A., MacLean, R.C., 2017. Fitness costs of plasmids: a limit to plasmid transmission. *Microbiol. Spectr.* 5. <https://doi.org/10.1128/microbiolspec.MTBP-0016-2017>.
- San Millan, A., Toll-Riera, M., Escudero, J.A., Cantón, R., Coque, T.M., MacLean, R.C., 2015a. Sequencing of plasmids pAMBL1 and pAMBL2 from *Pseudomonas aeruginosa* reveals a *bla*_{VIM-1} amplification causing high-level carbapenem resistance. *J. Antimicrob. Chemother.* 70, 3000–3003. <https://doi.org/10.1093/jac/dkv222>.
- San Millan, A., Toll-Riera, M., Qi, Q., Betts, A., Hopkinson, R.J., McCullagh, J., MacLean, R.C., 2018. Integrative analysis of fitness and metabolic effects of plasmids in *Pseudomonas aeruginosa* PAO1. *ISME J.* 12, 3014–3024. <https://doi.org/10.1038/s41396-018-0224-8>.
- San Millan, A., Toll-Riera, M., Qi, Q., MacLean, R.C., 2015b. Interactions between horizontally acquired genes create a fitness cost in *Pseudomonas aeruginosa*. *Nat. Commun.* 6, 6845. <https://doi.org/10.1038/ncomms7845>.
- Sánchez-Diener, I., Zamorano, L., López-Causapé, C., Cabot, G., Mulet, X., Peña, C., del Campo, R., Cantón, R., Doménech-Sánchez, A., Martínez-Martínez, L., Arcos, S.C., Navas, A., Oliver, A., 2017. Interplay among resistance profiles, high-risk clones, and virulence in the *Caenorhabditis elegans* *Pseudomonas aeruginosa* infection model. *Antimicrob. Agents Chemother.* 61, e01586-17. <https://doi.org/10.1128/AAC.01586-17>.
- Sander, P., Springer, B., Prammananan, T., Sturmfels, A., Kappler, M., Pletschette, M., Böttger, E.C., 2002. Fitness cost of chromosomal drug resistance-conferring mutations. *Antimicrob. Agents Chemother.* 46, 1204–1211. <https://doi.org/10.1128/aac.46.5.1204-1211.2002>.
- Sawa, T., Hamaoka, S., Kinoshita, M., Kainuma, A., Naito, Y., Akiyama, K., Kato, H., 2016. *Pseudomonas aeruginosa* Type III secretory toxin ExoU and its predicted homologs. *Toxins (Basel)* 8, 307. <https://doi.org/10.3390/toxins8110307>.
- Schenk, M.F., Szendro, I.G., Salverda, M.L.M., Krug, J., de Visser, J.A.G.M., 2013. Patterns of epistasis between beneficial mutations in an antibiotic resistance gene. *Mol. Biol. Evol.* 30, 1779–1787. <https://doi.org/10.1093/molbev/mst096>.
- Schick, A., Kassen, R., 2018. Rapid diversification of *Pseudomonas aeruginosa* in cystic fibrosis lung-like conditions. *Proc. Natl. Acad. Sci. U. S. A.* 115, 10714–10719. <https://doi.org/10.1073/pnas.1721270115>.
- Scotta, C., Juan, C., Cabot, G., Oliver, A., Lalucat, J., Bennisar, A., Alberti, S., 2011. Environmental microbiota represents a natural reservoir for dissemination of clinically relevant metallo-beta-lactamases. *Antimicrob. Agents Chemother.* 55, 5376–5379. <https://doi.org/10.1128/AAC.00716-11>.
- Secor, P.R., Michaels, L.A., Ratjen, A., Jennings, L.K., Singh, P.K., 2018. Entropically driven aggregation of bacteria by host polymers promotes antibiotic tolerance in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. U. S. A.* 115, 10780–10785. <https://doi.org/10.1073/pnas.1806005115>.
- Sevillano, E., Gallego, L., García-Lobo, J.M., 2009. First detection of the OXA-40 carbapenemase in *P. aeruginosa* isolates, located on a plasmid also found in *A. baumannii*. *Pathol. Biol.* 57, 493–495. <https://doi.org/10.1016/j.patbio.2008.05.002>.
- Sheppard, A.E., Stoesser, N., German-Mesner, I., Vegesana, K., Sarah Walker, A., Crook, D.W., Mathers, A.J., 2018. TETyper: a bioinformatic pipeline for classifying variation and genetic contexts of transposable elements from short-read whole-genome sequencing data. *Microb. Genomics* 4. <https://doi.org/10.1099/mgen.0.000232>.
- Shimizu, W., Kayama, S., Kouda, S., Ogura, Y., Kobayashi, K., Shigemoto, N., Shimada, N., Yano, R., Hisatsune, J., Kato, F., Hayashi, T., Sueda, T., Ohge, H., Sugai, M., 2015. Persistence and epidemic propagation of a *Pseudomonas aeruginosa* sequence type 235 clone harboring an IS26 composite transposon carrying the blaIMP-1 integron in Hiroshima, Japan, 2005 to 2012. *Antimicrob. Agents Chemother.* 59, 2678–2687. <https://doi.org/10.1128/AAC.04207-14>.
- Shortridge, D., Gales, A.C., Streit, J.M., Huband, M.D., Tsakris, A., Jones, R.N., 2019. Geographic and temporal patterns of antimicrobial resistance in *Pseudomonas aeruginosa* over 20 years from the SENTRY Antimicrobial Surveillance Program, 1997–2016. *Open Forum Infect. Dis.* 6, S63–S68. <https://doi.org/10.1093/ofid/ofy343>.
- Silby, M.W., Winstanley, C., Godfrey, S.A.C., Levy, S.B., Jackson, R.W., 2011. *Pseudomonas* genomes: diverse and adaptable. *FEMS Microbiol. Rev.* 35, 652–680. <https://doi.org/10.1111/j.1574-6976.2011.00269.x>.
- Silveira, M.C., Albano, R.M., Asensi, M.D., Carvalho-Assef, A.P.D., 2016. Description of clone ST277. *Infect. Genet. Evol.* 42, 60–65. <https://doi.org/10.1016/j.meegid.2016.04.024>.
- Skurnik, D., Roux, D., Cattoir, V., Danilchanka, O., Lu, X., Yoder-Himes, D.R., Han, K., Guillard, T., Jiang, D., Gaultier, C., Guerin, F., Aschard, H., Leclercq, R., Mekalanos, J.J., Lory, S., Pier, G.B., 2013. Enhanced in vivo fitness of carbapenem-resistant oprD mutants of *Pseudomonas aeruginosa* revealed through high-throughput sequencing. *Proc. Natl. Acad. Sci. U. S. A.* 110, 20747–20752. <https://doi.org/10.1073/pnas.1221552110>.
- Smillie, C., Garcillán-Barcia, M.P., Francia, M.V., Rocha, E.P.C., de la Cruz, F., 2010. Mobility of plasmids. *Microbiol. Mol. Biol. Rev.* 74, 434–452. <https://doi.org/10.1128/MMBR.00020-10>.
- Smith, J., 2011. Superinfection drives virulence evolution in experimental populations of bacteria and plasmids. *Evolution (N. Y.)* 65, 831–841. <https://doi.org/10.1111/j.1558-5646.2010.01178.x>.
- Snesrud, E., Maybank, R., Kwak, Y.I., Jones, A.R., Hinkle, M.K., McGann, P., 2018. Chromosomally encoded mcr-5 in colistin-nonsusceptible *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 62, e00679–e00718. <https://doi.org/10.1128/AAC.00679-18>.
- Sommer, M.O.A., Dantas, G., Church, G.M., 2009. Functional characterization of the antibiotic resistance reservoir in the human microflora. *Science* 325, 1128–1131. <https://doi.org/10.1126/science.1176950>.
- Sommer, M.O.A., Munck, C., Toft-Kehler, R.V., Andersson, D.I., 2017. Prediction of antibiotic resistance: time for a new preclinical paradigm? *Nat. Rev. Microbiol.* 15, 689–696. <https://doi.org/10.1038/nrmicro.2017.75>.
- Soucy, S.M., Huang, J., Gogarten, J.P., 2015. Horizontal gene transfer: building the web of life. *Nat. Rev. Genet.* 16, 472–482. <https://doi.org/10.1038/nrg3962>.
- Spalding, C., Keen, E., Smith, D.J., Krachler, A.-M., Jabbari, S., 2018. Mathematical modelling of the antibiotic-induced morphological transition of *Pseudomonas aeruginosa*. *PLOS Comput. Biol.* 14, e1006012. <https://doi.org/10.1371/journal.pcbi.1006012>.
- Starikova, I., Al-Haroni, M., Werner, G., Roberts, A.P., Sorum, V., Nielsen, K.M., Johnsen, P.J., 2013. Fitness costs of various mobile genetic elements in *Enterococcus faecium* and *Enterococcus faecalis*. *J. Antimicrob. Chemother.* 68, 2755–2765. <https://doi.org/10.1093/jac/dkt270>.
- Stevenson, C., Hall, J.P.J., Brockhurst, M.A., Harrison, E., 2018. Plasmid stability is enhanced by higher-frequency pulses of positive selection. *Proc. Biol. Sci.* 285. <https://doi.org/10.1098/rspb.2017.2497>.
- Stokes, H.W., Martinez, E., Chowdhury, P.R., Djordjevic, S., 2012. Class 1 integron-associated spread of resistance regions in *Pseudomonas aeruginosa*: plasmid or chromosomal platforms? *J. Antimicrob. Chemother.* 67, 1799–1800. <https://doi.org/10.1093/jac/dks116>.
- Stover, C.K., Pham, X.Q., Erwin, A.L., Mizoguchi, S.D., Warrenner, P., Hickey, M.J., Brinkman, F.S.L., Hufnagle, W.O., Kowalik, D.J., Lagrou, M., Garber, R.L., Goltry, L., Tolentino, E., Westbrook-Wadman, S., Yuan, Y., Brody, L.L., Coulter, S.N., Folger, K.R., Kas, A., Larbig, K., Lim, R., Smith, K., Spencer, D., Wong, G.K.-S., Wu, Z., Paulsen, I.T., Reizer, J., Saier, M.H., Hancock, R.E.W., Lory, S., Olson, M.V., 2000.

- Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an opportunistic pathogen. *Nature* 406, 959–964. <https://doi.org/10.1038/35023079>.
- Su, M., Satola, S.W., Read, T.D., 2018. Genome-based prediction of bacterial antibiotic resistance. *J. Clin. Microbiol.* 57, e01405-18. <https://doi.org/10.1128/JCM.01405-18>.
- Subedi, D., Vijay, A.K., Kohli, G.S., Rice, S.A., Willcox, M., 2018. Nucleotide sequence analysis of NPS-1 β -lactamase and a novel integron (In1427)-carrying transposon in an MDR *Pseudomonas aeruginosa* keratitis strain. *J. Antimicrob. Chemother.* 73, 1724–1726. <https://doi.org/10.1093/jac/dky073>.
- Sun, F., Zhou, D., Wang, Q., Feng, J., Feng, W., Luo, W., Zhang, D., Liu, Y., Qiu, X., Yin, Z., Chen, W., Xia, P., 2016. The first report of detecting the blaSIM-2 gene and determining the complete sequence of the SIM-encoding plasmid. *Clin. Microbiol. Infect.* 22, 347–351. <https://doi.org/10.1016/j.cmi.2015.12.001>.
- Sundqvist, E., Geli, P., Andersson, D.I., Sjölund-Karlsson, M., Runeheger, A., Cars, H., Abelson-Storby, K., Cars, O., Kahlmeter, G., 2010. Little evidence for reversibility of trimethoprim resistance after a drastic reduction in trimethoprim use. *J. Antimicrob. Chemother.* 65, 350–360. <https://doi.org/10.1093/jac/dkp387>.
- Suzuki, M., Suzuki, S., Matsui, M., Hiraki, Y., Kawano, F., Shibayama, K., 2014. A subclass B3 metallo- β -lactamase found in *Pseudomonas alcaligenes*. *J. Antimicrob. Chemother.* 69, 1430–1432. <https://doi.org/10.1093/jac/dkt498>.
- Svara, F., Rankin, D.J., 2011. The evolution of plasmid-carried antibiotic resistance. *BMC Evol. Biol.* 11, 130. <https://doi.org/10.1186/1471-2148-11-130>.
- Tacconelli, E., Carrara, E., Savoldi, A., Harbarth, S., Mendelson, M., Monnet, D.L., Pulcini, C., Kahlmeter, G., Kluytmans, J., Carmeli, Y., Ouellette, M., Outtersson, K., Patel, J., Cavaleri, M., Cox, E.M., Houchens, C.R., Grayson, M.L., Hansen, P., Singh, N., Theuretzbacher, U., Magnini, N., Aboderin, A.O., Al-Abri, S.S., Awang Jalil, N., Benzonana, N., Bhattacharya, S., Brink, A.J., Burkert, F.R., Cars, O., Cornaglia, G., Dyar, O.J., Friedrich, A.W., Gales, A.C., Gandra, S., Giske, C.G., Goff, D.A., Goossens, H., Gottlieb, T., Guzman Blanco, M., Hryniewicz, W., Kattula, D., Jinks, T., Kanj, S.S., Kerr, L., Kieny, M.-P., Kim, Y.S., Kozlov, R.S., Labarca, J., Laxminarayan, R., Leder, K., Leibovici, L., Levy-Hara, G., Littman, J., Malhotra-Kumar, S., Manchanda, V., Moja, L., Ndoye, B., Pan, A., Paterson, D.L., Paul, M., Qiu, H., Ramon-Pardo, P., Rodríguez-Baño, J., Sanguinetti, M., Sengupta, S., Sharland, M., Si-Mehand, M., Silver, L.L., Song, W., Steinbakk, M., Thomsen, J., Thwaites, G.E., van der Meer, J.W., Van Kinh, N., Vega, S., Villegas, M.V., Wechsler-Fördös, A., Wertheim, H.F.L., Wesangula, E., Woodford, N., Yilmaz, F.O., Zorzet, A., 2018. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *Lancet Infect. Dis.* 18, 318–327. [https://doi.org/10.1016/S1473-3099\(17\)30753-3](https://doi.org/10.1016/S1473-3099(17)30753-3).
- Tada, T., Shimada, K., Satou, K., Hirano, T., Pokhrel, B.M., Sherchand, J.B., Kirikae, T., 2017. *Pseudomonas aeruginosa* clinical isolates in Nepal coproducing metallo- β -lactamases and 16S rRNA methyltransferases. *Antimicrob. Agents Chemother.* 61, e00694–e00717. <https://doi.org/10.1128/AAC.00694-17>.
- Talá, L., Fineberg, A., Kukura, P., Persat, A., 2019. *Pseudomonas aeruginosa* orchestrates twitching motility by sequential control of type IV pili movements. *Nat. Microbiol.* 4, 774–780. <https://doi.org/10.1038/s41564-019-0378-9>.
- Tazzyman, S.J., Bonhoeffer, S., 2014. Why there are no essential genes on plasmids. *Mol. Biol. Evol.* PLoS One 32, 3079–3088. <https://doi.org/10.1093/molbev/msu293>.
- Temkin, E., Fallach, N., Almogor, J., Gladstone, B.P., Tacconelli, E., Carmeli, Y., DRIVE-AB Consortium, 2018. Estimating the number of infections caused by metallo-resistant *Escherichia coli* and *Klebsiella pneumoniae* in 2014: a modelling study. *Lancet Glob. Heal.* 6, e969–e979. [https://doi.org/10.1016/S2214-109X\(18\)30278-X](https://doi.org/10.1016/S2214-109X(18)30278-X).
- Thaller, M.C., Borgianni, L., Di Lallo, G., Chong, Y., Lee, K., Dajcs, J., Stroman, D., Rossolini, G.M., 2011. Metallo-beta-lactamase production by *Pseudomonas otitidis*: a species-related trait. *Antimicrob. Agents Chemother.* 55, 118–123. <https://doi.org/10.1128/AAC.01062-10>.
- The Global Risks Report, 2017. [WWW Document]. URL <https://www.weforum.org/reports/the-global-risks-report-2017> (accessed 21.07.19).
- Thomas, C.M., Haines, A.S., 2004. Plasmids of the genus *Pseudomonas*. *Pseudomonas*. Springer US, Boston, MA, pp. 197–231. https://doi.org/10.1007/978-1-4419-9086-0_7.
- Tognon, M., Köhler, T., Luscher, A., van Delden, C., 2019. Transcriptional profiling of *Pseudomonas aeruginosa* and *Staphylococcus aureus* during in vitro co-culture. *BMC Genom.* 20, 30. <https://doi.org/10.1186/s12864-018-5398-y>.
- Toleman, M.A., Biedenbach, D., Bennett, D., Jones, R.N., Walsh, T.R., 2003. Genetic characterization of a novel metallo- β -lactamase gene, blaIMP-13, harboured by a novel Tn5051-type transposon disseminating carbapenemase genes in Europe: report from the SENTRY worldwide antimicrobial surveillance programme. *J. Antimicrob. Chemother.* 52, 583–590. <https://doi.org/10.1093/jac/dkg410>.
- Toleman, M.A., Walsh, T.R., 2011. Combinatorial events of insertion sequences and ICE in Gram-negative bacteria. *FEMS Microbiol. Rev.* 35, 912–935. <https://doi.org/10.1111/j.1574-6976.2011.00294.x>.
- Touchon, M., Hoede, C., Tenailon, O., Barbe, V., Baeriswyl, S., Bidet, P., Bingen, E., Bonacorsi, S., Bouchier, C., Bouvet, O., Calteau, A., Chiappello, H., Clermont, O., Cruveiller, S., Danchin, A., Diard, M., Dossat, C., Karoui, M.E., Frapy, E., Garry, L., Ghigo, J.M., Gilles, A.M., Johnson, J., Le Bouguédec, C., Lescat, M., Mangenot, S., Martinez-Jéhanne, V., Matic, I., Nassif, X., Oztas, S., Petit, M.A., Pichon, C., Rouy, Z., Ruf, C.S., Schneider, D., Tourret, J., Vacherie, B., Vallenet, D., Médigue, C., Rocha, E.P.C., Denamur, E., 2009. Organised genome dynamics in the *Escherichia coli* species results in highly diverse adaptive paths. *PLoS Genet.* 5, e1000344. <https://doi.org/10.1371/journal.pgen.1000344>.
- Treangen, T.J., Rocha, E.P.C., 2011. Horizontal transfer, not duplication, drives the expansion of protein families in prokaryotes. *PLoS Genet.* 7, e1001284. <https://doi.org/10.1371/journal.pgen.1001284>.
- Treepong, P., Kos, V.N., Guyeux, C., Blanc, D.S., Bertrand, X., Valot, B., Hocquet, D., 2018. Global emergence of the widespread *Pseudomonas aeruginosa* ST235 clone. *Clin. Microbiol. Infect.* 24, 258–266. <https://doi.org/10.1016/j.cmi.2017.06.018>.
- Trindade, S., Sousa, A., Xavier, K.B., Dionisio, F., Ferreira, M.G., Gordo, I., 2009. Positive epistasis drives the acquisition of multidrug resistance. *PLoS Genet.* 5, e1000578. <https://doi.org/10.1371/journal.pgen.1000578>.
- Tseng, S.-P., Tsai, J.-C., Teng, L.-J., Hsueh, P.-R., 2009. Dissemination of transposon Tn6001 in carbapenem-non-susceptible and extensively drug-resistant *Pseudomonas aeruginosa* in Taiwan. *J. Antimicrob. Chemother.* 64, 1170–1174. <https://doi.org/10.1093/jac/dkp341>.
- Tuller, T., Girshovich, Y., Sella, Y., Kreimer, A., Freilich, S., Kupiec, M., Gophna, U., Ruppin, E., 2011. Association between translation efficiency and horizontal gene transfer within microbial communities. *Nucleic Acids Res.* 39, 4743–4755. <https://doi.org/10.1093/nar/gkr054>.
- Turner, K.H., Everett, J., Trivedi, U., Rumbaugh, K.P., Whiteley, M., 2014. Requirements for *Pseudomonas aeruginosa* acute burn and chronic surgical wound infection. *PLoS Genet.* 10, e1004518. <https://doi.org/10.1371/journal.pgen.1004518>.
- Turton, J.F., Wright, L., Underwood, A., Witney, A.A., Chan, Y.-T., Al-Shahib, A., Arnold, C., Doumith, M., Patel, B., Planche, T.D., Green, J., Holliman, R., Woodford, N., 2015. High-resolution analysis by whole-genome sequencing of an international lineage (sequence type 111) of *Pseudomonas aeruginosa* associated with metallo-carbapenemases in the United Kingdom. *J. Clin. Microbiol.* 53, 2622–2631. <https://doi.org/10.1128/JCM.00505-15>.
- Uemura, S., Yokota, S.-I., Mizuno, H., Sakawaki, E., Sawamoto, K., Maekawa, K., Tanno, K., Mori, K., Asai, Y., Fujii, N., 2010. Acquisition of a transposon encoding extended-spectrum β -lactamase SHV-12 by *Pseudomonas aeruginosa* isolates during the clinical course of a burn patient. *Antimicrob. Agents Chemother.* 54, 3956–3959. <https://doi.org/10.1128/AAC.00110-10>.
- Urbanowicz, P., Izdebski, R., Baraniak, A., Żabicka, D., Ziolkowski, G., Hryniewicz, W., Gniadkowski, M., 2019. *Pseudomonas aeruginosa* with NDM-1, DIM-1 and PME-1 β -lactamases, and RmtD3 16S rRNA methylase, encoded by new genomic islands. *J. Antimicrob. Chemother.* <https://doi.org/10.1093/jac/dkz262>.
- Valot, B., Guyeux, C., Rolland, J.Y., Mazouzi, K., Bertrand, X., Hocquet, D., 2015. What it takes to be a *Pseudomonas aeruginosa*? The core genome of the opportunistic pathogen updated. *PLoS One* 10, e0126468. <https://doi.org/10.1371/journal.pone.0126468>.
- van Belkum, A., Soriaga, L.B., LaFave, M.C., Akella, S., Veyrieras, J.-B., Barbu, E.M., Shortridge, D., Blanc, B., Hannum, G., Zambardi, G., Miller, K., Enright, M.C., Mugnier, N., Brami, D., Schicklin, S., Felderman, M., Schwartz, A.S., Richardson, T.H., Peterson, T.C., Hubby, B., Cady, K.C., 2015. Phylogenetic distribution of CRISPR-Cas systems in antibiotic-resistant *Pseudomonas aeruginosa*. *MBio* 6, e01796–e01815. <https://doi.org/10.1128/mBio.01796-15>.
- van der Zee, A., Kraak, W.B., Burggraaf, A., Goossens, W.H.F., Pirovano, W., Ossewaarde, J.M., Tommassen, J., 2018. Spread of carbapenem resistance by transposition and conjugation among *Pseudomonas aeruginosa*. *Front. Microbiol.* 9, 2057. <https://doi.org/10.3389/fmicb.2018.02057>.
- Vernikos, G.S., Parkhill, J., 2008. Resolving the structural features of genomic islands: a machine learning approach. *Genome Res.* 18, 331–342. <https://doi.org/10.1101/gr.7004508>.
- Versporten, A., Zarb, P., Caniaux, I., Gros, M.-F., Drapier, N., Miller, M., Jarlier, V., Nathwani, D., Goossens, H., Global-PPS Network, 2018. Antimicrobial consumption and resistance in adult hospital inpatients in 53 countries: results of an internet-based global point prevalence survey. *Lancet Glob. Heal.* 6, e619–e629. [https://doi.org/10.1016/S2214-109X\(18\)30186-4](https://doi.org/10.1016/S2214-109X(18)30186-4).
- Vielva, L., de Toro, M., Lanza, V.F., de la Cruz, F., Berger, B., 2017. PLACNETw: a web-based tool for plasmid reconstruction from bacterial genomes. *Bioinformatics* 33, 3796–3798. <https://doi.org/10.1093/bioinformatics/btx462>.
- Vilacoba, E., Quiroga, C., Pistorio, M., Famiglietti, A., Rodríguez, H., Kovensky, J., Deraspe, M., Raymond, F., Roy, P.H., Centrón, D., 2014. A blaVIM-2 plasmid disseminating in extensively drug-resistant clinical *Pseudomonas aeruginosa* and *Serratia marcescens* isolates. *Antimicrob. Agents Chemother.* 58, 7017–7018. <https://doi.org/10.1128/AAC.02934-14>.
- Vogwill, T., MacLean, R.C., 2015. The genetic basis of the fitness costs of antimicrobial resistance: a meta-analysis approach. *Evol. Appl.* 8, 284–295. <https://doi.org/10.1111/eva.12202>.
- Wagelchner, N., Wright, G.D., 2017. Antibiotic resistance: it's bad, but why isn't it worse? *BMC Biol.* 15, 84. <https://doi.org/10.1186/s12915-017-0423-1>.
- Wang, Y., Lu, J., Mao, L., Li, J., Yuan, Z., Bond, P.L., Guo, J., 2018. Antiepileptic drug carbamazepine promotes horizontal transfer of plasmid-borne multi-antibiotic resistance genes within and across bacterial genera. *ISME J.* 13, 509–522. <https://doi.org/10.1038/s41396-018-0275-x>.
- Ward, H., Perron, G.G., Maclean, R.C., 2009. The cost of multiple drug resistance in *Pseudomonas aeruginosa*. *J. Evol. Biol.* 22, 997–1003. <https://doi.org/10.1111/j.1420-9101.2009.01712.x>.
- Wattam, A.R., Davis, J.J., Assaf, R., Boisvert, S., Brettin, T., Bun, C., Conrad, N., Dietrich, E.M., Disz, T., Gabbard, J.L., Gerdes, S., Henry, C.S., Kenyon, R.W., Machi, D., Mao, C., Nordberg, E.K., Olsen, G.J., Murphy-Olson, D.E., Olson, R., Overbeek, R., Parrello, B., Pusch, G.D., Shukla, M., Vonstein, V., Warren, A., Xia, F., Yoo, H., Stevens, R.L., 2017. Improvements to PATRIC, the all-bacterial bioinformatics database and analysis resource center. *Nucleic Acids Res.* 45, D535–D542. <https://doi.org/10.1093/nar/gkw1017>.
- Wein, T., Hülter, N.F., Mizrahi, I., Dagan, T., 2019. Emergence of plasmid stability under non-selective conditions maintains antibiotic resistance. *Nat. Commun.* 10, 2595. <https://doi.org/10.1038/s41467-019-10600-7>.
- Weinert, L.A., Welch, J.J., 2017. Why might bacterial pathogens have small genomes?

- Trends Ecol. Evol. 32, 936–947. <https://doi.org/10.1016/j.tree.2017.09.006>.
- Winsor, G.L., Griffiths, E.J., Lo, R., Dhillon, B.K., Shay, J.A., Brinkman, F.S.L., 2016. Enhanced annotations and features for comparing thousands of *Pseudomonas* genomes in the *Pseudomonas* genome database. Nucleic Acids Res. 44, D646–D653. <https://doi.org/10.1093/nar/gkv1227>.
- Winstanley, C., Langille, M.G.I., Fothergill, J.L., Kukavica-Ibrulj, I., Paradis-Bleau, C., Sanschagrin, F., Thomson, N.R., Winsor, G.L., Quail, M.A., Lennard, N., Bignell, A., Clarke, L., Seeger, K., Saunders, D., Harris, D., Parkhill, J., Hancock, R.E.W., Brinkman, F.S.L., Levesque, R.C., 2009. Newly introduced genomic prophage islands are critical determinants of in vivo competitiveness in the Liverpool Epidemic Strain of *Pseudomonas aeruginosa*. Genome Res. 19, 12–23. <https://doi.org/10.1101/gr.086082.108>.
- Wistrand-Yuen, E., Knopp, M., Hjort, K., Koskiniemi, S., Berg, O.G., Andersson, D.I., 2018. Evolution of high-level resistance during low-level antibiotic exposure. Nat. Commun. 9, 1599. <https://doi.org/10.1038/s41467-018-04059-1>.
- Wong, A., Kassen, R., 2011. Parallel evolution and local differentiation in quinolone resistance in *Pseudomonas aeruginosa*. Microbiology 157, 937–944. <https://doi.org/10.1099/mic.0.046870-0>.
- Woodford, N., Turton, J.F., Livermore, D.M., 2011. Multiresistant Gram-negative bacteria: the role of high-risk clones in the dissemination of antibiotic resistance. FEMS Microbiol. Rev. 35, 736–755. <https://doi.org/10.1111/j.1574-6976.2011.00268.x>.
- World Health Organization (WHO), 2015. Global Action Plan on Antimicrobial Resistance. WHO. <https://www.who.int/antimicrobial-resistance/global-action-plan/en/>.
- Wozniak, R.A.F., Waldor, M.K., 2010. Integrative and conjugative elements: mosaic mobile genetic elements enabling dynamic lateral gene flow. Nat. Rev. Microbiol. 8, 552–563. <https://doi.org/10.1038/nrmicro2382>.
- Wurtzel, O., Yoder-Himes, D.R., Han, K., Dandekar, A.A., Edelheit, S., Greenberg, E.P., Sorek, R., Lory, S., 2012. The single-nucleotide resolution transcriptome of *Pseudomonas aeruginosa* grown in body temperature. PLoS Pathog. 8, e1002945. <https://doi.org/10.1371/journal.ppat.1002945>.
- Xiong, J., Alexander, D.C., Ma, J.H., Déraspe, M., Low, D.E., Jamieson, F.B., Roy, P.H., 2013. Complete sequence of pOZ176, a 500-kilobase IncP-2 plasmid encoding IMP-9-mediated carbapenem resistance, from outbreak isolate *Pseudomonas aeruginosa* 96. Antimicrob. Agents Chemother. 57, 3775–3782. <https://doi.org/10.1128/AAC.00423-13>.
- Yen, P., Papin, J.A., 2017. History of antibiotic adaptation influences microbial evolutionary dynamics during subsequent treatment. PLOS Biol. 15, e2001586. <https://doi.org/10.1371/journal.pbio.2001586>.
- Yokoyama, M., Stevens, E., Laabei, M., Bacon, L., Heesom, K., Bayliss, S., Ooi, N., O'Neill, A.J., Murray, E., Williams, P., Lubben, A., Reeksting, S., Meric, G., Pascoe, B., Sheppard, S.K., Recker, M., Hurst, L.D., Massey, R.C., 2018. Epistasis analysis uncovers hidden antibiotic resistance-associated fitness costs hampering the evolution of MRSA. Genome Biol. 19, 94. <https://doi.org/10.1186/s13059-018-1469-2>.
- Yoon, S.-H., Ha, S.-M., Kwon, S., Lim, J., Kim, Y., Seo, H., Chun, J., 2017. Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. Int. J. Syst. Evol. Microbiol. 67, 1613–1617. <https://doi.org/10.1099/ijsem.0.001755>.
- Zhan, Z., Hu, L., Jiang, X., Zeng, L., Feng, J., Wu, W., Chen, W., Yang, H., Yang, W., Gao, B., Yin, Z., Zhou, D., 2018. Plasmid and chromosomal integration of four novel blaIMP-carrying transposons from *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* and an *Enterobacter* sp. J. Antimicrob. Chemother. 73, 3005–3015. <https://doi.org/10.1093/jac/dky288>.
- Zwanzig, M., Harrison, E., Brockhurst, M.A., Hall, J.P.J., Berendonk, T.U., Berger, U., 2019. Mobile compensatory mutations promote plasmid survival. mSystems 4, e00186–e00218. <https://doi.org/10.1128/mSystems.00186-18>.