
2012-2013 Influenza Report Week 01

December 30 - January 5, 2013

About our flu activity reporting

MSDH relies upon selected sentinel health practitioners across the state to report the percentage of non-trauma visits consistent with an influenza-like illness (ILI: fever of 100°F or higher AND cough and/or sore throat). Also, providers are supplied with specimen collection kits. Samples are submitted to the Mississippi Public Health Laboratory for influenza PCR testing. Reports are used to estimate the state's ILI rate and the magnitude of the state's influenza activity. Reports represent only the distribution of flu in the state, not an actual count of all flu cases statewide. **Information is provisional only and may change depending on additional reporting from sentinel providers.**

State ILI Surveillance



For week 01 (12/30/2012-01/05/2013), the overall state ILI rate (9.4%) decreased from last week (13.3%), but was higher than this time last year (3.6%). |
Figure 1

Total number of non-trauma patients treated by sentinel providers in the last three weeks. | Table 1


2012-2013 Influenza Season

CDC Week	Week Ending	Non-trauma patients	ILI symptoms	ILI Rate (%)
01	Jan. 5	14288	1338	9.4%
52	Dec. 29	13767	1831	13.3%
51	Dec. 22	13314	1724	12.9%

During week **01**, **one** district (3) had an increase in ILI activity, while **six** districts (2, 4, 6, 7, 8, and 9)

had a decrease. **Two** districts (1 and 5) remained about the same. *Information is provisional only and may change depending on additional reporting from sentinel providers.* / **Table 2**


MSDH District ILI Rates (%) 2012-2013				
District	Week 52	Week 01		
State	13.3	9.4		
1	6.4	6.0		
2	17.7	13.8		
3	3.3	10.2		
4	23.4	13.8		
5	4.9	4.1		
6	23.4	13.7		
7	15.4	7.8		
8	11.1	8.8		
9	11.1	9.4		




Since week
35, the
number of
reported ILI
cases has
been highest
among people
in the 5 to 24
years of age
group,
followed by
those in the 0

to 4 years of age group. During Week **01**, the number of reported ILI cases was highest among individuals in the **0 to 4 years** of age group. | **Figure 2**

Figure 3




Mississippi ILI Rates 2010-2013 | Figure 4



Flu testing Reports

From week **35** (week ending September 1st) through week **01** (week ending January 5th), **108** positive influenza samples were identified by MSDH. 102 were influenza A and 6 were influenza B. Ninety-six were subtyped as influenza A (H3) and five were subtyped as influenza A (H1). One sample did not have subtyping performed.


The influenza cases were identified from the following counties: Calhoun, Carroll, Chickasaw (4), Clarke (2), Copiah (2), Covington (4), DeSoto, Harrison (8), Hinds (8), Itawamba, Jefferson Davis (5), Kemper (2), Lamar (5), Lauderdale, Lawrence, Lee (3), Leflore (3), Marion (4), Marshall (14), Monroe (6), Neshoba (2), Oktibbeha (6), Rankin (5), Sharkey, Tate (4), Tishomingo (2), Union (2), Washington (7), and Yazoo (3). | Figure 5




National ILI Surveillance

Nationally, there were **two** influenza-associated pediatric deaths reported to CDC during week **01**. One death was associated with an influenza A (H3) virus and occurred during week 52 (week ending December 29th). The second death was associated with an influenza A virus for which the subtype was not determined and occurred during week 01 (week ending January 5th). The total number of influenza-associated pediatric deaths reported during the 2012-2013 season is **20**. Additional data can be found at: http://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html. | **Figure 6**

Mississippi has had no reported influenza-associated pediatric deaths during this influenza season.






Mississippi reported "Regional" for week 01. | Table 4

Level of Flu Activity	Definition
No Activity	Overall clinical activity remains low and there are no lab confirmed cases.
Sporadic	Isolated cases of lab confirmed influenza in the state; ILI activity is not increased <u>OR</u> A lab-confirmed outbreak in a single institution in the state; ILI activity is not increased.
Local	Increased ILI within a single region AND recent (within the past 3 weeks) laboratory evidence of influenza in that region. ILI activity in other regions is not increased <u>OR</u> two of more institutional outbreaks (ILI or lab confirmed) within a single region AND recent (within the past 3 weeks) lab confirmed influenza in that region. Other regions do not have increased ILI and virus activity is no greater than sporadic in those regions
Regional	Increased ILI in at least 2 regions but fewer than half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the affected regions <u>OR</u> Institutional outbreaks (ILI or lab confirmed) in at least 2 regions but fewer than half of the regions AND recent lab confirmed influenza in the affected regions.
Widespread	Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the state.

2012- 2013 Influenza Season | Week 01 Influenza Report| December 30, 2012 - January 5, 2013 During week **01**, influenza activity remained elevated in the United States, but may be decreasing in some areas. | **Figure 8**



¹For up-to-date information on flu activity nationwide, please refer to the CDC's website: http://www.cdc.gov/flu/weekly/fluactivitysurv.htm.

Additional information:

Centers for Disease Control and Prevention	http://cdc.gov/flu/
Flu.gov	http://www.flu.gov/
MSDH Flu and Pneumonia	http://msdh.ms.gov/msdhsite/_static/14,0,199.html
Google Flu Trends	http://www.google.org/flutrends/
World Health Organization FluNet	http://www.who.int/influenza/gisrs_laboratory/flunet/en/

Page intentionally blank

Appendix

Figure 1






Figure 2












Figure 5




Figure 6




Figure 7




Figure 8

A Weekly Influenza Surveillance Report Prepared by the Influenza Division Weekly Influenza Activity Estimates Reported by State and Territorial Epidemiologists* Week Ending January 05, 2013- Week 1 District of Columbia ■ No Report Guam / No Activity Sporadic Local Regional ■ Widespread Alaska **US Virgin Islands** Puerto Rico Hawaii

^{*}This map indicates geographic spread and does not measure the severity of influenza activity.