2013-2014 Influenza Report


Week 40

September 29 – October 5, 2013

About our flu activity reporting

MSDH relies upon selected sentinel health practitioners across the state to report the percentage of total patient visits consistent with an influenza-like illness (ILI: fever of 100°F or higher AND cough and/or sore throat). Also, providers are supplied with specimen collection kits. Samples are submitted to the Mississippi Public Health Laboratory for influenza PCR testing. Reports are used to estimate the state's ILI rate and the magnitude of the state's influenza activity. Reports represent only the distribution of flu in the state, not an actual count of all flu cases statewide. *Information is provisional only and may change depending on additional reporting from sentinel providers.*

State ILI Surveillance



For week 40 (09/29/2013-10/05/2013), the overall state ILI rate (3.1%) was comparable to last week (2.6%) and to this time last year (3.3%). | Figure 1

Total number of patients treated by sentinel providers in the last three weeks. | Table 1


2013-2014 Influenza Season

| CDC Week | Week Ending | Total patients | ILI symptoms | ILI Rate (%) |
|----------|-------------|----------------|--------------|--------------|
| 40 | Oct. 5 | 16768 | 514 | 3.1% |
| 39 | Sept. 28 | 16915 | 438 | 2.6% |
| 38 | Sept. 21 | 17542 | 478 | 2.7% |

During week 40, four districts (2, 6, 8 and 9) had a slight increase in ILI activity, while five districts (1,


3, 4, 5 and 7) remained about the same. *Information is* provisional only and may change depending on additional reporting from sentinel providers. | **Table 2**

| MSDH District ILI Rates (%) 2013-2014 | | | | | |
|---------------------------------------|---------|---------|--|--|--|
| District | Week 39 | Week 40 | | | |
| State | 2.6 | 3.1 | | | |
| 1 | 1.8 | 1.6 | | | |
| 2 | 1.5 | 2.3 | | | |
| 3 | 6.2 | 5.7 | | | |
| 4 | 2.4 | 2.7 | | | |
| 5 | 0.3 | 0.3 | | | |
| 6 | 9.6 | 10.3 | | | |
| 7 | 2.4 | 2.2 | | | |
| 8 | 1.1 | 2.3 | | | |
| 9 | 3.1 | 4.0 | | | |




years of age groups. During week **40**, the percentage of reported ILI cases was highest among individuals in the **5 to 24 years** of age group. | <u>Figure 2</u>

2013- 2014 Influenza Season | Week 40 Influenza Report| Sept. 29 – Oct. 5, 2013 The 2013-14 state ILI rate was **above** the national and Region 4 baselines, but was **below** the state baseline for week **40**. | Figure 3



Mississippi ILI Rates 2010-2013 | Figure 4



Flu Testing Reports

Since week **40** (week ending October 5th), **3** positive influenza samples were identified by MSDH. All three samples were identified as influenza A (H1).

The influenza cases were identified from the following counties: Forrest (1) and Washington (2).

National and Mississippi Pediatric Mortality Surveillance

National data not currently available due to the Federal shutdown.

Mississippi has had **no** influenza-associated pediatric deaths reported during this influenza season.

National ILI Surveillance

Mississippi reported "Sporadic" flu activity for week 40. | Table 3

| Level of Flu Activity | Definition |
|--------------------------|--|
| No Activity | Overall clinical activity remains low and there are no lab confirmed cases. |
| Sporadic | Isolated cases of lab confirmed influenza in the state; ILI activity is not increased <u>OR</u> A lab-confirmed outbreak in a single institution in the state; ILI activity is not increased. |
| Local | Increased ILI within a single region AND recent (within the past 3 weeks) laboratory evidence of influenza in that region. ILI activity in other regions is not increased <u>OR</u> two of more institutional outbreaks (ILI or lab confirmed) within a single region AND recent (within the past 3 weeks) lab confirmed influenza in that region. Other regions do not have increased ILI and virus activity is no greater than sporadic in those regions |
| Regional | Increased ILI in at least 2 regions but fewer than half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the affected regions <u>OR</u> Institutional outbreaks (ILI or lab confirmed) in at least 2 regions but fewer than half of the regions AND recent lab confirmed influenza in the affected regions. |
| Widespread | Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the state. |

Additional information:

| Centers for Disease Control and Prevention | http://cdc.gov/flu/ |
|--|--|
| Flu.gov | http://www.flu.gov/ |
| MSDH Flu and Pneumonia | http://msdh.ms.gov/msdhsite/_static/14,0,199.html |
| Google Flu Trends | http://www.google.org/flutrends/ |
| World Health Organization FluNet | http://www.who.int/influenza/gisrs_laboratory/flunet/en/ |

2013- 2014 Influenza Season | Week 40 Influenza Report| Sept. 29 – Oct. 5, 2013

Page intentionally blank

Appendix

Figure 1




Figure 2




Figure 3




Figure 4

