



planetmath.org

Math for the people, by the people.

time invariant

Canonical name	TimeInvariant
Date of creation	2013-03-22 15:02:14
Last modified on	2013-03-22 15:02:14
Owner	Mathprof (13753)
Last modified by	Mathprof (13753)
Numerical id	5
Author	Mathprof (13753)
Entry type	Definition
Classification	msc 00A05
Related topic	AutonomousSystem
Defines	time-invariant
Defines	shift-invariant

A dynamical system is **time-invariant** if its generating formula is dependent on state only, and independent of time. A synonym for time-invariant is autonomous. The complement of time-invariant is time-varying (or nonautonomous).

For example, the continuous-time system $\dot{x} = f(x, t)$ is time-invariant if and only if $f(x, t_1) \equiv f(x, t_2)$ for all valid states x and times t_1 and t_2 . Thus $\dot{x} = \sin x$ is time-invariant, while $\dot{x} = \frac{\sin x}{1+t}$ is time-varying.

Likewise, the discrete-time system $x[n] = f[x, n]$ is time-invariant (also called shift-invariant) if and only if $f[x, n_1] \equiv f[x, n_2]$ for all valid states x and time indices n_1 and n_2 . Thus $x[n] = 2x[n - 1]$ is time-invariant, while $x[n] = 2nx[n - 1]$ is time-varying.