



planetmath.org

Math for the people, by the people.

property

Canonical name	Property
Date of creation	2013-03-22 14:01:29
Last modified on	2013-03-22 14:01:29
Owner	drini (3)
Last modified by	drini (3)
Numerical id	15
Author	drini (3)
Entry type	Definition
Classification	msc 00A05
Synonym	attribute
Synonym	propositional function
Related topic	Subset
Related topic	CharacteristicFunction
Related topic	Relation
Related topic	ClosureOfARelationWithRespectToAProperty
Defines	unary relation
Defines	predicate

Let X be a set. A *property* p of X is a function

$$p: X \rightarrow \{\text{true}, \text{false}\}.$$

An element $x \in X$ is said to *have* or *does not have the property* p depending on whether $p(x) = \text{true}$ or $p(x) = \text{false}$. Any property gives rise in a natural way to the set

$$X(p) := \{x \in X \mid x \text{ has property } p\}$$

and the corresponding <http://planetmath.org/node/CharacteristicFunction> characteristic function $1_{X(p)}$. The identification of p with $X(p) \subseteq X$ enables us to think of a property of X as a 1-ary, or a *unary relation* on X . Therefore, one may treat all these notions equivalently.

Usually, a property p of X can be identified with a so-called *propositional function*, or a *predicate* $\varphi(v)$, where v is a variable or a tuple of variables whose values range over X . The values of a propositional function is a proposition, which can be interpreted as being either “true” or “false”, so that $X(p) = \{x \mid \varphi(x) \text{ is true}\}$.

Below are a few examples:

- Let $X = \mathbb{Z}$. Let $\varphi(v)$ be the propositional function “ v is divisible by 3”. If p is the property identified with $\varphi(v)$, then $X(p) = 3\mathbb{Z}$.
- Again, let $X = \mathbb{Z}$. Let $\varphi(v_1, v_2) :=$ “ v_1 is divisible by v_2 ” and p the corresponding property. Then

$$X(p) = \{(m, n) \mid m = np, \text{ for some } p \in \mathbb{Z}\},$$

which is a subset of $X \times X$. So p is a property of $X \times X$.

- The reflexive property of a binary relation on X can be identified with the propositional function $\varphi(V) :=$ “ $\forall a \in X, (a, a) \in V$ ”, and therefore

$$X(p) = \{R \subseteq X \times X \mid \varphi(R) \text{ is true}\},$$

which is a subset of $2^{X \times X}$. Thus, p is a property of $2^{X \times X}$.

- In point set topology, we often encounter the finite intersection property on a family of subsets of a given set X . Let

$$\varphi(\mathcal{V}) := \text{“}\forall n \in \mathbb{N}, \forall E_1 \in \mathcal{V}, \dots, \forall E_n \in \mathcal{V}, \exists x \in X (x \in E_1 \cap \dots \cap E_n)\text{”}$$

and p the corresponding property, then

$$X(p) = \{\mathcal{F} \subseteq 2^X \mid \varphi(\mathcal{F}) \text{ is } \textit{true}\},$$

which is a subset of 2^{2^X} . Thus p is a property of 2^{2^X} .