



planetmath.org

Math for the people, by the people.

satisfaction relation

Canonical name	SatisfactionRelation
Date of creation	2013-03-22 12:43:56
Last modified on	2013-03-22 12:43:56
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	14
Author	CWoo (3771)
Entry type	Definition
Classification	msc 03C07
Related topic	Model
Defines	satisfaction
Defines	satisfy

Alfred Tarski was the first mathematician to give a formal definition of what it means for a formula to be “true” in a structure. To do this, we need to provide a meaning to terms, and truth-values to the formulas. In doing this, free variables cause a problem : what value are they going to have ? One possible answer is to supply temporary values for the free variables, and define our notions in terms of these temporary values.

Let  $\mathcal{A}$  be a structure with signature  $\tau$ . Suppose  $\mathcal{I}$  is an interpretation, and  $\sigma$  is a function that assigns elements of  $A$  to variables, we define the function  $\text{Val}_{\mathcal{I},\sigma}$  inductively on the construction of terms :

$$\begin{aligned}\text{Val}_{\mathcal{I},\sigma}(c) &= \mathcal{I}(c) & c \text{ a constant symbol} \\ \text{Val}_{\mathcal{I},\sigma}(x) &= \sigma(x) & x \text{ a variable} \\ \text{Val}_{\mathcal{I},\sigma}(f(t_1, \dots, t_n)) &= \mathcal{I}(f)(\text{Val}_{\mathcal{I},\sigma}(t_1), \dots, \text{Val}_{\mathcal{I},\sigma}(t_n)) & f \text{ an } n\text{-ary function symbol}\end{aligned}$$

Now we are set to define satisfaction. Again we have to take care of free variables by assigning temporary values to them via a function  $\sigma$ . We define the relation  $\mathcal{A}, \sigma \models \varphi$  by induction on the construction of formulas :

$$\begin{aligned}\mathcal{A}, \sigma &\models t_1 = t_2 \text{ if and only if } \text{Val}_{\mathcal{I},\sigma}(t_1) = \text{Val}_{\mathcal{I},\sigma}(t_2) \\ \mathcal{A}, \sigma &\models R(t_1, \dots, t_n) \text{ if and only if } (\text{Val}_{\mathcal{I},\sigma}(t_1), \dots, \text{Val}_{\mathcal{I},\sigma}(t_n)) \in \mathcal{I}(R) \\ \mathcal{A}, \sigma &\models \neg\varphi \text{ if and only if } \mathcal{A}, \sigma \not\models \varphi \\ \mathcal{A}, \sigma &\models \varphi \vee \psi \text{ if and only if either } \mathcal{A}, \sigma \models \varphi \text{ or } \mathcal{A}, \sigma \models \psi \\ \mathcal{A}, \sigma &\models \exists x.\varphi(x) \text{ if and only if for some } a \in A, \mathcal{A}, \sigma[x/a] \models \varphi\end{aligned}$$

Here

$$\sigma[x/a](y) \begin{cases} a & \text{if } x = y \\ \sigma(y) & \text{else.} \end{cases}$$

In case for some  $\varphi$  of  $L$ , we have  $\mathcal{A}, \sigma \models \varphi$ , we say that  $\mathcal{A}$  **models**, or **is a model of**, or **satisfies**  $\varphi$ . If  $\varphi$  has the free variables  $x_1, \dots, x_n$ , and  $a_1, \dots, a_n \in A$ , we also write  $\mathcal{A} \models \varphi(a_1, \dots, a_n)$  or  $\mathcal{A} \models \varphi(a_1/x_1, \dots, a_n/x_n)$  instead of  $\mathcal{A}, \sigma[x_1/a_1] \cdots [x_n/a_n] \models \varphi$ . In case  $\varphi$  is a sentence (formula with no free variables), we write  $\mathcal{A} \models \varphi$ .