



negative translation

Canonical name	NegativeTranslation
Date of creation	2013-03-22 19:35:48
Last modified on	2013-03-22 19:35:48
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	9
Author	CWoo (3771)
Entry type	Definition
Classification	msc 03B20
Classification	msc 03F55
Synonym	double negative translation
Defines	Kolmogorov negative translation
Defines	Gödel negative translation
Defines	Kuroda negative translation
Defines	Krivine negative translation

It is well-known that classical propositional logic PL_c can be considered as a subsystem of intuitionistic propositional logic PL_i by translating any wff A in PL_c into $\neg\neg A$ in PL_i . According to Glivenko's theorem, A is a theorem of PL_c iff $\neg\neg A$ is a theorem of PL_i . This translation, however, fails to preserve theoremhood in the corresponding predicate logics. For example, if A is of the form $\exists xB$, then $\vdash_c A$ no longer implies $\vdash_i \neg\neg A$. A number of translations have been devised to overcome this defect. They are collectively known as *negative translations* or *double negative translations* of classical logic into intuitionistic logic. Below is a list of the most commonly mentioned negative translations:

- *Kolmogorov negative translation:*

- | | |
|---|---|
| 1. $p^\circ := \neg\neg p$ with p atomic and $\perp^\circ := \perp$ | 4. $(A \rightarrow B)^\circ := \neg\neg(A^\circ \rightarrow B^\circ)$ |
| 2. $(A \wedge B)^\circ := \neg\neg(A^\circ \wedge B^\circ)$ | 5. $(\forall xA)^\circ := \neg\neg\forall xA^\circ$ |
| 3. $(A \vee B)^\circ := \neg\neg(A^\circ \vee B^\circ)$ | 6. $(\exists xA)^\circ := \neg\neg\exists xA^\circ$ |

- *God el negative translation:*

- | | |
|---|---|
| 1. $p^- := \neg\neg p$ with p atomic and $\perp^- := \perp$ | 4. $(A \rightarrow B)^- := A^- \rightarrow B^-$ |
| 2. $(A \wedge B)^- := A^- \wedge B^-$ | 5. $(\forall xA)^- := \forall xA^-$ |
| 3. $(A \vee B)^- := \neg\neg(A^- \vee B^-)$ | 6. $(\exists xA)^- := \neg\neg\exists xA^-$ |

- *Kuroda negative translation:*

- | | |
|--|---|
| 1. $p_u := p$ with p atomic and $\perp_u := \perp$ | 4. $(A \rightarrow B)_u := A_u \rightarrow B_u$ |
| 2. $(A \wedge B)_u := A_u \wedge B_u$ | 5. $(\forall xA)_u := \forall x\neg\neg A_u$ |
| 3. $(A \vee B)_u := A_u \vee B_u$ | 6. $(\exists xA)_u := \exists xA_u$ |

And then $A^u := \neg\neg A_u$.

- *Krivine negative translation:*

1. $p_r := \neg p$ with p atomic and $\perp_r := \neg \perp$
2. $(A \wedge B)_r := A_r \vee B_r$
3. $(A \vee B)_r := A_r \wedge B_r$
4. $(A \rightarrow B)_r := \neg A_r \wedge B_r$
5. $(\forall x A)_r := \exists x A_r$
6. $(\exists x A)_r := \neg \exists x \neg A_r$

And then $A^r := \neg A_r$.

Remark. It can be shown that for any wff A :

$$\vdash_i A^* \leftrightarrow A^\# \quad \text{and} \quad \vdash_c A \quad \text{iff} \quad \vdash_i A^*$$

where $*, \# \in \{\circ, -, u, r\}$.