



planetmath.org

Math for the people, by the people.

inverse function theorem

Canonical name	InverseFunctionTheorem
Date of creation	2013-03-22 12:58:30
Last modified on	2013-03-22 12:58:30
Owner	azdbacks4234 (14155)
Last modified by	azdbacks4234 (14155)
Numerical id	9
Author	azdbacks4234 (14155)
Entry type	Theorem
Classification	msc 03E20
Related topic	DerivativeOfInverseFunction
Related topic	LegendreTransform
Related topic	DerivativeAsParameterForSolvingDifferentialEquations
Related topic	TheoryForSeparationOfVariables

Let \mathbf{f} be a continuously differentiable, vector-valued function mapping the open set $E \subset \mathbb{R}^n$ to \mathbb{R}^n and let $S = \mathbf{f}(E)$. If, for some point $\mathbf{a} \in E$, the Jacobian, $|J_{\mathbf{f}}(\mathbf{a})|$, is non-zero, then there is a uniquely defined function \mathbf{g} and two open sets $X \subset E$ and $Y \subset S$ such that

1. $\mathbf{a} \in X$, $\mathbf{f}(\mathbf{a}) \in Y$;
2. $Y = \mathbf{f}(X)$;
3. $\mathbf{f} : X \rightarrow Y$ is one-one;
4. \mathbf{g} is continuously differentiable on Y and $\mathbf{g}(\mathbf{f}(\mathbf{x})) = \mathbf{x}$ for all $\mathbf{x} \in X$.

0.0.1 Simplest case

When $n = 1$, this theorem becomes: Let f be a continuously differentiable, real-valued function defined on the open interval I . If for some point $a \in I$, $f'(a) \neq 0$, then there is a neighbourhood $[\alpha, \beta]$ of a in which f is strictly monotonic. Then $y \rightarrow f^{-1}(y)$ is a continuously differentiable, strictly monotonic function from $[f(\alpha), f(\beta)]$ to $[\alpha, \beta]$. If f is increasing (or decreasing) on $[\alpha, \beta]$, then so is f^{-1} on $[f(\alpha), f(\beta)]$.

0.0.2 Note

The inverse function theorem is a special case of the implicit function theorem where the dimension of each variable is the same.