

planetmath.org

Math for the people, by the people.

function

Canonical name Function

Date of creation 2013-03-22 11:48:58 Last modified on 2013-03-22 11:48:58

Owner djao (24)Last modified by djao (24)

Numerical id 23

Author djao (24)
Entry type Definition
Classification msc 03E20
Classification msc 44A20
Classification msc 33E20
Classification msc 30D15

Synonym map Related topic Mapping

Related topic InjectiveFunction

Related topic Surjective
Related topic Bijection
Related topic Relation
Defines domain
Defines codomain
Defines composition

Defines image Defines range

Defines composite function

A function is a triplet (f, A, B) where:

- 1. A is a set (called the *domain* of the function).
- 2. B is a set (called the *codomain* of the function).
- 3. f is a binary relation between A and B.
- 4. For every $a \in A$, there exists $b \in B$ such that $(a, b) \in f$.
- 5. If $a \in A$, $b_1, b_2 \in B$, and $(a, b_1) \in f$ and $(a, b_2) \in f$, then $b_1 = b_2$.

The triplet (f, A, B) is usually written with the specialized notation $f: A \to B$. This notation visually conveys the fact that f maps elements of A into elements of B.

Other standard notations for functions are as follows:

- For $a \in A$, one denotes by f(a) the unique element $b \in B$ such that $(a,b) \in f$.
- The *image* of (f, A, B), denoted f(A), is the set

$$\{b \in B \mid f(a) = b \text{ for some } a \in A\}$$

consisting of all elements of B which equal f(a) for some element $a \in A$. Note that, by abuse of notation, the set f(A) is almost always called the image of f, rather than the image of (f, A, B).

- In cases where the function f is clear from context, the notation $a \mapsto b$ is equivalent to the statement f(a) = b.
- Given two functions $f: A \to B$ and $g: B \to C$, there exists a unique function $g \circ f: A \to C$ satisfying the equation $g \circ f(a) = g(f(a))$. The function $g \circ f$ is called the *composition* of f and g, and a function constructed in this manner is called a *composite function*. Composition is associative, meaning that $h \circ (g \circ f) = (h \circ g) \circ f$ provided that either expression is defined.
- When a function $f: A \to A$ has its domain equal to its codomain, one often writes f^n for the *n*-fold composition

$$\underbrace{f \circ f \circ \cdots \circ f}_{n \text{ times}}$$

where n is any natural number. Occasionally this can be confused with ordinary exponentiation (for example the function $x \mapsto (\sin x)(\sin x)$ is conventionally written as \sin^2); in such cases one usually writes $f^{[n]}$ to denote the n-fold composition.

There is no universal agreement as to the definition of the *range* of a function. Some authors define the range of a function to be equal to the codomain, and others define the range of a function to be equal to the image.

Remark. In set theory, a function is defined as a relation f, such that whenever $(a,b), (a,c) \in f$, then b=c. Notice that the sets A,B are not specified in advance, unlike the defintion given in the beginning of the article. The *domain* and *range* of the function f is the domain and range of f as a relation. Using this definition of a function, we may recapture the defintion at the top of the entry by saying that a function f maps from a set A into a set B, if the domain of f is A, and the range of f is a subset of B.