



planetmath.org

Math for the people, by the people.

iterated forcing

Canonical name	IteratedForcing
Date of creation	2013-03-22 12:54:47
Last modified on	2013-03-22 12:54:47
Owner	Henry (455)
Last modified by	Henry (455)
Numerical id	5
Author	Henry (455)
Entry type	Definition
Classification	msc 03E35
Classification	msc 03E40
Defines	FS
Defines	CS
Defines	finite support
Defines	finite support iterated forcing
Defines	countable support
Defines	countable support iterated forcing
Defines	support iterated forcing

We can define an *iterated forcing* of length α by induction as follows:

Let $P_0 = \emptyset$.

Let \hat{Q}_0 be a forcing notion.

For $\beta \leq \alpha$, P_β is the set of all functions f such that $\text{dom}(f) \subseteq \beta$ and for any $i \in \text{dom}(f)$, $f(i)$ is a P_i -name for a member of \hat{Q}_i . Order P_β by the rule $f \leq g$ iff $\text{dom}(g) \subseteq \text{dom}(f)$ and for any $i \in \text{dom}(f)$, $g \restriction i \Vdash f(i) \leq_{\hat{Q}_i} g(i)$. (Translated, this means that any generic subset including g restricted to i forces that $f(i)$, an element of \hat{Q}_i , be less than $g(i)$.)

For $\beta < \alpha$, \hat{Q}_β is a forcing notion in P_β (so $\Vdash_{P_\beta} \hat{Q}_\beta$ is a forcing notion).

Then the sequence $\langle \hat{Q}_\beta \rangle_{\beta < \alpha}$ is an iterated forcing.

If P_β is restricted to finite functions that it is called a *finite support iterated forcing* (FS), if P_β is restricted to countable functions, it is called a *countable support iterated function* (CS), and in general if each function in each P_β has size less than κ then it is a *$< \kappa$ -support iterated forcing*.

Typically we construct the sequence of \hat{Q}_β 's by induction, using a function F such that $F(\langle \hat{Q}_\beta \rangle_{\beta < \gamma}) = \hat{Q}_\gamma$.