



planetmath.org

Math for the people, by the people.

ring of sets

Canonical name	RingOfSets
Date of creation	2013-03-22 15:47:46
Last modified on	2013-03-22 15:47:46
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	18
Author	CWoo (3771)
Entry type	Definition
Classification	msc 03E20
Classification	msc 28A05
Synonym	lattice of sets
Synonym	algebra of sets
Related topic	SigmaAlgebra
Related topic	AbsorbingElement
Related topic	RepresentingADistributiveLatticeByRingOfSets
Related topic	RepresentingABooleanLatticeByFieldOfSets
Defines	field of sets

Ring of Sets

Let S be a set and 2^S be the power set of S . A subset \mathcal{R} of 2^S is said to be a *ring of sets* of S if it is a lattice under the intersection and union operations. In other words, \mathcal{R} is a ring of sets if

- for any $A, B \in \mathcal{R}$, then $A \cap B \in \mathcal{R}$,
- for any $A, B \in \mathcal{R}$, then $A \cup B \in \mathcal{R}$.

A ring of sets is a distributive lattice. The word “ring” in the name has nothing to do with the ordinary ring found in algebra. Rather, it is an abelian semigroup with respect to each of the binary set operations. If $S \in \mathcal{R}$, then (\mathcal{R}, \cap, S) becomes an abelian monoid. Similarly, if $\emptyset \in \mathcal{R}$, then $(\mathcal{R}, \cup, \emptyset)$ is an abelian monoid. If both $S, \emptyset \in \mathcal{R}$, then $(\mathcal{R}, \cup, \cap)$ is a commutative semiring, since $\emptyset \cap A = A \cap \emptyset = \emptyset$, and \cap distributes over \cup . Dualizing, we see that $(\mathcal{R}, \cap, \cup)$ is also a commutative semiring. It is perhaps with this connection that the name “ring of sets” is so chosen.

Since S is not required to be in \mathcal{R} , a ring of sets can in theory be the empty set. Even if \mathcal{R} may be non-empty, it may be a singleton. Both cases are not very interesting to study. To avoid such examples, some authors, particularly measure theorists, define a ring of sets to be a non-empty set with the first condition above replaced by

- for any $A, B \in \mathcal{R}$, then $A - B \in \mathcal{R}$.

This is indeed a stronger condition, as $A \cap B = A - (A - B) \in \mathcal{R}$. However, we shall stick with the more general definition here.

Field of Sets

An even stronger condition is to insist that not only is \mathcal{R} non-empty, but that $S \in \mathcal{R}$. Such a ring of sets is called a field, or algebra of sets. Formally, given a set S , a *field of sets* \mathcal{F} of S satisfies the following criteria

- \mathcal{F} is a ring of sets of S ,
- $S \in \mathcal{F}$, and
- if $A \in \mathcal{F}$, then the complement $\overline{A} \in \mathcal{F}$.

The three conditions above are equivalent to the following three conditions:

- $\emptyset \in \mathcal{F}$,
- if $A, B \in \mathcal{F}$, then $A \cup B \in \mathcal{F}$, and
- if $A \in \mathcal{F}$, then $\overline{A} \in \mathcal{F}$.

A field of sets is also known as an *algebra of sets*.

It is easy to see that \mathcal{F} is a distributive complemented lattice, and hence a Boolean lattice. From the discussion earlier, we also see that \mathcal{F} (of S) is a commutative semiring, with S acting as the multiplicative identity and \emptyset both the additive identity and the multiplicative absorbing element.

Remark. Two remarkable theorems relating to of certain lattices as rings or fields of sets are the following:

1. a lattice is distributive iff it is <http://planetmath.org/LatticeIsomorphismLattice> isomorphic to a ring of sets (G. Birkhoff and M. Stone);
2. a lattice is <http://planetmath.org/BooleanLatticeBoolean> iff it is lattice to a field of sets (M. Stone).

References

- [1] P. R. Halmos: *Lectures on Boolean Algebras*, Springer-Verlag (1970).
- [2] P. R. Halmos: *Measure Theory*, Springer-Verlag (1974).
- [3] G. Grätzer: *General Lattice Theory*, Birkhäuser, (1998).