

planetmath.org

Math for the people, by the people.

polyadic algebra with equality

Canonical name PolyadicAlgebraWithEquality

Date of creation 2013-03-22 17:51:37 Last modified on 2013-03-22 17:51:37

Owner CWoo (3771) Last modified by CWoo (3771)

Numerical id 10

Author CWoo (3771)
Entry type Definition
Classification msc 03G15
Synonym equality algebra
Related topic CylindricAlgebra
Defines equality predicate

Defines substitutive
Defines reflexive
Defines symmetric
Defines transitive

Let $A = (B, V, \exists, S)$ be a polyadic algebra. An equality predicate on A is a function $E: V \times V \to B$ such that

- 1. $S(f) \circ E(x,y) = E(f(x),f(y))$ for any $f:V \to V$ and any $x,y \in V$
- 2. E(x,x) = 1 for every $x \in V$, and
- 3. $E(x,y) \wedge a \leq S(x/y)a$, where $a \in B$, and (x/y) denotes the function $V \to V$ that maps x to y, and constant everywhere else.

Heuristically, we can interpret the conditions above as follows:

- 1. if x = y and if we replace x by, say x_1 , and y by y_1 , then $x_1 = y_1$.
- 2. x = x for every variable x
- 3. if we have a propositional function a that is true, and x = y, then the proposition obtained from a by replacing all occurrences of x by y is also true.

The second condition is also known as the *reflexive property* of the equality predicate E, and the third is known as the *substitutive property* of E

A polyadic algebra with equality is a pair (A, E) where A is a polyadic algebra and E is an equality predicate on A. Paul Halmos introduced this concept and called this simply an equality algebra.

Below are some basic properties of the equality predicate E in an equality algebra (A, E):

- (symmetric property) $E(x, y) \leq E(y, x)$
- (transitive property) $E(x,y) \wedge E(y,z) \leq E(x,z)$
- $E(x,y) \land a = E(x,y) \land S(x,y)a$, where (x,y) in the S is the transposition on V that swaps x and y and leaves everything else fixed.
- if a variable $x \in V$ is not in the support of $a \in A$, then $a = \exists (x)(E(x,y) \land S(y/x)a)$.
- $\exists (x)(E(x,y) \land a) \land \exists (x)(E(x,y) \land a') = 0 \text{ for all } a \in A \text{ and all } x,y \in V$ whenever $x \neq y$.
- $\exists (x)(E(x,y) \land E(x,z)) = E(y,z) \text{ for all } x,y,z \in V \text{ where } x \notin \{y,z\}.$

Remarks

- The degree and local finiteness of a polyadic algebra (A, E) are defined as the degree and the local finiteness and degree of its underlying polyadic algebra A.
- It can be shown that every locally finite polyadic algebra of infinite degree can be embedded (as a polyadic subalgebra) in a locally finite polyadic algebra with equality of infinite degree.
- Like cylindric algebras, polyadic algebras with equality is an attempt at "converting" a first order logic (with equality) into algebraic form, so that the logic can be studied using algebraic means.

References

- [1] P. Halmos, Algebraic Logic, Chelsea Publishing Co. New York (1962).
- [2] B. Plotkin, *Universal Algebra, Algebraic Logic, and Databases*, Kluwer Academic Publishers (1994).