



planetmath.org

Math for the people, by the people.

discontinuity of characteristic function

Canonical name	DiscontinuityOfCharacteristicFunction
Date of creation	2015-02-03 21:23:33
Last modified on	2015-02-03 21:23:33
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	3
Author	pahio (2872)
Entry type	Theorem
Classification	msc 03-00
Classification	msc 26-00
Classification	msc 26A09

**Theorem.** For a subset  $A$  of  $\mathbb{R}^n$ , the set of the <http://planetmath.org/Continuousdiscontinuity> points of the characteristic function  $\chi_A$  is the <http://planetmath.org/BoundaryFrontierboundary> of  $A$ .

*Proof.* Let  $a$  be a discontinuity point of  $\chi_A$ . Then any <http://planetmath.org/Neighborhood> of  $a$  contains the points  $b$  and  $c$  such that  $\chi_A(b) = 1$  and  $\chi_A(c) = 0$ . Thus  $b \in A$  and  $c \notin A$ , whence  $a$  is a boundary point of  $A$ .

If, on the contrary,  $a$  is a boundary point of  $A$  and  $U(a)$  an arbitrary neighborhood of  $a$ , it follows that  $U(a)$  contains both points belonging to  $A$  and points not belonging to  $A$ . So we have in  $U(a)$  the points  $b$  and  $c$  such that  $\chi_A(b) = 1$  and  $\chi_A(c) = 0$ . This means that  $\chi_A$  cannot be continuous at the point  $a$  (N.B. that one does not need to know the value  $\chi_A(a)$ ).