



planetmath.org

Math for the people, by the people.

substitution theorem for propositional logic

Canonical name	SubstitutionTheoremForPropositionalLogic
Date of creation	2013-03-22 19:33:08
Last modified on	2013-03-22 19:33:08
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	17
Author	CWoo (3771)
Entry type	Result
Classification	msc 03B05
Related topic	AxiomSystemForPropositionalLogic
Defines	substitution theorem

In this entry, we will prove the substitution theorem for propositional logic based on the axiom system found <http://planetmath.org/AxiomSystemForPropositionalLogic>. Besides the deduction theorem, below are some additional results we will need to prove the theorem:

1. If $\Delta \vdash A \rightarrow B$ and $\Gamma \vdash B \rightarrow C$, then $\Delta, \Gamma \vdash A \rightarrow C$.
2. $\Delta \vdash A$ and $\Delta \vdash B$ iff $\Delta \vdash A \wedge B$.
3. $\vdash A \leftrightarrow A$.
4. $\vdash A \leftrightarrow \neg\neg A$ (law of double negation).
5. $\perp \rightarrow A$ (ex falso quodlibet)
6. $\Delta \vdash A$ implies $\Delta \vdash B$ iff $\Delta \vdash A \rightarrow B$.

The proofs of these results can be found <http://planetmath.org/SomeTheoremSchemasOfPropositionalLogic>.

Theorem 1. (*Substitution Theorem*) Suppose p_1, \dots, p_m are all the propositional variables, not necessarily distinct, that occur in order in A , and if $B_1, \dots, B_m, C_1, \dots, C_m$ are wff's such that $\vdash B_i \leftrightarrow C_i$, then

$$\vdash A[B_1/p_1, \dots, B_m/p_m] \leftrightarrow A[C_1/p_1, \dots, C_m/p_m]$$

where $A[X_1/p_1, \dots, X_m/p_m]$ is the wff obtained from A by replacing p_i by the wff X_i via simultaneous substitution.

Proof. We do induction on the number n of \rightarrow in wff A .

If $n = 0$, A is either a propositional variable, say p , or \perp , which respectively means that $A[B/p] \leftrightarrow A[C/p]$ is either $B \leftrightarrow C$ or $\perp \leftrightarrow \perp$. The former is the assumption and the latter is a theorem.

Suppose now A has $n + 1$ occurrences of \rightarrow . We may write A as $X \rightarrow Y$ uniquely by unique readability. Also, both X and Y have at most n occurrences of \rightarrow .

Let A_1 be $A[B_1/p_1, \dots, B_m/p_m]$ and A_2 be $A[C_1/p_1, \dots, C_m/p_m]$. Then A_1 is $X_1 \rightarrow Y_1$ and $X_2 \rightarrow Y_2$, where X_1 is $X[B_1/p_1, \dots, B_k/p_k]$, Y_1 is $Y[B_{k+1}/p_{k+1}, \dots, B_m/p_m]$, X_2 is $X[C_1/p_1, \dots, C_k/p_k]$, and Y_2 is $Y[C_{k+1}/p_{k+1}, \dots, C_m/p_m]$.

Then	
by induction	$\vdash X_1 \leftrightarrow X_2$ (1)
by 2 above	$\vdash X_1 \rightarrow X_2$ and $\vdash X_2 \rightarrow X_1$ (2)
by induction	$\vdash Y_1 \leftrightarrow Y_2$ (3)
by 2 above	$\vdash Y_1 \rightarrow Y_2$ and $\vdash Y_2 \rightarrow Y_1$ (4)
since A_1 is $X_1 \rightarrow Y_1$	$A_1 \vdash X_1 \rightarrow Y_1$ (5)
by applying 1 to $\vdash X_2 \rightarrow X_1$ and (5)	$A_1 \vdash X_2 \rightarrow Y_1$ (6)
by applying 1 to (6) and $\vdash Y_1 \rightarrow Y_2$	$A_1 \vdash X_2 \rightarrow Y_2$ (7)
by the deduction theorem	$\vdash A_1 \rightarrow A_2$ (8)
by a similar reasoning as above	$\vdash A_2 \rightarrow A_1$ (9)
by applying 2 to (8) and (9)	$\vdash A_1 \leftrightarrow A_2$ (10)
	□

As a corollary, we have

Corollary 1. *If $\vdash B \leftrightarrow C$, then $\vdash A[B/s(p)] \leftrightarrow A[C/s(p)]$, where p is a propositional variable that occurs in A , $s(p)$ is a set of positions of occurrences of p in A , and the wff $A[X/s(p)]$ is obtained by replacing all p that occur in the positions in $s(p)$ in A by wff X .*

Proof. For any propositional variable q not being replaced, use the corresponding theorem $\vdash q \leftrightarrow q$, and then apply the substitution theorem. □

Remark. What about $\vdash B[A/p] \leftrightarrow C[A/p]$, given $\vdash B \leftrightarrow C$? Here, $B[A/p]$ and $C[A/p]$ are wff's obtained by uniform substitution of p (all occurrences of p) in B and C respectively. Since $B[A/p] \leftrightarrow C[A/p]$ is just

$(B \leftrightarrow C)[A/p]$, an instance of the schema $B \leftrightarrow C$ by assumption, the result follows directly if we assume $B \leftrightarrow C$ is a theorem schema.

Using the substitution theorem, we can easily derive more theorem schemas, such as

7. $(A \rightarrow B) \leftrightarrow (\neg B \rightarrow \neg A)$ (Law of Contraposition)

8. $A \rightarrow (\neg B \rightarrow \neg(A \rightarrow B))$

9. $((A \rightarrow B) \rightarrow A) \rightarrow A$ (Peirce's Law)

Proof. 7. Since $(\neg B \rightarrow \neg A) \rightarrow (A \rightarrow B)$ is already a theorem schema, we only need to show $\vdash (A \rightarrow B) \rightarrow (\neg B \rightarrow \neg A)$. By law of double negation (4 above) and the substitution theorem, it is enough to show that $\vdash (\neg\neg A \rightarrow \neg\neg B) \rightarrow (\neg B \rightarrow \neg A)$. But this is just an instance of an axiom schema. Combining the two schemas, we get $\vdash (A \rightarrow B) \leftrightarrow (\neg B \rightarrow \neg A)$.

8. First, observe that $A, A \rightarrow B \vdash B$ by modus ponens. Since $\vdash B \leftrightarrow \neg\neg B$, we have $A, A \rightarrow B \vdash \neg\neg B$ by the substitution theorem. So $A, A \rightarrow B, \neg B \vdash \perp$ by the deduction theorem, and $A, \neg B \vdash (A \rightarrow B) \rightarrow \perp$ by the deduction theorem again. Apply the deduction two more times, we get $\vdash A \rightarrow (\neg B \rightarrow \neg(A \rightarrow B))$.

9. To show $\vdash ((A \rightarrow B) \rightarrow A) \rightarrow A$, it is enough to show $\vdash \neg A \rightarrow \neg((A \rightarrow B) \rightarrow A)$ by 7 and modus ponens, or $\neg A \vdash \neg((A \rightarrow B) \rightarrow A)$ by the deduction theorem. Now, since $\vdash X \wedge Y \leftrightarrow \neg(X \rightarrow \neg Y)$ (as they are the same thing, and because $C \leftrightarrow C$ is a theorem schema), by the law of double negation and the substitution theorem, $\vdash X \wedge \neg Y \leftrightarrow \neg(X \rightarrow Y)$, and we have $\vdash (A \rightarrow B) \wedge \neg A \leftrightarrow \neg((A \rightarrow B) \rightarrow A)$. So to show $\neg A \vdash \neg((A \rightarrow B) \rightarrow A)$, it is enough to show $\neg A \vdash (A \rightarrow B) \wedge \neg A$, which is enough to show that $\neg A \vdash A \rightarrow B$ and $\neg A \vdash \neg A$, according to a meta-theorem found <http://planetmath.org/SomeTheoremSchemasOfPropositionalLogic> here. To show $\neg A \vdash A \rightarrow B$, it is enough to show $\{\neg A, A\} \vdash B$, and $A, \neg A, \perp, \perp \rightarrow B, B$ is such a deduction. The second statement $\neg A \vdash \neg A$ is clear. \square

As an application, we prove the following useful meta-theorems of propositional logic:

Proposition 1. *There is a wff A such that $\Delta \vdash A$ and $\Delta \vdash \neg A$ iff $\Delta \vdash \perp$*

Proof. Assume the former. Let \mathcal{E}_1 be a deduction of A from Δ and \mathcal{E}_2 a deduction of $\neg A$ from Δ , then

$$\mathcal{E}_1, \mathcal{E}_2, \perp$$

is a deduction of \perp from Δ . Conversely, assume the later. Pick any wff A (if necessary, pick \perp). Then $\perp \rightarrow A$ by ex falso quodlibet. By modus ponens, we have $\Delta \vdash A$. Similarly, $\Delta \vdash \neg A$. \square

Proposition 2. *If $\Delta, A \vdash B$ and $\Delta, \neg A \vdash B$, then $\Delta \vdash B$*

Proof. By assumption, we have $\Delta \vdash A \rightarrow B$ and $\Delta \vdash \neg A \rightarrow B$. Using modus ponens and the theorem schema $(A \rightarrow B) \rightarrow (\neg B \rightarrow \neg A)$, we have $\Delta \vdash \neg B \rightarrow \neg A$, or

$$\Delta, \neg B \vdash \neg A.$$

Similarly, $\Delta \vdash \neg B \rightarrow \neg \neg A$. By the law of double negation and the substitution theorem, we have $\Delta \vdash \neg B \rightarrow A$, or

$$\Delta, \neg B \vdash A.$$

By the previous proposition, $\Delta, \neg B \vdash \perp$, or $\Delta \vdash \neg \neg B$. Applying the substitution theorem and the law of double negation, we have

$$\Delta \vdash B.$$

\square