



planetmath.org

Math for the people, by the people.

substitutability

Canonical name	Substitutability
Date of creation	2013-03-22 19:12:15
Last modified on	2013-03-22 19:12:15
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	24
Author	CWoo (3771)
Entry type	Definition
Classification	msc 03B10
Classification	msc 03B05
Synonym	substitutable for
Related topic	Subformula
Related topic	FreeAndBoundVariables
Defines	free for
Defines	free substitution instance

In any logical system, the way to obtain (well-formed) formulas from existing ones is by attaching logical connectives to existing ones. For example, in classical propositional logic, if φ and ψ are formulas, the following

$$\varphi \vee \psi, \quad \neg\varphi, \quad \psi \wedge \varphi$$

are formed by attaching logical connectives \vee , \neg , and \wedge appropriately to φ and ψ .

Another convenient device is substitution:

replace all occurrences of some symbol x in a formula φ by an expression ψ .

We denote the resulting expression by

$$\varphi[\psi/x].$$

For example, in classical propositional logic, if φ is $p \wedge (\neg r \vee p)$, then $\varphi[(p \vee q)/p]$ is

$$(p \vee q) \wedge (\neg r \vee (p \vee q))$$

On the other hand, the expressions

- $\varphi[\neg/p]$, which is $\neg \wedge (\neg r \vee \neg)$, and
- $\varphi[q/\wedge]$, which is $p \vee q(\neg r \vee q)$,

are not formulas. Thus, one must be careful when performing substitutions on formulas lest the resulting expressions are ill-formed. In other words, conditions must be placed on x and ψ in $\varphi[\psi/x]$ in order that $\varphi[\psi/x]$ is a (well-formed) formula. These conditions are called the *substitutability conditions*. In this entry, we will concentrate on substitutability conditions on predicate logic. Details on substitutions in propositional logic can be found in <http://planetmath.org/SubstitutionsInLogichere>.

Substitution in First-Order Logic

Substitution works pretty much the same way for first-order logic as in propositional logic. However, the substitutability conditions are more subtle. Take a look at the following example:

$$\exists x (x = 1)$$

If we replace x by 0, we end up with

$$\exists 0 (0 = 1),$$

which is non-sensical (not a wff). This is because x occurs in the formula as a bound variable. (one reason why we distinguish the variables occurring in first order formulas into two types: free and bound).

We now formalize the notion of substitution in first-order logic. There are two parts: substitution for terms, and substitution for formulas.

Definition. For any term t , any symbol x , and any expression s , define $t[s/x]$ inductively, as follows:

1. if t is an individual variable or a constant symbol, then $t[s/x]$ is s if t is x , and $t[s/x]$ is t otherwise;
2. if t is $f(t_1, \dots, t_n)$, where f is an n -ary function symbol, and each t_i is a term, then $t[s/x]$ is $f(t_1[s/x], \dots, t_n[s/x])$.

For example, if t is $x + y$, then $t[(x - y)/x]$ is $(x - y) + y$.

It is easy to see that s is a term and x an individual variable, then $t[s/x]$ is a term. In addition, by induction, one can easily show that if the formula $s_1 = s_2$ is true, so is the formula $t[s_1/x] = t[s_2/x]$.

Next, we define substitution for formulas. In light of the last example at the beginning of this section, we need to be a little careful.

Definition. Let φ be a formula, x a symbol, and s an expression. The expression $\varphi[s/x]$ is again define inductively:

1. if φ is $t_1 = t_2$, then $\varphi[s/x]$ is $t_1[s/x] = t_2[s/x]$;
2. if φ is $R(t_1, \dots, t_n)$, then $\varphi[s/x]$ is $R(t_1[s/x], \dots, t_n[s/x])$;
3. if φ is $\neg\psi$, then $\varphi[s/x]$ is $\neg(\psi[s/x])$;
4. if φ is $\psi \vee \sigma$, then $\varphi[s/x]$ is $\psi[s/x] \vee \sigma[s/x]$;
5. if φ is $\exists y\psi$, then $\varphi[s/x]$ is $\exists y(\psi[s/x])$ if $x \neq y$, and $\varphi[s/x]$ is φ otherwise.

Again, substitutions involving logical connectives \rightarrow , \wedge , and the universal quantifier \forall can be derived from the rules given above.

For example, if φ is $\exists x(x = y \vee y = z)$, then $\varphi[t/y]$ is $\exists x(x = t \vee t = z)$, whereas $\varphi[t/x]$ is just φ .

Given that φ is a formula, it is easy to see that if x is an individual variable, and s is a term, then $\varphi[s/x]$ is a formula.

In addition, it is easy to see that sentences are not affected by substitutions: if φ is a sentence, then $\varphi[s/x]$ is just φ . In other words, sentences can not be changed into formulas with free variables.

Conversely, can a formula with free variables be changed into a sentence by substitution? Certainly. For example, if φ is

$$\exists x(y < x),$$

then $\varphi[x/y]$ is

$$\exists x(x < x).$$

Although syntactically correct, this is undesirable in many situations, particularly when we are interested in the interpretations of these formulas. In the example above, we have changed $\exists x(y < x)$, which many very well be true in many interpretations, into $\exists x(x < x)$, something with a fixed meaning (and always false if $<$ is interpreted as the usual less than relation).

The problem with the situation described in the last paragraph arises because a free variable in t becomes bound in $\varphi[t/x]$. To eliminate this undesirable situation, we define the notion of “free for”:

Definition. Let x be an individual variable, t a term, and φ a formula. We define the relation t is free for, or substitutable for x in φ , inductively, as follows:

1. φ is an atomic formula;
2. φ is $\neg\psi$, and t is free for x in ψ ;
3. φ is $\psi \vee \sigma$, and t is free for x in ψ and in σ ;
4. φ is $\exists y\psi$, and either
 - $x \notin \text{FV}(\varphi)$ (x does not occur free in φ), or
 - y does not occur in t , and t is free for x in ψ .

In words, t is free for x in φ iff whenever z is a variable in t , no literal subformula of φ of the form $\exists z\psi$ contains an occurrence of x which is free in φ .

For example, $f(x, y)$ is free for x in the following formulas:

$$P(x, y), \quad P(x) \vee \neg Q(z), \quad \neg \exists x \neg R(x, y), \quad \text{and} \quad \neg S(y) \vee \exists y T(y, z),$$

but not in the following formulas:

$$\exists y P(x, y), \quad Q(x) \vee \exists z \exists y R(x, y), \quad \text{and} \quad S(y) \rightarrow \forall y (T(y, y) \wedge \neg Q(x)).$$

Given any formula φ , we again write $\varphi(x)$ to mean that variable x occurs in φ . A *substitution instance* of $\varphi(x)$ is just $\varphi[t/x]$, or $\varphi(t)$ for short. Furthermore, if t is free for x in φ , then $\varphi(t)$ is called a *free substitution instance* of $\varphi(x)$.

It is easy, by induction, to show that if terms t_1 and t_2 are free for x in φ , and that the formula $t_1 = t_2$ is true, the substitution instances $\varphi(t_1)$ and $\varphi(t_2)$ are logically equivalent, as intended.