



planetmath.org

Math for the people, by the people.

creating an infinite model

Canonical name	CreatingAnInfiniteModel
Date of creation	2013-03-22 12:44:29
Last modified on	2013-03-22 12:44:29
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	9
Author	CWoo (3771)
Entry type	Example
Classification	msc 03B10
Classification	msc 03C07
Related topic	CompactnessTheoremForFirstOrderLogic
Related topic	GettingModelsIModelsConstructedFromConstants

From the syntactic compactness theorem for first order logic, we get this nice (and useful) result:

Let  $\mathbf{T}$  be a theory of first-order logic. If  $\mathbf{T}$  has finite models of unboundedly large sizes, then  $\mathbf{T}$  also has an infinite model.

*Proof.* Define the propositions

$$\Phi_n \equiv \underline{\exists x_1 \cdots \exists x_n. (x_1 \neq x_2) \wedge \cdots \wedge (x_1 \neq x_n) \wedge (x_2 \neq x_3) \wedge \cdots \wedge (x_{n-1} \neq x_n)}$$

( $\Phi_n$  says “*there exist (at least)  $n$  different elements in the world*”). Note that

$$\cdots \vdash \Phi_n \vdash \cdots \vdash \Phi_2 \vdash \Phi_1.$$

Define a new theory

$$\mathbf{T}_\infty = \mathbf{T} \cup \{\Phi_1, \Phi_2, \dots\}.$$

For any *finite* subset  $\mathbf{T}' \subset \mathbf{T}_\infty$ , we claim that  $\mathbf{T}'$  is consistent: Indeed,  $\mathbf{T}'$  contains axioms of  $\mathbf{T}$ , along with finitely many of  $\{\Phi_n\}_{n \geq 1}$ . Let  $\Phi_m$  correspond to the largest index appearing in  $\mathbf{T}'$ . If  $\mathcal{M}_m \models \mathbf{T}$  is a model of  $\mathbf{T}$  with at least  $m$  elements (and by hypothesis, such a model exists), then  $\mathcal{M}_m \models \mathbf{T} \cup \{\Phi_m\} \vdash \mathbf{T}'$ .

So every finite subset of  $\mathbf{T}_\infty$  is consistent; by the compactness theorem for first-order logic,  $\mathbf{T}_\infty$  is consistent, and by Gödel’s completeness theorem for first-order logic it has a model  $\mathcal{M}$ . Then  $\mathcal{M} \models \mathbf{T}_\infty \vdash \mathbf{T}$ , so  $\mathcal{M}$  is a model of  $\mathbf{T}$  with infinitely many elements ( $\mathcal{M} \models \Phi_n$  for any  $n$ , so  $\mathcal{M}$  has at least  $\geq n$  elements for all  $n$ ).  $\square$