



models constructed from constants

Canonical name	ModelsConstructedFromConstants
Date of creation	2013-03-22 12:44:42
Last modified on	2013-03-22 12:44:42
Owner	ratboy (4018)
Last modified by	ratboy (4018)
Numerical id	17
Author	ratboy (4018)
Entry type	Definition
Classification	msc 03C07
Synonym	completeness theorem
Synonym	Gödel completeness theorem
Related topic	UpwardsSkolemLowenheimTheorem
Defines	set of witnesses

The definition of a structure and of the satisfaction relation is nice, but it raises the following question : how do we get models in the first place? The most basic construction for models of first-order theory is the construction that uses constants. Throughout this entry, L is a fixed first-order language.

Let C be a set of constant symbols of L , and T be a theory in L . Then we say C is a *set of witnesses* for T if and only if for every formula φ with at most one free variable x , we have $T \vdash \exists x(\varphi) \Rightarrow \varphi(c)$ for some $c \in C$.

Lemma. Let T is any consistent set of sentences of L , and C is a set of new symbols such that $|C| = |L|$. Let $L' = L \cup C$. Then there is a consistent set $T' \subseteq L'$ extending T and which has C as set of witnesses.

Lemma. If T is a consistent theory in L , and C is a set of witnesses for T in L , then T has a model whose elements are the constants in C .

Proof: Let Σ be the signature for L . If T is a consistent set of sentences of L , then there is a maximal consistent $T' \supseteq T$. Note that T' and T have the same sets of witnesses. As every model of T' is also a model of T , we may assume T is maximal consistent.

We let the universe of \mathfrak{M} be the set of equivalence classes C/\sim , where $a \sim b$ if and only if “ $a = b$ ” $\in T$. As T is maximal consistent, this is an equivalence relation. We interpret the non-logical symbols as follows :

1. $[a] =^{\mathfrak{M}} [b]$ if and only if $a \sim b$;
2. Constant symbols are interpreted in the obvious way, i.e. if $c \in \Sigma$ is a constant symbol, then $c^{\mathfrak{M}} = [c]$;
3. If $R \in \Sigma$ is an n -ary relation symbol, then $([a_1], \dots, [a_n]) \in R^{\mathfrak{M}}$ if and only if $R(a_1, \dots, a_n) \in T$;
4. If $F \in \Sigma$ is an n -ary function symbol, then $F^{\mathfrak{M}}([a_0], \dots, [a_n]) = [b]$ if and only if “ $F(a_1, \dots, a_n) = b$ ” $\in T$.

From the fact that T is maximal consistent, and \sim is an equivalence relation, we get that the operations are well-defined (it is not so simple, i'll write it out later). The proof that $\mathfrak{M} \models T$ is a straightforward induction on the complexity of the formulas of T . \diamond

Corollary. (The extended completeness theorem) A set T of formulas of L is consistent if and only if it has a model (regardless of whether or not L has witnesses for T).

Proof: First add a set C of new constants to L , and expand T to T' in such a way that C is a set of witnesses for T' . Then expand T' to a maximal

consistent set T'' . This set has a model \mathfrak{M} consisting of the constants in C , and \mathfrak{M} is also a model of T . \diamond

Corollary. (Compactness theorem) A set T of sentences of L has a model if and only if every finite subset of T has a model.

Proof: Replace “has a model” by “is consistent”, and apply the syntactic compactness theorem. \diamond

Corollary. (Gödel’s completeness theorem) Let T be a consistent set of formulas of L . Then A sentence φ is a theorem of T if and only if it is true in every model of T .

Proof: If φ is not a theorem of T , then $\neg\varphi$ is consistent with T , so $T \cup \{\neg\varphi\}$ has a model \mathfrak{M} , in which φ cannot be true. \diamond

Corollary. (Downward Löwenheim-Skolem theorem) If $T \subseteq L$ has a model, then it has a model of power at most $|L|$.

Proof: If T has a model, then it is consistent. The model constructed from constants has power at most $|L|$ (because we must add at most $|L|$ many new constants). \diamond

Most of the treatment found in this entry can be read in more details in Chang and Keisler’s book *Model Theory*.