



Math for the people, by the people.

König's lemma

Canonical name	KonigsLemma
Date of creation	2013-03-22 15:38:26
Last modified on	2013-03-22 15:38:26
Owner	mps (409)
Last modified by	mps (409)
Numerical id	6
Author	mps (409)
Entry type	Theorem
Classification	msc 05C05
Synonym	Koenig's lemma

Theorem (König's lemma). *Let T be a rooted directed tree. If each vertex has finite degree but there are arbitrarily long rooted paths in T , then T contains an infinite path.*

Proof. For each $n \geq 1$, let P_n be a rooted path in T of length n , and let c_n be the child of the root appearing in P_n . By assumption, the set $\{c_n \mid n \geq 1\}$ is finite. Since the set $\{P_n \mid n \geq 1\}$ is infinite, the pigeonhole principle implies that there is a child c of the root such that $c = c_n$ for infinitely many n .

Now let us look at the subtree T_c of T rooted at c . Each vertex has finite degree, and since there are paths P_n of arbitrarily long length in T passing through c , there are arbitrarily long paths in T_c rooted at c . Hence if T satisfies the hypothesis of the lemma, the root has a child c such that T_c also satisfies the hypothesis of the lemma. Hence we may inductively build up a path in T of infinite length, at each stage selecting a child so that the subtree rooted at that vertex still has arbitrarily long paths. \square

References

- [1] Kleene, Stephen., *Mathematical Logic*, New York: Wiley, 1967.