

planetmath.org

Math for the people, by the people.

projective basis

Canonical name ProjectiveBasis

Date of creation 2013-03-22 19:14:38 Last modified on 2013-03-22 19:14:38

Owner CWoo (3771) Last modified by CWoo (3771)

Numerical id 10

Author CWoo (3771)
Entry type Definition
Classification msc 05B35
Classification msc 06C10
Classification msc 51A05
Synonym independent

Defines span

Defines projective independence
Defines projectively independent

Defines basis

In the parent entry, we see how one may define dimension of a projective space inductively, from its subspaces starting with a point, then a line, and working its way up. Another way to define dimension start with defining dimensions of the empty set, a point, a line, and a plane to be -1, 0, 1, and 2, and then use the fact that any other projective space is isomorphic to the projective space P(V) associated with a vector space V, and then define the dimension to be the dimension of V, minus 1. In this entry, we introduce a more natural way of defining dimensions, via the concept of a basis.

Throughout the discussion, \mathbf{P} is a projective space (as in any model satisfying the axioms of projective geometry).

Given a subset S of \mathbf{P} , the span of S, written $\langle S \rangle$, is the smallest subspace of \mathbf{P} containing S. In other words, $\langle S \rangle$ is the intersection of all subspaces of \mathbf{P} containing S. Thus, if S is itself a subspace of \mathbf{P} , $\langle S \rangle = S$. We also say that S spans $\langle S \rangle$.

One may think of $\langle \cdot \rangle$ as an operation on the powerset of **P**. It is easy to verify that this operation is a closure operator. In addition, $\langle \cdot \rangle$ is *algebraic*, in the sense that any point in $\langle S \rangle$ is in the span of a finite subset of S. In other words,

$$\langle S \rangle = \{ P \mid P \in \langle F \rangle \text{ for some finite } F \subseteq S \}.$$

Another property of $\langle \cdot \rangle$ is the exchange property: for any subspace U, if $P \notin U$, then for any point Q, $\langle U \cup \{P\} \rangle = \langle U \cup \{Q\} \rangle$ iff $Q \in \langle U \cup \{P\} \rangle - U$.

A subset S of \mathbf{P} is said to be *projectively independent*, or simply *independent*, if, for any proper subset S' of S, the span of S' is a proper subset of the span of S: $\langle S' \rangle \subset \langle S \rangle$. This is the same as saying that S is a *minimal* spanning set for $\langle S \rangle$, in the sense that no proper subset of S spans $\langle S \rangle$. Equivalently, S is independent iff for any $x \in S$, $\langle S - \{x\} \rangle \neq \langle S \rangle$.

S is called a *projective basis*, or simply *basis* for \mathbf{P} , if S is independent and spans \mathbf{P} .

All of the properties about spanning sets, independent sets, and bases for vector spaces have their projective counterparts. We list some of them here:

- 1. Every projective space has a basis.
- 2. If S_1, S_2 are independent, then $\langle S_1 \cap S_2 \rangle = \langle S_1 \rangle \cap \langle S_2 \rangle$.
- 3. If S is independent and $P \in \langle S \rangle$, then there is $Q \in S$ such that $(\{P\} \cup S) \{Q\} \text{ spans } \langle S \rangle$.

- 4. Let B be a basis for **P**. If S spans **P**, then $|B| \leq |S|$. If S is independent, then $|S| \leq |B|$. As a result, all bases for **P** have the same cardinality.
- 5. Every independent subset in P may be extended to a basis for P.
- 6. Every spanning set for **P** may be reduced to a basis for **P**.

In light of items 1 and 4 above, we may define the *dimension* of \mathbf{P} to be the cardinality of its basis.

One of the main result on dimension is the dimension formula: if U, V are subspaces of \mathbf{P} , then

$$\dim(U) + \dim(V) = \dim(U \cup V) + \dim(U \cap V),$$

which is the counterpart of the same formula for vector subspaces of a vector space (see http://planetmath.org/DimensionFormulaeForVectorSpacesthis entry).

References

[1] A. Beutelspacher, U. Rosenbaum *Projective Geometry, From Foundations to Applications*, Cambridge University Press (2000)