

divisibility of central binomial coefficient

 ${\bf Canonical\ name} \quad {\bf Divisibility Of Central Binomial Coefficient}$

Date of creation 2013-03-22 17:41:22 Last modified on 2013-03-22 17:41:22

Owner rspuzio (6075) Last modified by rspuzio (6075)

Numerical id 8

Author rspuzio (6075)

Entry type Proof
Classification msc 05A10
Classification msc 11B65

In this entry, we shall prove two results about the divisibility of central binomial coefficients which were stated in the main entry.

Theorem 1. If $n \geq 3$ is an integer and p is a prime number such that n , then <math>p divides $\binom{2n}{n}$.

Proof. We will examine the following expression for our binomial coefficient:

$$\binom{2n}{n} = \frac{2n(2n-1)\cdots(n+2)(n+1)}{n(n-1)\cdots3\cdot2\cdot1}.$$

Since n , we find <math>p appearing in the numerator. However, p cannot appear in the denominator because the terms there are all smaller than n. Hence, p cannot be cancelled, so it must divide $\binom{2n}{n}$.

Theorem 2. If $n \geq 3$ is an integer and p is a prime number such that 2n/3 , then <math>p does not divide $\binom{2n}{n}$.

Proof. We will again examine our expression for our binomial coefficient:

$$\binom{2n}{n} = \frac{2n(2n-1)\cdots(n+2)(n+1)}{n(n-1)\cdots3\cdot2\cdot1}.$$

This time, because 2n/3 , we find <math>p appearing in the denominator and 2p appearing in the numerator. No other multiples will appear because, if m > 2, then mp > 2n. The two occurrences of p noted above cancel, hence p is not a prime factor of $\binom{2n}{n}$.