



Math for the people, by the people.

## Erdős-Rado theorem

|                  |                     |
|------------------|---------------------|
| Canonical name   | ErdHosRadoTheorem   |
| Date of creation | 2013-03-22 12:59:30 |
| Last modified on | 2013-03-22 12:59:30 |
| Owner            | Henry (455)         |
| Last modified by | Henry (455)         |
| Numerical id     | 9                   |
| Author           | Henry (455)         |
| Entry type       | Theorem             |
| Classification   | msc 05D10           |
| Classification   | msc 03E05           |
| Related topic    | arrowsrelation      |
| Related topic    | ArrowsRelation      |

Repeated exponentiation for cardinals is denoted  $\exp_i(\kappa)$ , where  $i < \omega$ . It is defined by:

$$\exp_0(\kappa) = \kappa$$

and

$$\exp_{i+1}(\kappa) = 2^{\exp_i(\kappa)}$$

The Erdős-Rado theorem states that:

$$\exp_i(\kappa)^+ \rightarrow (\kappa^+)_\kappa^{i+1}$$

That is, if  $f : [\exp_i(\kappa)^+]^{i+1} \rightarrow \kappa$  then there is a homogeneous set of size  $\kappa^+$ .

As special cases,  $(2^\kappa)^+ \rightarrow (\kappa^+)_\kappa^2$  and  $(2^{\aleph_0})^+ \rightarrow (\aleph_1)_{\aleph_0}^2$ .