



Math for the people, by the people.

Vandermonde identity

Canonical name	VandermondeIdentity
Date of creation	2013-03-22 14:08:49
Last modified on	2013-03-22 14:08:49
Owner	mps (409)
Last modified by	mps (409)
Numerical id	8
Author	mps (409)
Entry type	Theorem
Classification	msc 05A19
Related topic	PascalsRule

Theorem 1 ([?] 24.1.1 formula A. II.). *For any p and q and any k with $0 \leq k \leq p + q$,*

$$\binom{p+q}{k} = \sum_{i=0}^k \binom{p}{i} \binom{q}{k-i}. \quad (*)$$

Proof. Let P and Q be disjoint sets with $|P| = p$ and $|Q| = q$. Then the left-hand side of Equation (*) is equal to the number of subsets of $P \cup Q$ of size k . To build a subset of $P \cup Q$ of size k , we first decide how many elements, say i with $0 \leq i \leq k$, we will select from P . We can then select those elements in $\binom{p}{i}$ ways. Once we have done so, we must select the remaining $k - i$ elements from Q , which we can do in $\binom{q}{k-i}$ ways. Thus there are $\binom{p}{i} \binom{q}{k-i}$ ways to select a subset of $P \cup Q$ of size k subject to the restriction that exactly i elements come from P . Summing over all possible i completes the proof. \square

References

- [1] Abramowitz, M., and I. A. Stegun, eds. *Handbook of Mathematical Functions*. National Bureau of Standards, Dover, New York, 1974.