



planetmath.org

Math for the people, by the people.

reduction algorithm for symmetric  
polynomials

Canonical name	ReductionAlgorithmForSymmetricPolynomials
Date of creation	2013-03-22 12:11:17
Last modified on	2013-03-22 12:11:17
Owner	djao (24)
Last modified by	djao (24)
Numerical id	6
Author	djao (24)
Entry type	Proof
Classification	msc 05E05
Classification	msc 12F10
Defines	height

We give here an algorithm for reducing a symmetric polynomial into a polynomial in the elementary symmetric polynomials.

We define the *height* of a monomial  $x_1^{e_1} \cdots x_n^{e_n}$  in  $R[x_1, \dots, x_n]$  to be  $e_1 + 2e_2 + \cdots + ne_n$ . The height of a polynomial is defined to be the maximum height of any of its monomial terms, or 0 if it is the zero polynomial.

Let  $f$  be a symmetric polynomial. We reduce  $f$  into elementary symmetric polynomials by induction on the height of  $f$ . Let  $cx_1^{e_1} \cdots x_n^{e_n}$  be the monomial term of maximal height in  $f$ . Consider the polynomial

$$g := f - cs_1^{e_n - e_{n-1}} s_2^{e_{n-1} - e_{n-2}} \cdots s_{n-1}^{e_2 - e_1} s_n^{e_1}$$

where  $s_k$  is the  $k$ -th elementary symmetric polynomial in the  $n$  variables  $x_1, \dots, x_n$ . Then  $g$  is a symmetric polynomial of lower height than  $f$ , so by the induction hypothesis,  $g$  is a polynomial in  $s_1, \dots, s_n$ , and it follows immediately that  $f$  is also a polynomial in  $s_1, \dots, s_n$ .