



planetmath.org

Math for the people, by the people.

order of six means

Canonical name	OrderOfSixMeans
Date of creation	2013-03-22 18:45:28
Last modified on	2013-03-22 18:45:28
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	4
Author	pahio (2872)
Entry type	Theorem
Classification	msc 06A05
Classification	msc 26B35
Classification	msc 26D07
Related topic	Mean3
Related topic	ComparisonOfPythagoreanMeans
Related topic	InequalityWithAbsoluteValues
Related topic	LehmerMean

The of the six usual means of two positive numbers (a and b) is from the least to the greatest one

1. harmonic mean,
 2. geometric mean,
 3. Heronian mean,
 4. arithmetic mean,
 5. quadratic mean,
 6. contraharmonic mean,
- i. e.

$$\frac{2ab}{a+b} \leq \sqrt{ab} \leq \frac{a+\sqrt{ab}+b}{3} \leq \frac{a+b}{2} \leq \sqrt{\frac{a^2+b^2}{2}} \leq \frac{a^2+b^2}{a+b}.$$

The equality signs are valid iff $a = b$.

Proof. If $x^2 - y^2 \geq 0$ for nonnegative x and y , then $x \geq y$.

“1 \leq 2”:

$$\left(\sqrt{ab}\right)^2 - \left(\frac{a+b}{2}\right)^2 = ab - \frac{4a^2b^2}{(a+b)^2} = ab \left(1 - \frac{4ab}{(a+b)^2}\right) = ab \cdot \frac{(a+b)^2 - 4ab}{(a+b)^2} = \frac{ab(a-b)^2}{(a+b)^2} \geq 0$$

“2 \leq 3” and “3 \leq 4”: proven in Heronian mean is between geometric and arithmetic mean

“4 \leq 5”:

$$\left(\sqrt{\frac{a^2+b^2}{2}}\right)^2 - \left(\frac{a+b}{2}\right)^2 = \frac{2a^2+2b^2-a^2-2ab-b^2}{4} = \left(\frac{a-b}{2}\right)^2 \geq 0$$

“5 \leq 6”:

$$\begin{aligned} \left(\frac{a^2+b^2}{a+b}\right)^2 - \left(\sqrt{\frac{a^2+b^2}{2}}\right)^2 &= \frac{2(a^2+b^2)^2 - (a^2+b^2)(a+b)^2}{2(a+b)^2} = \frac{(a^2+b^2)(2a^2+2b^2-a^2-2ab-b^2)}{2(a+b)^2} \\ &= \frac{(a^2+b^2)(a-b)^2}{2(a+b)^2} \geq 0 \end{aligned}$$