

## planetmath.org

Math for the people, by the people.

## basic facts about ordered rings

 ${\bf Canonical\ name} \quad {\bf BasicFactsAboutOrderedRings}$ 

Date of creation 2013-03-22 16:17:21 Last modified on 2013-03-22 16:17:21 Owner Wkbj79 (1863) Last modified by Wkbj79 (1863)

Numerical id 11

Author Wkbj79 (1863)

Entry type Result
Classification msc 06F25
Classification msc 12J15
Classification msc 13J25

Related topic MathbbCIsNotAnOrderedField

Throughout this entry,  $(R, \leq)$  is an ordered ring.

**Lemma 1.** If  $a, b, c \in R$  with a < b, then a + c < b + c.

*Proof.* The contrapositive will be proven.

Let  $a, b, c \in R$  with  $a + c \ge b + c$ . Note that  $-c \in R$ . Thus,

$$b = b + 0$$

$$= b + c + (-c)$$

$$\leq a + c + (-c)$$

$$= a + 0$$

$$= a. \square$$

**Lemma 2.** If  $|R| \neq 1$  and R has a characteristic, then it must be 0.

*Proof.* Suppose not. Let n be a positive integer such that char R = n. Since  $|R| \neq 1$ , it must be the case that n > 1.

Let  $r \in R$  with r > 0. By the previous lemma,  $0 < r \le ... \le \sum_{i=1}^{n-1} r \le ...$ 

$$\sum_{j=1}^{n} r = 0, \text{ a contradiction.}$$

**Lemma 3.** If  $a, b \in R$  with  $a \le b$  and  $c \in R$  with c < 0, then  $ac \ge bc$ .

*Proof.* Note that  $-c \in R$  and 0 = c + (-c) < 0 + (-c) = -c. Since  $a \le b$ ,  $-(ac) = a(-c) \le b(-c) = -(bc)$ . Thus,

$$bc = bc + 0$$

$$= bc + (ac + (-(ac)))$$

$$= (bc + ac) + (-(ac))$$

$$\leq (bc + ac) + (-(bc))$$

$$= -(bc) + (bc + ac)$$

$$= (-(bc) + bc) + ac$$

$$= 0 + ac$$

$$= ac. \square$$

**Lemma 4.** Suppose further that R is a ring with multiplicative identity  $1 \neq 0$ . Then 0 < 1.

*Proof.* Suppose that  $0 \not< 1$ . Since R is an ordered ring, it must be the case that 1 < 0. By the previous lemma,  $1 \cdot 1 \ge 0 \cdot 1$ . Thus,  $1 \ge 0$ , a contradiction.