



**planetmath.org**

Math for the people, by the people.

# Birkhoff Recurrence Theorem

Canonical name	BirkhoffRecurrenceTheorem
Date of creation	2015-03-20 0:56:48
Last modified on	2015-03-20 0:56:48
Owner	Filipe (28191)
Last modified by	Filipe (28191)
Numerical id	2
Author	Filipe (28191)
Entry type	Theorem
Related topic	Poincar Recurrence Theorem

Let  $T : X \rightarrow X$  be a continuous transformation in a compact metric space  $X$ . Then, there exists some point  $x \in X$  that is recurrent to  $T$ , that is, there exists a sequence  $(n_k)_k$  such that  $T^{n_k}(x) \rightarrow x$  when  $k \rightarrow \infty$ .

Several proofs of this theorem are available. It may be obtained from topological arguments together with Zorn's lemma. It is also a consequence of Krylov-Bogolyubov theorem, or existence of invariant probability measures theorem, which asserts that every continuous transformation in a compact metric space admits an invariant probability measure, and an application of Poincaré Recurrence theorem to that invariant probability measure yields Birkhoff Recurrence theorem.

There is also a generalization of Birkhoff recurrence theorem for multiple commuting transformations, known as Birkhoff Multiple Recurrence theorem.