



Math for the people, by the people.

Birkhoff prime ideal theorem

Canonical name	BirkhoffPrimeIdealTheorem
Date of creation	2013-03-22 17:02:18
Last modified on	2013-03-22 17:02:18
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	11
Author	CWoo (3771)
Entry type	Theorem
Classification	msc 06D05
Classification	msc 03E25
Related topic	DistributiveLattice

Birkhoff Prime Ideal Theorem. Let L be a distributive lattice and I a proper lattice ideal of L . Pick any element $a \notin I$. Then there is a prime ideal P in L such that $I \subseteq P$ and $a \notin P$.

Proof. If I is prime, then we are done. Let $S := \{J \mid J \text{ is an ideal in } L, \text{ and } a \notin J\}$. Then $I \in S$. Order S by inclusion. This turns S into a poset. Let C be a chain in S . Let $K = \bigcup C$. If $x, y \in K$, then $x \in J_1$ and $y \in J_2$ for some ideals $J_1, J_2 \in C$. Since C is a chain, we may assume that $J_1 \subseteq J_2$, so that $x \in J_2$ as well. This means $x \vee y \in J_2 \subseteq K$. Next, assume $x \in K$ and $y \leq x$. Then $x \in J$ for some ideal $J \in C$, so that $y \in J \subseteq K$ also. This shows that K is an ideal. If $a \in K$, then $a \in J$ for some $J \in C \subseteq S$, contradicting the definition of S . So $a \notin K$ and $K \in S$ also. This shows that every chain in S has an upper bound. We can now appeal to Zorn's lemma, and conclude that S has a maximal element, say P .

We now want to show that P is the candidate that we are seeking: P is a prime ideal in L and $a \notin P$. Since $P \in S$, P is an ideal such that $a \notin P$. So the only thing left to prove is that P is prime. This amounts to showing that if $x \wedge y \in P$, then $x \in P$ or $y \in P$. Suppose not: $x, y \notin P$. Let Q_1 be the ideal generated by elements of P and x , and Q_2 the ideal generated by P and y . Since Q_1 and Q_2 properly contain P , $a \in Q_1$ and $a \in Q_2$. Write $a \leq p_1 \vee x$ and $a \leq p_2 \vee y$, where $p_1, p_2 \in P$. Then $a \vee p_2 \leq (p_1 \vee p_2) \vee x$ and $a \vee p_1 \leq (p_1 \vee p_2) \vee y$. Take the meet of these two expressions, and we obtain $(a \vee p_2) \wedge (a \vee p_1) \leq ((p_1 \vee p_2) \vee x) \wedge ((p_1 \vee p_2) \vee y)$. Since L is distributive, on the left hand side, we get $a \vee (p_1 \wedge p_2)$. On the right hand side, we have $(p_1 \vee p_2) \vee (x \wedge y) \in P$. As the left hand side is less than or equal to the right hand side, we get that $a \vee (p_1 \wedge p_2) \in P$. Since $a \leq a \vee (p_1 \wedge p_2) \in P$, $a \in P$, a contradiction. Therefore, P is prime and the proof is complete. \square

In the proof, we use the fact that, an element $a \in L$ belongs to the ideal generated by ideals I_k iff a is less than or equal to a finite join of elements, each of which belongs to some I_k .

Remarks.

1. The theorem can be generalized: if we use a subset $S \cap I = \emptyset$ instead of an element $a \notin I$, there is a prime ideal P containing I but excluding S .
2. Birkhoff's prime ideal theorem has been shown to be equivalent to the axiom of choice, under ZF.