

## planetmath.org

Math for the people, by the people.

## meet continuous

Canonical name MeetContinuous
Date of creation 2013-03-22 16:36:41
Last modified on 2013-03-22 16:36:41

Owner CWoo (3771) Last modified by CWoo (3771)

Numerical id 12

Author CWoo (3771)
Entry type Definition
Classification msc 06A12
Classification msc 06B35

Synonym order convergence

Related topic CriteriaForAPosetToBeACompleteLattice

Related topic JoinInfiniteDistributive

Defines join continuous Defines order converges Let L be a meet semilattice. We say that L is meet continuous if

- 1. for any monotone net  $D = \{x_i \mid i \in I\}$  in L, its supremum  $\bigvee D$  exists, and
- 2. for any  $a \in L$  and any monotone net  $\{x_i \mid i \in I\}$ ,

$$a \wedge \bigvee \{x_i \mid i \in I\} = \bigvee \{a \wedge x_i \mid i \in I\}.$$

A monotone net  $\{x_i \mid i \in I\}$  is a net  $x: I \to L$  such that x is a non-decreasing function; that is, for any  $i \leq j$  in I,  $x_i \leq x_j$  in L.

Note that we could have replaced the first condition by saying simply that  $D \subseteq L$  is a directed set. (A monotone net is a directed set, and a directed set is a trivially a monotone net, by considering the identity function as the net). It's not hard to see that if D is a directed subset of L, then  $a \wedge D := \{a \wedge x \mid x \in D\}$  is also directed, so that the right hand side of the second condition makes sense.

Dually, a join semilattice L is join continuous if its dual (as a meet semilattice) is meet continuous. In other words, for any antitone net  $D = \{x_i \mid i \in I\}$ , its infimum  $\bigwedge D$  exists and that

$$a \vee \bigwedge \{x_i \mid i \in I\} = \bigwedge \{a \vee x_i \mid i \in I\}.$$

An antitone net is just a net  $x: I \to L$  such that for  $i \leq j$  in  $I, x_j \leq x_i$  in L. Remarks.

- A meet continuous lattice is a complete lattice, since a poset such that finite joins and directed joins exist is a complete lattice (see the link below for a proof of this).
- Let a lattice L be both meet continuous and join continuous. Let  $\{x_i \mid i \in I\}$  be any net in L. We define the following:

$$\overline{\lim} x_i = \bigwedge_{j \in I} \{ \bigvee_{j \le i} x_i \}$$
 and  $\underline{\lim} x_i = \bigvee_{j \in I} \{ \bigwedge_{i \le j} x_i \}$ 

If there is an  $x \in L$  such that  $\overline{\lim} x_i = x = \underline{\lim} x_i$ , then we say that the net  $\{x_i\}$  order converges to x, and we write  $x_i \to x$ , or  $x = \lim x_i$ . Now, define a subset  $C \subseteq L$  to be closed (in L) if for any net  $\{x_i\}$  in C such that  $x_i \to x$  implies that  $x \in C$ , and open if its set complement is closed, then L becomes a topological lattice. With respect to this topology, meet  $\wedge$  and join  $\vee$  are easily seen to be continuous.

## References

- [1] G. Birkhoff, *Lattice Theory*, 3rd Edition, Volume 25, AMS, Providence (1967).
- [2] G. Gierz, K. H. Hofmann, K. Keimel, J. D. Lawson, M. W. Mislove, D. S. Scott, *Continuous Lattices and Domains*, Cambridge University Press, Cambridge (2003).
- [3] G. Grätzer, General Lattice Theory, 2nd Edition, Birkhäuser (1998).