

Let R be a commutative ring. A *Newtonian coalgebra* over R is an R -module C which is simultaneously a coalgebra with comultiplication $\Delta: C \rightarrow C \otimes C$ and an algebra with multiplication $\cdot: C \otimes C \rightarrow C$ such that Δ is a derivation over \cdot , that is, such that the identity

$$\Delta(u \cdot v) = \Delta(u) \cdot v + u \cdot \Delta(v)$$

holds for any u and v in C . Newtonian coalgebras were introduced by Joni and Rota in [?], where they were called *infinitesimal coalgebras*. They reserved the term “Newtonian coalgebra” for the special case of the coalgebra of divided differences. This example was studied in more detail by Hirschhorn and Raphael [?]. Joni and Rota also showed that Newtonian coalgebras provide a language which can explain iterated differentiation of trigonometric functions as well as Faà di Bruno’s formula. See also the paper of Nichols and Sweedler [?] for more on trigonometric coalgebras.

A Newtonian coalgebra cannot have both a unit and a counit, so no Newtonian coalgebra is a Hopf algebra. However, Aguiar [?] developed a notion of antipode that makes sense for Newtonian coalgebras, leading to what he calls an infinitesimal Hopf algebra. Ehrenborg and Readdy [?] used Newtonian coalgebras to give an algebraic structure to the <http://planetmath.org/CdIndexcd-index>, a poset invariant generalizing the f -vector of polytopes.

One example of a Newtonian coalgebra is the free associative algebra $R\langle \mathbf{a}, \mathbf{b} \rangle$ of polynomials on the noncommuting variables \mathbf{a} and \mathbf{b} with coefficients in R . The product is the ordinary noncommutative polynomial product, and the comultiplication is defined by setting

$$\Delta(u_1 \cdots u_n) = \sum_{j \in [n]} u_1 \cdots u_{j-1} \otimes u_{j+1} \cdots u_n$$

for each monomial and extending by linearity.

References

- [1] M. Aguiar, Infinitesimal Hopf algebras. *New trends in Hopf algebra theory: (La Falda, 1999)*, 1–29, Contemp. Math., 267, Amer. Math. Soc., Providence, RI, 2000.
- [2] M. Aguiar, Infinitesimal Hopf algebras and the **cd**-index of polytopes. *Discrete Comput. Geom.*, 27 (2002), no. 1, 3–28.

- [3] R. Ehrenborg and M. Readdy, Coproducts and the **cd**-index, *J. Algebr. Comb.*, 8 (1998), 273–299.
- [4] P. S. Hirschhorn and L. A. Raphael, Coalgebraic foundation of the method of divided differences, *Adv. Math.*, 91 (1992), 75–135.
- [5] S. A. Joni and G.-C. Rota, *Coalgebras and bialgebras in combinatorics*, Stud. Appl. Math., 61 (1979), pp. 93–139.
- [6] W. Nichols and M. Sweedler, *Hopf algebras and combinatorics*, in *Proceedings of the conference on umbral calculus and Hopf algebras*, ed. R. Morris, AMS, 1982.