



sectionally complemented lattice

Canonical name	SectionallyComplementedLattice
Date of creation	2013-03-22 17:58:46
Last modified on	2013-03-22 17:58:46
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	6
Author	CWoo (3771)
Entry type	Definition
Classification	msc 06C15
Classification	msc 06B05
Related topic	DifferenceOfLatticeElements
Defines	sectionally complemented
Defines	dually sectionally complemented lattice

Proposition 1. *Let L be a lattice with the least element 0 . Then the following are equivalent:*

1. *Every pair of elements have a difference.*
2. *for any $a \in L$, the lattice interval $[0, a]$ is a complemented lattice.*

Proof. Suppose first that every pair of elements have a difference. Let $b \in [0, a]$ and let c be a difference between a and b . So $0 = b \wedge c$ and $c \vee b = b \vee a = a$, since $b \leq a$. This shows that c is a complement of b in $[0, a]$.

Next suppose that $[0, a]$ is complemented for every $a \in L$. Let $x, y \in L$ be any two elements in L . Let $a = x \vee y$. Since $[0, a]$ is complemented, y has a complement, say $z \in [0, a]$. This means that $y \wedge z = 0$ and $y \vee z = a = x \vee y$. Therefore, z is a difference of x and y . \square

Definition. A lattice L with the least element 0 satisfying either of the two equivalent conditions above is called a *sectionally complemented lattice*.

Every relatively complemented lattice is sectionally complemented. Every sectionally complemented distributive lattice is relatively complemented.

Dually, one defines a *dually sectionally complemented lattice* to be a lattice L with the top element 1 such that for every $a \in L$, the interval $[a, 1]$ is complemented, or, equivalently, the lattice dual L^∂ is sectionally complemented.

References

- [1] G. Grätzer, *General Lattice Theory*, 2nd Edition, Birkhäuser (1998)