



planetmath.org

Math for the people, by the people.

proof of Schroeder-Bernstein theorem using
Tarski-Knaster theorem

Canonical name	ProofOfSchroederBernsteinTheoremUsingTarskiKnasterTheorem
Date of creation	2013-03-22 15:30:24
Last modified on	2013-03-22 15:30:24
Owner	kompik (10588)
Last modified by	kompik (10588)
Numerical id	7
Author	kompik (10588)
Entry type	Proof
Classification	msc 06B99
Related topic	SchroederBernsteintheorem
Related topic	TarskiKnastertheorem
Related topic	TarskiKnasterTheorem
Related topic	SchroederBernsteinTheorem

The Tarski-Knaster theorem can be used to give a short, elegant proof of the Schroeder-Bernstein theorem.

Proof. Suppose $f : S \rightarrow T$ and $g : T \rightarrow S$ are injective. Define a function $\varphi : P(S) \rightarrow P(S)$ by $\varphi(X) = S \setminus g(T \setminus f(X))$.

If $X \subseteq Y \subseteq S$, then $S \setminus g(T \setminus f(X)) \subseteq S \setminus g(T \setminus f(Y))$, and so φ is monotone. Since $P(S)$ is a complete lattice, we may apply the Tarski-Knaster theorem to conclude that the set of fixed points of φ is a complete lattice and thus nonempty.

Let C be a fixed point of φ . We have

$$S \setminus C = g(T \setminus f(C)).$$

Hence $g|_{T \setminus f(C)} : T \setminus f(C) \rightarrow S \setminus C$ and $f|_C : C \rightarrow f(C)$ are bijections. We can therefore construct the desired bijection $h : S \rightarrow T$ by defining

$$h(x) = \begin{cases} f(x) & \text{if } x \in C \\ (g|_{T \setminus f(C)})^{-1}(x) & \text{if } x \notin C. \quad \square \end{cases}$$

The usual proof of Schroeder-Bernstein theorem explicitly constructs a fixed point of φ .

References

- [1] Thomas Forster, *Logic, induction and sets*, Cambridge University Press, Cambridge, 2003.
- [2] M. Kolibiar, A. Legéň, T. Šalát, and Š. Znáťm, *Algebra a príbuzné disciplíny*, Alfa, Bratislava, 1992 (Slovak).