

planetmath.org

Math for the people, by the people.

atom

Canonical name Atom

Date of creation 2013-03-22 15:20:09 Last modified on 2013-03-22 15:20:09

Owner CWoo (3771) Last modified by CWoo (3771)

Numerical id 13

Author CWoo (3771)
Entry type Definition
Classification msc 06A06
Classification msc 06B99
Defines atomic poset
Defines atomic lattice
Defines atomistic lattice

Defines atomistic

Let P be a poset, partially ordered by \leq . An element $a \in P$ is called an atom if it covers some minimal element of P. As a result, an atom is never minimal. A poset P is called atomic if for every element $p \in P$ that is not minimal has an atom a such that $a \leq p$.

Examples.

- 1. Let A be a set and $P = 2^A$ its power set. P is a poset ordered by \subseteq with a unique minimal element \varnothing . Thus, all singleton subsets $\{a\}$ of A are atoms in P.
- 2. \mathbb{Z}^+ is partially ordered if we define $a \leq b$ to mean that $a \mid b$. Then 1 is a minimal element and any prime number p is an atom.

Remark. Given a lattice L with underlying poset P, an element $a \in L$ is called an atom (of L) if it is an atom in P. A lattice is a called an atomic lattice if its underlying poset is atomic. An atomistic lattice is an atomic lattice such that each element that is not minimal is a join of atoms. If a is an atom in a semimodular lattice L, and if a is not under x, then $a \vee x$ is an atom in any interval lattice I where $x = \bigwedge I$.

Examples.

- 1. $P = 2^A$, with the usual intersection and union as the lattice operations meet and join, is atomistic: every subset B of A is the union of all the singleton subsets of B.
- 2. \mathbb{Z}^+ , partially ordered as above, with lattice binary operations defined by $a \wedge b = \gcd(a,b)$, and $a \vee b = \operatorname{lcm}(a,b)$, is a lattice that is atomic, as we have seen earlier. But it is not atomistic: 4 is not a join of 2's; 36 is not a join of 2 and 3 are just two counterexamples.