



planetmath.org

Math for the people, by the people.

supremum over closure

Canonical name	SupremumOverClosure
Date of creation	2013-03-22 17:08:22
Last modified on	2013-03-22 17:08:22
Owner	Wkbj79 (1863)
Last modified by	Wkbj79 (1863)
Numerical id	9
Author	Wkbj79 (1863)
Entry type	Theorem
Classification	msc 06A05
Classification	msc 26A15

Theorem 1. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function and $A \subseteq \mathbb{R}$. Then $\sup_{x \in A} f(x) = \sup_{x \in \overline{A}} f(x)$, where \overline{A} denotes the closure of A .

Proof. The theorem is clearly true for $A = \emptyset$. Thus, it will be assumed that $A \neq \emptyset$.

Since $A \subseteq \overline{A}$, we have $\sup_{x \in A} f(x) \leq \sup_{x \in \overline{A}} f(x)$.

Suppose first that $\sup_{x \in \overline{A}} f(x) = \infty$. Let $r \in \mathbb{R}$. Then there exists $x_0 \in \overline{A}$ with $f(x_0) \geq r + 1$. Since f is continuous, there exists $\delta > 0$ such that, for any $x \in \mathbb{R}$ with $-\delta < x - x_0 < \delta$, we have $-1 < f(x) - f(x_0) < 1$. Since $x_0 \in \overline{A}$, there exists $x_1 \in A$ with $-\delta < x_1 - x_0 < \delta$. (Recall that $x \in \overline{A}$ if and only if every neighborhood of x intersects A .) Thus, $f(x_1) - f(x_0) > -1$. Therefore, $f(x_1) > f(x_0) - 1 \geq r + 1 - 1 = r$. Hence, $\sup_{x \in A} f(x) = \infty$.

Now suppose that $\sup_{x \in \overline{A}} f(x) = R$ for some $R \in \mathbb{R}$. Let $\varepsilon > 0$. Then there exists $x_2 \in \overline{A}$ with $f(x_2) \geq R - \frac{\varepsilon}{2}$. Since f is continuous, there exists $\delta' > 0$ such that, for any $x \in \mathbb{R}$ with $-\delta' < x - x_2 < \delta'$, we have $-\frac{\varepsilon}{2} < f(x) - f(x_2) < \frac{\varepsilon}{2}$. Since $x_2 \in \overline{A}$, there exists $x_3 \in A$ with $-\delta' < x_3 - x_2 < \delta'$. Thus, $f(x_3) - f(x_2) > -\frac{\varepsilon}{2}$. Therefore, $f(x_3) > f(x_2) - \frac{\varepsilon}{2} \geq R - \frac{\varepsilon}{2} - \frac{\varepsilon}{2} = R - \varepsilon$. Hence, $\sup_{x \in A} f(x) \geq R$.

In either case, it follows that $\sup_{x \in A} f(x) = \sup_{x \in \overline{A}} f(x)$. □

Note that this theorem also holds for continuous functions $f: X \rightarrow \mathbb{R}$, where X is an arbitrary topological space. To prove this fact, one would need to slightly adjust the proof supplied here.