



planetmath.org

Math for the people, by the people.

dual of Dilworth’s theorem

Canonical name	DualOfDilworthsTheorem
Date of creation	2013-03-22 15:01:49
Last modified on	2013-03-22 15:01:49
Owner	justice (4961)
Last modified by	justice (4961)
Numerical id	7
Author	justice (4961)
Entry type	Theorem
Classification	msc 06A06
Related topic	DilworthsTheorem

Theorem 1. *Let P be a poset of height h . Then P can be partitioned into h antichains and furthermore at least h antichains are required.*

Proof. Induction on h . If $h = 1$, then no elements of P are comparable, so P is an antichain. Now suppose that P has height $h \geq 2$ and that the theorem is true for $h - 1$. Let A_1 be the set of maximal elements of P . Then A_1 is an antichain in P and $P - A_1$ has height $h - 1$ since we have removed precisely one element from every chain. Hence, $P - A_1$ can be partitioned into $h - 1$ antichains A_2, A_3, \dots, A_h . Now we have the partition $A_1 \cup A_2 \cup \dots \cup A_h$ of P into h antichains as desired.

The necessity of h antichains is trivial by the pigeonhole principle; since P has height h , it has a chain of length h , and each element of this chain must be placed in a different antichain of our partition. \square