



planetmath.org

Math for the people, by the people.

direct limit of algebraic systems

Canonical name	DirectLimitOfAlgebraicSystems
Date of creation	2013-03-22 16:53:56
Last modified on	2013-03-22 16:53:56
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	7
Author	CWoo (3771)
Entry type	Definition
Classification	msc 08B25
Synonym	direct system of algebraic systems
Synonym	inverse system of algebraic systems
Synonym	projective system of algebraic systems
Related topic	DirectLimitOfSets
Defines	direct family of algebraic systems
Defines	inverse family of algebraic systems
Defines	inverse limit of algebraic systems

An immediate generalization of the concept of the direct limit of a direct family of sets is the direct limit of a direct family of algebraic systems.

Direct Family of Algebraic Systems

The definition is almost identical to that of a direct family of sets, except that functions ϕ_{ij} are now homomorphisms. For completeness, we will spell out the definition in its entirety.

Let $\mathcal{A} = \{A_i \mid i \in I\}$ be a family of algebraic systems of the same type (say, they are all O -algebras), indexed by a non-empty set I . \mathcal{A} is said to be a *direct family* if

1. I is a directed set,
2. whenever $i \leq j$ in I , there is a homomorphism $\phi_{ij} : A_i \rightarrow A_j$,
3. ϕ_{ii} is the identity on A_i ,
4. if $i \leq j \leq k$, then $\phi_{jk} \circ \phi_{ij} = \phi_{ik}$.

An example of this is a direct family of sets. A homomorphism between two sets is just a function between the sets.

Direct Limit of Algebraic Systems

Let \mathcal{A} be a direct family of algebraic systems A_i , indexed by I ($i \in I$). Take the disjoint union of the underlying sets of each algebraic system, and call it A . Next, a binary relation \sim is defined on A as follows:

given that $a \in A_i$ and $b \in A_j$, $a \sim b$ iff there is A_k such that $\phi_{ik}(a) = \phi_{jk}(b)$.

It is shown <http://planetmath.org/DirectLimitOfSetshere> that \sim is an equivalence relation on A , so we can take the quotient A/\sim , and denote it by A_∞ . Elements of A_∞ are denoted by $[a]_I$ or $[a]$ when there is no confusion, where $a \in A$. So A_∞ is just the direct limit of A_i *considered as sets*.

Next, we want to turn A_∞ into an O -algebra. Corresponding to each set of n -ary operations ω_i defined on A_i for all $i \in I$, we define an n -ary operation ω on A_∞ as follows:

for $i = 1, \dots, n$, pick $a_i \in A_{j(i)}$, $j(i) \in I$. Let $J := \{j(i) \mid i = 1, \dots, n\}$. Since I is directed and J is finite, J has an upper bound $j \in I$. Let $\alpha_i = \phi_{j(i)j}(a_i)$. Define

$$\omega([a_1], \dots, [a_n]) := [\omega_j(\alpha_1, \dots, \alpha_n)].$$

Proposition 1. ω is a well-defined n -ary operation on A_∞ .

Proof. Suppose $[b_1] = [a_1], \dots, [b_n] = [a_n]$. Let α_i be defined as above, and let $a := \omega_j(\alpha_1, \dots, \alpha_n) \in A_j$. Similarly, β_i are defined: $\beta_i := \phi_{k(i)k}(b_i) \in A_k$, where $b_i \in A_{k(i)}$. Let $b := \omega_k(\beta_1, \dots, \beta_n) \in A_k$. We want to show that $a \sim b$.

Since $a_i \sim b_i$, $\alpha_i \sim \beta_i$. So there is $c_i := \phi_{j\ell(i)}(\alpha_i) = \phi_{k\ell(i)}(\beta_i) \in A_{\ell(i)}$. Let ℓ be the upper bound of the set $\{\ell(1), \dots, \ell(n)\}$ and define $\gamma_i := \phi_{\ell(i)\ell}(c_i) \in A_\ell$. Then

$$\begin{aligned} \phi_{j\ell}(a) &= \phi_{j\ell}(\omega_j(\alpha_1, \dots, \alpha_n)) \\ &= \omega_\ell(\phi_{j\ell}(\alpha_1), \dots, \phi_{j\ell}(\alpha_n)) \\ &= \omega_\ell(\phi_{\ell(1)\ell} \circ \phi_{j\ell(1)}(\alpha_1), \dots, \phi_{\ell(n)\ell} \circ \phi_{j\ell(n)}(\alpha_n)) \\ &= \omega_\ell(\phi_{\ell(1)\ell}(c_1), \dots, \phi_{\ell(n)\ell}(c_n)) \\ &= \omega_\ell(\phi_{\ell(1)\ell} \circ \phi_{k\ell(1)}(\beta_1), \dots, \phi_{\ell(n)\ell} \circ \phi_{k\ell(n)}(\beta_n)) \\ &= \omega_\ell(\phi_{k\ell}(\beta_1), \dots, \phi_{k\ell}(\beta_n)) \\ &= \phi_{k\ell}(\omega_k(\beta_1, \dots, \beta_n)) \\ &= \phi_{k\ell}(b), \end{aligned}$$

which shows that $a \sim b$. □

Definition. Let \mathcal{A} be a direct family of algebraic systems of the same type (say O) indexed by I . The O -algebra A_∞ constructed above is called the *direct limit* of \mathcal{A} . A_∞ is alternatively written $\varinjlim A_i$.

Remark. Dually, one can define an *inverse family of algebraic systems*, and its inverse limit. The inverse limit of an inverse family \mathcal{A} is written A^∞ or $\varprojlim A_i$.