



planetmath.org

Math for the people, by the people.

## Jordan-Banach and Jordan-Lie algebras

Canonical name	JordanBanachAndJordanLieAlgebras
Date of creation	2013-03-22 18:14:05
Last modified on	2013-03-22 18:14:05
Owner	bci1 (20947)
Last modified by	bci1 (20947)
Numerical id	32
Author	bci1 (20947)
Entry type	Topic
Classification	msc 08A99
Classification	msc 08A05
Classification	msc 08A70
Synonym	quantum operator algebras
Related topic	Algebras2
Related topic	CAlgebra3
Related topic	AlgebraicCategoryOfLMnLogicAlgebras
Related topic	NonAbelianStructures
Related topic	AbelianCategory
Related topic	AxiomsForAnAbelianCategory
Related topic	GeneralizedVanKampenTheoremsHigherDimensional
Related topic	AxiomaticTheoryOfSupercategories
Related topic	AlgebraicCategoryOfLMnLogicAlgebras
Related topic	Categorical
Defines	Jordan algebra
Defines	Jordan-Banach algebra
Defines	Jordan-Lie algebra

### 0.0.1 Definitions of Jordan-Banach, Jordan-Lie, and Jordan-Banach-Lie algebras

Firstly, a specific *algebra* consists of a vector space  $E$  over a ground field (typically  $\mathbb{R}$  or  $\mathbb{C}$ ) equipped with a bilinear and distributive multiplication  $\circ$ . Note that  $E$  is not necessarily commutative or associative.

A *Jordan algebra* (over  $\mathbb{R}$ ), is an algebra over  $\mathbb{R}$  for which:

$$\begin{aligned} S \circ T &= T \circ S, \\ S \circ (T \circ S^2) &= (S \circ T) \circ S^2, \end{aligned}$$

for all elements  $S, T$  of the algebra.

It is worthwhile noting now that in the algebraic theory of Jordan algebras, an important role is played by the *Jordan triple product*  $\{STW\}$  as defined by:

$$\{STW\} = (S \circ T) \circ W + (T \circ W) \circ S - (S \circ W) \circ T,$$

which is linear in each factor and for which  $\{STW\} = \{WTS\}$ . Certain examples entail setting  $\{STW\} = \frac{1}{2}\{STW + WTS\}$ .

A *Jordan Lie algebra* is a real vector space  $\mathfrak{A}_{\mathbb{R}}$  together with a *Jordan product*  $\circ$  and *Poisson bracket*

$\{, \}$ , satisfying :

$$1. \text{ for all } S, T \in \mathfrak{A}_{\mathbb{R}}, \quad \begin{aligned} S \circ T &= T \circ S \\ \{S, T\} &= -\{T, S\} \end{aligned}$$

2. the *Leibniz rule* holds

$$\{S, T \circ W\} = \{S, T\} \circ W + T \circ \{S, W\} \text{ for all } S, T, W \in \mathfrak{A}_{\mathbb{R}}, \text{ along with}$$

3. the *Jacobi identity* :

$$\{S, \{T, W\}\} = \{\{S, T\}, W\} + \{T, \{S, W\}\}$$

4. for some  $\hbar^2 \in \mathbb{R}$ , there is the *associator identity* :

$$(S \circ T) \circ W - S \circ (T \circ W) = \frac{1}{4}\hbar^2\{\{S, W\}, T\}.$$

### 0.0.2 Poisson algebra

By a *Poisson algebra* we mean a Jordan algebra in which  $\circ$  is associative. The usual algebraic types of morphisms automorphism, isomorphism, etc.) apply to Jordan-Lie (Poisson) algebras (see Landsman, 2003).

Consider the classical configuration space  $Q = \mathbb{R}^3$  of a moving particle whose phase space is the cotangent bundle  $T^*\mathbb{R}^3 \cong \mathbb{R}^6$ , and for which the space of (classical) observables is taken to be the real vector space of smooth functions

$$\mathfrak{A}_{\mathbb{R}}^0 = C^\infty(T^*R^3, \mathbb{R})$$

. The usual pointwise multiplication of functions  $fg$  defines a bilinear map on  $\mathfrak{A}_{\mathbb{R}}^0$ , which is seen to be commutative and associative. Further, the Poisson bracket on functions

$$\{f, g\} := \frac{\partial f}{\partial p^i} \frac{\partial g}{\partial q_i} - \frac{\partial f}{\partial q_i} \frac{\partial g}{\partial p^i} ,$$

which can be easily seen to satisfy the Liebniz rule above. The axioms above then set the stage of passage to quantum mechanical systems which the parameter  $k^2$  suggests.

### 0.0.3 C\*-algebras (C\*-A), JLB and JBW Algebras

An *involution* on a complex algebra  $\mathfrak{A}$  is a real-linear map  $T \mapsto T^*$  such that for all

$$S, T \in \mathfrak{A} \text{ and } \lambda \in \mathbb{C}, \text{ we have } T^{**} = T, (ST)^* = T^*S^*, (\lambda T)^* = \bar{\lambda}T^* .$$

A *\*-algebra* is said to be a complex associative algebra together with an involution  $*$  .

A *C\*-algebra* is a simultaneously a \*-algebra and a Banach space  $\mathfrak{A}$ , satisfying for all  $S, T \in \mathfrak{A}$  :

$$\begin{aligned} \|S \circ T\| &\leq \|S\| \|T\| , \\ \|T^*T\|^2 &= \|T\|^2 . \end{aligned}$$

We can easily see that  $\|A^*\| = \|A\|$  . By the above axioms a C\*-algebra is a special case of a Banach algebra where the latter requires the above norm property but not the involution (\*) property. Given Banach spaces  $E, F$  the space  $\mathcal{L}(E, F)$  of (bounded) linear operators from  $E$  to  $F$  forms a Banach space, where for  $E = F$ , the space  $\mathcal{L}(E) = \mathcal{L}(E, E)$  is a Banach algebra with respect to the norm

$$\|T\| := \sup\{\|Tu\| : u \in E, \|u\| = 1\} .$$

In quantum field theory one may start with a Hilbert space  $H$ , and consider the Banach algebra of bounded linear operators  $\mathcal{L}(H)$  which given to be closed under the usual algebraic operations and taking adjoints, forms a \*-algebra of bounded operators, where the adjoint operation functions as the involution, and for  $T \in \mathcal{L}(H)$  we have :

$$\|T\| := \sup\{(Tu, Tu) : u \in H, (u, u) = 1\} , \text{ and } \|Tu\|^2 = (Tu, Tu) = (u, T^*Tu) \leq \|T^*T\| \|u\|^2 .$$

By a morphism between C\*-algebras  $\mathfrak{A}, \mathfrak{B}$  we mean a linear map  $\phi : \mathfrak{A} \rightarrow \mathfrak{B}$ , such that for all  $S, T \in \mathfrak{A}$ , the following hold :

$$\phi(ST) = \phi(S)\phi(T) , \phi(T^*) = \phi(T)^* ,$$

where a bijective morphism is said to be an isomorphism (in which case it is then an isometry). A fundamental relation is that any norm-closed  $*$ -algebra  $\mathcal{A}$  in  $\mathcal{L}(H)$  is a  $C^*$ -algebra, and conversely, any  $C^*$ -algebra is isomorphic to a norm-closed  $*$ -algebra in  $\mathcal{L}(H)$  for some Hilbert space  $H$ .

For a  $C^*$ -algebra  $\mathfrak{A}$ , we say that  $T \in \mathfrak{A}$  is *self-adjoint* if  $T = T^*$ . Accordingly, the self-adjoint part  $\mathfrak{A}^{sa}$  of  $\mathfrak{A}$  is a real vector space since we can decompose  $T \in \mathfrak{A}^{sa}$  as :

$$T = T' + T'' := \frac{1}{2}(T + T^*) + \iota\left(\frac{-\iota}{2}\right)(T - T^*) .$$

A *commutative*  $C^*$ -algebra is one for which the associative multiplication is commutative. Given a commutative  $C^*$ -algebra  $\mathfrak{A}$ , we have  $\mathfrak{A} \cong C(Y)$ , the algebra of continuous functions on a compact Hausdorff space  $Y$ .

A *Jordan-Banach algebra* (a JB-algebra for short) is both a real Jordan algebra and a Banach space, where for all  $S, T \in \mathfrak{A}_{\mathbb{R}}$ , we have

$$\begin{aligned} \|S \circ T\| &\leq \|S\| \|T\| , \\ \|T\|^2 &\leq \|S^2 + T^2\| . \end{aligned}$$

A *JLB-algebra* is a JB-algebra  $\mathfrak{A}_{\mathbb{R}}$  together with a Poisson bracket for which it becomes a Jordan-Lie algebra for some  $\hbar^2 \geq 0$ . Such JLB-algebras often constitute the real part of several widely studied complex associative algebras.

For the purpose of quantization, there are fundamental relations between  $\mathfrak{A}^{sa}$ , JLB and Poisson algebras.

For further details see Landsman (2003) (Thm. 1.1.9).

A JB-algebra which is monotone complete and admits a separating set of normal sets is called a *JBW-algebra*. These appeared in the work of von Neumann who developed a (orthomodular) lattice theory of projections on  $\mathcal{L}(H)$  on which to study quantum logic (see later). BW-algebras have the following property: whereas  $\mathfrak{A}^{sa}$  is a J(L)B-algebra, the self adjoint part of a von Neumann algebra is a JBW-algebra.

A *JC-algebra* is a norm closed real linear subspace of  $\mathcal{L}(H)^{sa}$  which is closed under the bilinear product  $S \circ T = \frac{1}{2}(ST + TS)$  (non-commutative and nonassociative). Since any norm closed Jordan subalgebra of  $\mathcal{L}(H)^{sa}$  is a JB-algebra, it is natural to specify the exact relationship between JB and JC-algebras, at least in finite dimensions. In order to do this, one introduces the ‘exceptional’ algebra  $H_3(\mathbb{O})$ , the algebra of  $3 \times 3$  Hermitian matrices with values in the octonions  $\mathbb{O}$ . Then a finite dimensional JB-algebra is a JC-algebra if and only if it does not contain  $H_3(\mathbb{O})$  as a (direct) summand [?].

The above definitions and constructions follow the approach of Alfsen and Schultz (2003) and Landsman (1998).

## References

- [1] Alfsen, E.M. and F. W. Schultz: Geometry of State Spaces of Operator Algebras, Birkhäuser, Boston-Basel-Berlin.(2003).