

planetmath.org

Math for the people, by the people.

implicational class

Canonical name ImplicationalClass
Date of creation 2013-03-22 17:31:35
Last modified on 2013-03-22 17:31:35

Owner CWoo (3771) Last modified by CWoo (3771)

Numerical id 5

Author CWoo (3771)
Entry type Definition
Classification msc 08C15
Classification msc 03C05
Synonym quasivariety

Synonym quasiprimitive class
Defines algebraic class

Defines equational implication

In this entry, we extend the notion of an equational class (or a variety) to a more general notion known as an implicational class (or a quasivariety). Recall that an equational class K is a class of algebraic systems satisfying a set Σ of "equations" and that K is the smallest class satisfying Σ . Typical examples are the varieties of groups, rings, or lattices.

An implicational class, loosely speaking, is the *smallest* class of algebraic systems satisfying a set of "implications", where an implication has the form $P \to Q$, where P and Q are some sentences. Formally, we define an *equational implication* in an algebraic system to be a sentence of the form

$$(\forall x_1)\cdots(\forall x_n)(e_1\wedge\cdots\wedge e_p\to e_q),$$

where each e_i is an identity of the form $f_i(x_1, \ldots, x_n) = g_i(x_1, \ldots, x_n)$ for some *n*-ary polynomials f_i and g_i , and $i = 1, \ldots, p, q$.

Definition. A class K of algebraic systems of the same type (signature) is called an *implicational class* if there is a set Σ of equational implications such that

$$K = \{A \text{ is a structure } \mid A \text{ is a model in } \Sigma\} = \{A \mid (\forall q \in \Sigma) \to (A \models q)\}.$$

Examples

- 1. Any equational class is implicational. Each identity p=q can be thought of as an equational implication $(p=p) \rightarrow (p=q)$. In other words, every algebra satisfying the identity also satisfies the corresponding equational implication, and vice versa.
- 2. The class of all Dedekind-finite rings. In addition to satisfying the identities for being a (unital) ring, each ring also satisfies the equational implication

$$(\forall x)(\forall y)(xy=1) \to (yx=1).$$

3. The class of all cancellation semigroups. In addition to satisfying the identities for being a semigroup, each semigroup also satisfies the implications

$$(\forall x)(\forall y)(\forall z)(xy=xz)\to (y=z)\quad \text{and}\quad (\forall x)(\forall y)(\forall z)(yx=zx)\to (y=z).$$

4. The class K of all torsion free abelian groups. In addition to satisfying the identities for being abelian groups, each group also satisfies the set of all implications

$$\{\forall x(nx=0) \to (x=0) \mid n \text{ is a positive integer}\}.$$

There is an equivalent formulation of an implicational class. Again, let K be a class of algebraic systems of the same type (signature) τ . Define the following four "operations" on the classes of algebraic systems of type τ :

- 1. I(K) is the class of all isomorphic copies of algebras in K,
- 2. S(K) is the class of all subalgebras of algebras in K,
- 3. P(K) is the class of all product of algebras in K (including the empty products, which means P(K) includes the trivial algebra), and
- 4. U(K) is the class of all ultraproducts of algebras in K.

Suppose X is any one of the operations above, we say that K is *closed* under operation X if $X(K) \subseteq K$.

Definition. K is said to be an algebraic class if K is closed under I, and K is said to be a quasivariety if it is algebraic and is closed under S, P, U.

It can be shown that a class K of algebraic systems of the same type is implicational iff it is a quasivariety. Therefore, we may use the two terms interchangeably.

As we have seen earlier, a variety is a quasivariety. However, the converse is not true, as can be readily seen in the last example above, since a homomorphic image of a torsion free abelian is in general not torsion free: the homomorphic image of $\phi: \mathbb{Z} \to \mathbb{Z}_n$ is a subgroup of \mathbb{Z}_n , hence not torsion free.