



planetmath.org

Math for the people, by the people.

congruence relation on an algebraic system

Canonical name	CongruenceRelationOnAnAlgebraicSystem
Date of creation	2013-03-22 16:26:23
Last modified on	2013-03-22 16:26:23
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	33
Author	CWoo (3771)
Entry type	Definition
Classification	msc 08A30
Related topic	Congruence3
Related topic	Congruence2
Related topic	CongruenceInAlgebraicNumberField
Related topic	PolynomialCongruence
Related topic	QuotientCategory
Related topic	CategoryOfAdditiveFractions
Defines	congruence
Defines	congruence relation
Defines	quotient algebra
Defines	proper congruence
Defines	trivial congruence
Defines	non-trivial congruence
Defines	congruence restricted to a subalgebra
Defines	extension of a subalgebra by a congruence
Defines	principal congruence
Defines	congruence generated by

Let (A, O) be an algebraic system. A *congruence relation*, or simply a *congruence* \mathfrak{C} on A

1. is an equivalence relation on A ; if $(a, b) \in \mathfrak{C}$ we write $a \equiv b \pmod{\mathfrak{C}}$, and
2. respects every n -ary operator on A : if ω_A is an n -ary operator on A ($\omega \in O$), and for any $a_i, b_i \in A$, $i = 1, \dots, n$, we have

$$a_i \equiv b_i \pmod{\mathfrak{C}} \quad \text{implies} \quad \omega_A(a_1, \dots, a_n) \equiv \omega_A(b_1, \dots, b_n) \pmod{\mathfrak{C}}.$$

For example, A^2 and $\Delta_A := \{(a, a) \mid a \in A\}$ are both congruence relations on A . Δ_A is called the *trivial congruence* (on A). A *proper* congruence relation is one not equal to A^2 .

Remarks.

- \mathfrak{C} is a congruence relation on A if and only if \mathfrak{C} is an equivalence relation on A and a subalgebra of the <http://planetmath.org/DirectProductOfAlgebrasproduct> $A \times A$.
- The set of congruences of an algebraic system is a complete lattice. The meet is the usual set intersection. The join (of an arbitrary number of congruences) is the <http://planetmath.org/PartitionsFormALatticejoin> of the underlying equivalence relations. This join corresponds to the subalgebra (of $A \times A$) generated by the union of the underlying sets of the congruences. The *lattice of congruences* on A is denoted by $\text{Con}(A)$.
- **(restriction)** If \mathfrak{C} is a congruence on A and B is a subalgebra of A , then \mathfrak{C}_B defined by $\mathfrak{C} \cap (B \times B)$ is a congruence on B . The equivalence of \mathfrak{C}_B is obvious. For any n -ary operator ω_B inherited from A 's ω_A , if $a_i \equiv b_i \pmod{\mathfrak{C}_B}$, then $\omega_B(a_1, \dots, a_n) = \omega_A(a_1, \dots, a_n) \equiv \omega_A(b_1, \dots, b_n) = \omega_B(b_1, \dots, b_n) \pmod{\mathfrak{C}}$. Since both $\omega_B(a_1, \dots, a_n)$ and $\omega_B(b_1, \dots, b_n)$ are in B , $\omega_B(a_1, \dots, a_n) \equiv \omega_B(b_1, \dots, b_n) \pmod{\mathfrak{C}_B}$ as well. \mathfrak{C}_B is the congruence *restricted* to B .
- **(extension)** Again, let \mathfrak{C} be a congruence on A and B a subalgebra of A . Define $B^\mathfrak{C}$ by $\{a \in A \mid (a, b) \in \mathfrak{C} \text{ and } b \in B\}$. In other words, $a \in B^\mathfrak{C}$ iff $a \equiv b \pmod{\mathfrak{C}}$ for some $b \in B$. We assert that $B^\mathfrak{C}$ is a subalgebra of A . If ω_A is an n -ary operator on A and $a_1, \dots, a_n \in$

$B^{\mathfrak{C}}$, then $a_i \equiv b_i \pmod{\mathfrak{C}}$, so $\omega_A(a_1, \dots, a_n) \equiv \omega_A(b_1, \dots, b_n) \pmod{\mathfrak{C}}$. Since $\omega_A(b_1, \dots, b_n) \in B$, $\omega_A(a_1, \dots, a_n) \in B^{\mathfrak{C}}$. Therefore, $B^{\mathfrak{C}}$ is a subalgebra. Because $B \subseteq B^{\mathfrak{C}}$, we call it the *extension* of B by \mathfrak{C} .

- Let B be a subset of $A \times A$. The smallest congruence \mathfrak{C} on A such that $a \equiv b \pmod{\mathfrak{C}}$ for all $a, b \in B$ is called the *congruence generated by B* . \mathfrak{C} is often written $\langle B \rangle$. When B is a singleton $\{(a, b)\}$, then we call $\langle B \rangle$ a *principal congruence*, and denote it by $\langle (a, b) \rangle$.

Quotient algebra

Given an algebraic structure (A, O) and a congruence relation \mathfrak{C} on A , we can construct a new O -algebra $(A/\mathfrak{C}, O)$, as follows: elements of A/\mathfrak{C} are of the form $[a]$, where $a \in A$. We set

$$[a] = [b] \text{ iff } a \equiv b \pmod{\mathfrak{C}}.$$

Furthermore, for each n -ary operator ω_A on A , define $\omega_{A/\mathfrak{C}}$ by

$$\omega_{A/\mathfrak{C}}([a_1], \dots, [a_n]) := [\omega_A(a_1, \dots, a_n)].$$

It is easy to see that $\omega_{A/\mathfrak{C}}$ is a well-defined operator on A/\mathfrak{C} . The O -algebra thus constructed is called the *quotient algebra* of A over \mathfrak{C} .

Remark. The bracket $[\cdot] : A \rightarrow A/\mathfrak{C}$ is in fact an epimorphism, with <http://planetmath.org/KernelOfAHomomorphismBetweenAlgebraicSystems> kernel $([\cdot]) = \mathfrak{C}$. This means that every congruence of an algebraic system A is the kernel of some homomorphism from A . $[\cdot]$ is usually written $[\cdot]_{\mathfrak{C}}$ to signify its association with \mathfrak{C} .

References

- [1] G. Grätzer: *Universal Algebra*, 2nd Edition, Springer, New York (1978).