

planetmath.org

Math for the people, by the people.

de Moivre identity, proof of

Canonical name DeMoivreIdentityProofOf

Date of creation 2013-03-22 14:34:08 Last modified on 2013-03-22 14:34:08

Owner rspuzio (6075) Last modified by rspuzio (6075)

Numerical id 10

Author rspuzio (6075)

Entry type Proof Classification msc 12E10

Synonym proof of de Moivre's formula Synonym proof of de Moivre's theorem

Related topic AngleSumIdentity

To prove the de Moivre identity, we will first prove by induction on n that the identity holds for all natural numbers.

For the case n = 0, observe that

$$\cos(0\theta) + i\sin(0\theta) = 1 + i0 = (\cos(\theta) + i\sin(\theta))^{0}.$$

Assume that the identity holds for a certain value of n:

$$\cos(n\theta) + i\sin(n\theta) = (\cos(\theta) + i\sin(\theta))^{n}.$$

Multiply both sides of this identity by $\cos(\theta) + i\sin(\theta)$ and expand the left side to obtain

$$\cos(\theta)\cos(n\theta) - \sin(\theta)\sin(n\theta) + i\cos(\theta)\sin(n\theta) + i\sin(\theta)\cos(n\theta) = (\cos(\theta) + i\sin(\theta))(\cos(n\theta) + i\sin(\theta))\sin(n\theta) + i\sin(\theta)\sin(n\theta) + i\sin(\theta)\sin(n\theta) + i\sin(\theta)\cos(n\theta) = (\cos(\theta) + i\sin(\theta))(\cos(n\theta) + i\sin(\theta))\sin(n\theta) + i\sin(\theta)\sin(n\theta) + i\sin(\theta)\cos(n\theta) = (\cos(\theta) + i\sin(\theta))(\cos(n\theta) + i\sin(\theta))\sin(n\theta) + i\sin(\theta)\cos(n\theta) = (\cos(\theta) + i\sin(\theta))\cos(n\theta) = (\cos(\theta) + i\sin(\theta))\cos(n\theta) + i\sin(\theta)\cos(n\theta) = (\cos(\theta) + i\sin(\theta))\cos(n\theta) = (\cos(\theta) + i\cos(\theta))\cos(n\theta) = (\cos(\theta) + i\cos($$

By the angle sum identities,

$$\cos(\theta)\cos(n\theta) - \sin(\theta)\sin(n\theta) = \cos(n\theta + \theta)$$
$$\cos(\theta)\sin(n\theta) + \sin(\theta)\cos(n\theta) = \sin(n\theta + \theta)$$

Therefore,

$$\cos((n+1)\theta) + i\sin((n+1)\theta) = (\cos(\theta) + i\sin(\theta))^{n+1}.$$

Hence by induction de Moivre's identity holds for all natural n.

Now let -n be any negative integer. Then using the fact that cos is an even and sin an odd function, we obtain that

$$\cos(-n\theta) + i\sin(-n\theta) = \cos(n\theta) - i\sin(n\theta)$$

$$= \frac{\cos(n\theta) - i\sin(n\theta)}{\cos^2(n\theta) + \sin^2(n\theta)}$$

$$= \frac{1}{\cos(n\theta) + i\sin(n\theta)} \cdot \frac{\cos(n\theta) - i\sin(n\theta)}{\cos(n\theta) - i\sin(n\theta)}$$

$$= \frac{1}{\cos(n\theta) + i\sin(n\theta)},$$

the denominator of which is $(\cos(n\theta) + i\sin(n\theta))^n$. Hence

$$\cos(-n\theta) + i\sin(-n\theta) = (\cos(\theta) + i\sin(\theta))^{-n}.$$