



planetmath.org

Math for the people, by the people.

characterizations of integral

Canonical name	CharacterizationsOfIntegral
Date of creation	2013-03-22 14:56:54
Last modified on	2013-03-22 14:56:54
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	11
Author	pahio (2872)
Entry type	Theorem
Classification	msc 12E99
Classification	msc 13B21
Synonym	characterisations of integral

Theorem. Let R be a subring of a field K , $1 \in R$ and let α be a non-zero element of K . The following conditions are equivalent:

1. α is integral over R .
2. α belongs to $R[\alpha^{-1}]$.
3. α is unit of $R[\alpha^{-1}]$.
4. $\alpha^{-1}R[\alpha^{-1}] = R[\alpha^{-1}]$.

Proof. Supposing the first condition that an equation

$$\alpha^n + a_1\alpha^{n-1} + \dots + a_{n-1}\alpha + a_n = 0,$$

with a_j 's belonging to R , holds. Dividing both by α^{n-1} gives

$$\alpha = -a_1 - a_2\alpha^{-1} - \dots - a_n\alpha^{-n+1}.$$

One sees that α belongs to the ring $R[\alpha^{-1}]$ even being a unit of this (of course $\alpha^{-1} \in R[\alpha^{-1}]$). Therefore also the principal ideal $\alpha^{-1}R[\alpha^{-1}]$ of the ring $R[\alpha^{-1}]$ coincides with this ring. Conversely, the last circumstance implies that α is integral over R .

References

- [1] Emil Artin: . Lecture notes. Mathematisches Institut, Göttingen (1959).