



Math for the people, by the people.

variant of Cardano's derivation

Canonical name	VariantOfCardanosDerivation
Date of creation	2013-03-22 13:38:43
Last modified on	2013-03-22 13:38:43
Owner	mathcam (2727)
Last modified by	mathcam (2727)
Numerical id	9
Author	mathcam (2727)
Entry type	Proof
Classification	msc 12D10

By a linear change of variable, a cubic polynomial over \mathbb{C} can be given the form $x^3 + 3bx + c$. To find the zeros of this cubic in the form of surds in b and c , make the substitution $x = y^{1/3} + z^{1/3}$, thus replacing one unknown with two, and then write down identities which are suggested by the resulting equation in two unknowns. Specifically, we get

$$y + 3(y^{1/3} + z^{1/3})y^{1/3}z^{1/3} + z + 3b(y^{1/3} + z^{1/3}) + c = 0. \quad (1)$$

This will be true if

$$y + z + c = 0 \quad (2)$$

$$3y^{1/3}z^{1/3} + 3b = 0, \quad (3)$$

which in turn requires

$$yz = -b^3. \quad (4)$$

The pair of equations (2) and (4) is a quadratic system in y and z , readily solved. But notice that (3) puts a restriction on a certain choice of cube roots.