



planetmath.org

Math for the people, by the people.

proof of factor theorem using division

Canonical name	ProofOfFactorTheoremUsingDivision
Date of creation	2013-03-22 15:08:58
Last modified on	2013-03-22 15:08:58
Owner	alozano (2414)
Last modified by	alozano (2414)
Numerical id	8
Author	alozano (2414)
Entry type	Proof
Classification	msc 12D10
Classification	msc 12D05

Lemma (cf. factor theorem). *Let R be a commutative ring with identity and let $p(x) \in R[x]$ be a polynomial with coefficients in R . The element $a \in R$ is a root of $p(x)$ if and only if $(x - a)$ divides $p(x)$.*

Proof. Let $p(x)$ be a polynomial in $R[x]$ and let a be an element of R .

1. First we assume that $(x - a)$ divides $p(x)$. Therefore, there is a polynomial $q(x) \in R[x]$ such that $p(x) = (x - a) \cdot q(x)$. Hence, $p(a) = (a - a) \cdot q(a) = 0$ and a is a root of $p(x)$.
2. Assume that a is a root of $p(x)$, i.e. $p(a) = 0$. Since $x - a$ is a monic polynomial, we can perform the <http://planetmath.org/LongDivisionpolynomial> long division of $p(x)$ by $(x - a)$. Thus, there exist polynomials $q(x)$ and $r(x)$ such that:

$$p(x) = (x - a) \cdot q(x) + r(x)$$

and the degree of $r(x)$ is less than the degree of $x - a$ (so $r(x)$ is just a constant). Moreover, $0 = p(a) = 0 + r(a) = r(a) = r(x)$. Therefore $p(x) = (x - a) \cdot q(x)$ and $(x - a)$ divides $p(x)$.

□