



planetmath.org

Math for the people, by the people.

multiplication rule gives inverse ideal

Canonical name	MultiplicationRuleGivesInverseIdeal
Date of creation	2013-03-22 15:24:16
Last modified on	2013-03-22 15:24:16
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	5
Author	pahio (2872)
Entry type	Theorem
Classification	msc 13A15
Classification	msc 16D25
Related topic	PruferRing
Related topic	Characterization

Theorem. Let R be a commutative ring with non-zero unity. If an ideal (a, b) of R , with a or b <http://planetmath.org/RegularElementregular>, obeys the multiplication rule

$$(a, b)(c, d) = (ac, ad+bc, bd) \quad (1)$$

with all ideals (c, d) of R , then (a, b) is an invertible ideal.

Proof. The rule gives

$$(a, b)^2 = (a, -b)(a, b) = (a^2, ab-ba, b^2) = (a^2, b^2).$$

Thus the product ab may be written in the form

$$ab = ua^2 + vb^2,$$

where u and v are elements of R . Let's assume that e.g. a is regular. Then a has the multiplicative inverse a^{-1} in the total ring of fractions R . Again applying the rule yields

$$(a, b)(va, a-vb)(a^{-2}) = (va^2, a^2-vab+vab, ab-vb^2)(a^{-2}) = (va^2, a^2, ua^2)(a^{-2}) = (v, 1, u) = R.$$

Consequently the ideal (a, b) has an inverse ideal (which may be a <http://planetmath.org/FractionalIdeal> ideal); this settles the proof.

Remark. The rule (1) in the theorem may be replaced with the rule

$$(a, b)(c, d) = (ac, (a+b)(c+d), bd) \quad (2)$$

as is seen from the identical equation $(a+b)(c+d) - ac - bd = ad + bc$.