



Math for the people, by the people.

arithmetic ring

Canonical name	ArithmeticalRing
Date of creation	2013-03-22 15:23:58
Last modified on	2013-03-22 15:23:58
Owner	PrimeFan (13766)
Last modified by	PrimeFan (13766)
Numerical id	8
Author	PrimeFan (13766)
Entry type	Theorem
Classification	msc 13A99
Related topic	QuotientOfIdeals

Theorem. If R is a commutative ring, then the following three conditions are equivalent:

- For all ideals \mathfrak{a} , \mathfrak{b} and \mathfrak{c} of R , one has $\mathfrak{a} \cap (\mathfrak{b} + \mathfrak{c}) = (\mathfrak{a} \cap \mathfrak{b}) + (\mathfrak{a} \cap \mathfrak{c})$.
- For all ideals \mathfrak{a} , \mathfrak{b} and \mathfrak{c} of R , one has $\mathfrak{a} + (\mathfrak{b} \cap \mathfrak{c}) = (\mathfrak{a} + \mathfrak{b}) \cap (\mathfrak{a} + \mathfrak{c})$.
- For each maximal ideal \mathfrak{p} of R the set of all ideals of $R_{\mathfrak{p}}$, the <http://planetmath.org/Localization> of R at $R \setminus \mathfrak{p}$, is totally ordered by set inclusion.

The ring R satisfying the conditions of the theorem is called an *arithmetical ring*.